FERNANDA ANTUNES ALVES DA COSTA

Isolamento e caracterização de DETs (*Differentially Expressed Transcripts*) em espécies de peixes: *Leporinus macrocephalus* (Characiformes) e *Danio rerio* (Cypriniformes)

Tese apresentada ao Instituto de Biociências da Universidade Estadual Paulista "Júlio de Mesquita Filho", Campus de Botucatu, para obtenção do título de Doutor em Ciências Biológicas - Genética

Orientadora: Adriane Pinto Wasko

Botucatu - SP 2008

Livros Grátis

http://www.livrosgratis.com.br

Milhares de livros grátis para download.

FICHA CATALOGRÁFICA ELABORADA PELA SEÇÃO DE AQUIS. E TRAT. DA INFORMAÇÃO DIVISÃO TÉCNICA DE BIBLIOTECA E DOCUMENTAÇÃO - CAMPUS DE BOTUCATU - UNESP BIBLIOTECÁRIA RESPONSÁVEL: *ROSEMEIRE APARECIDA VICENTE*

BIBLIOTECARIA RESPONSAVEL. ROSEMEIRE AFARECIDA VICENTE
Costa, Fernanda Antunes Alves da.
Isolamento e caracterização de DETs (Differentially Expressed
Transcripts) em espécies de peixes: Leporinus macrocephalus
(Characiformes) e Danio rerio (Cypriniformes) / Fernanda Antunes
Alves da Costa. – Botucatu : [s.n.], 2008
Tese (doutorado) - Instituto de Biociências de Botucatu, Universidade
Estadual Paulista, 2008.
Orientador: Prof ^a . Dr ^a Adriane Pinto Wasko
Assunto CAPES: 20200005
1. Genética. 2. Peixes. 3. Actina.
CDD 597.55
Palavras-chave: Actina: Danio rerio: DETs: Fabp6: Leporinus

Palavras-chave: Actina; Danio rerio; DETS; Fabpo; Leporinus macrocephalus.

... Quando paramos de aprender e de progredir, começamos a morrer realmente...

C. Torres Pastorino

Aos meus pais, José Maria e Cecília e ao meu esposo, Rogério, que me amam e me compreendem, sempre me apoiando em todos os grandes e pequenos passos do nosso caminho.

À minha querida orientadora, Adriane, exemplo de pessoa e profissional.

AGRADECIMENTOS

Agradeço a todos que contribuíram direta ou indiretamente à realização deste trabalho, em especial:

À minha orientadora, Profa. Dra. Adriane Pinto Wasko, pela confiança, orientação e dedicação oferecidas para elaboração e conclusão deste trabalho e para minha formação profissional;

À FAPESP – Fundação de Amparo à Pesquisa do Estado de São Paulo, pelo apoio financeiro concedido;

Aos Profs. Drs. Cesar Martins e Maeli Dal Pai Silva pelo apoio direto na realização deste trabalho;

Aos Profs. Drs. Fausto Foresti, Claudio Oliveira, Ivan de Godoy Maia, Lígia Sousa Lima Silveira da Mota, João Pessoa, Paulo Eduardo Ribolla, Alexandre Secorun Borges, pelo apoio e sugestões;

À Dra. Debora Colombi, pela preciosas sugestões quanto às análises de PCR quantitativo;

Aos Laboratórios de Biologia Molecular Animal (Departamento de Genética), Biotecnologia e Genética Molecular – BIOGEM (Departamento de Genética) e Biologia e Genética de Peixes (Departamento de Morfologia). À Pós-graduação em Ciências Biológicas (Genética), ao Instituto de Biociências e à Universidade Estadual Paulista, pela oportunidade cedida para a realização deste trabalho;

Aos companheiros do Laboratório de Biologia Molecular Animal: Magali, Cassiane, Ana Teresa, Thaís, Juliana e alunos do Prof. Dr. Alexandre Secorun Borges – Fernanda, Peres, Paulo e Zé Filho;

Aos companheiros do Laboratório de Biologia e Genética de Peixes: Alex, Andréia Alves, Andréia Poletto, Celso, Claudinha, Cristiane, Daniela, Danillo, Emanuel, Guilherme (Kabelo), Irani, Jefferson, Juliana, Karina, Kelly, Lessandra, Lígia, Luciana, Luiz Henrique, Márcio, Marina, Marisa, Patrícia Elda, Ricardo, Tatiane e Welcy

Aos alunos da Profa. Maeli Dal Pai Silva, em especial: Eduardo, Fernanda Losi e Fernanda Carani, pela enorme ajuda;

Aos alunos do Laboratório de Biotecnologia e Genética Molecular – BIOGEM, em especial ao Flávio (Bonsai), Fábio, Tânia e Virgínia.

À colega Magali, pela amizade, carinho e compreensão e por estar sempre presente e disposta a me ajudar ou simplesmente para dizer "eu entendo", coisas que somente os amigos sabem o verdadeiro significado;

Ao Prof. Dr. Luis Fernando Marins e ao Sr. Odval Stradioti, Piscicultura Água Milagrosa, pelo gentil fornecimento dos animais;

Ao Prof. Dr. Luis Fernando Marins e seus alunos, pelo apoio e discussões e pela acolhida;

Aos Profs. Drs. Jonathan M. Wright e Eileen Denovan-Wright e Stephen Mockford pela grande oportunidade oferecida, pelos muitos ensinamentos e, em especial, pelo carinho e atenção com que me receberam;

À Profa. Dra. Silvia Regina Rogatto e seus alunos, em especial à Eliane e Grace, pelo auxílio no desenvolvimento das hibridações fluorescentes *in situ;*

Aos funcionários dos Departamentos de Genética e Morfologia pelo auxílio atribuído durante a realização deste trabalho;

Aos amigos e familiares pela presença e apoio durante os vários momentos de minha vida;

Aos meus pais pelo carinho e apoio constantes, por me ensinarem sobre a vida e a viver e por toda dedicação incondicional à minha felicidade;

Ao meu amado esposo, Rogério, por completar minha felicidade, por me apoiar em todos os momentos e em todas minha decisões, por sempre buscar o meu bem e por me amar como eu sou.

Muito obrigada!!

SUMÁRIO

RESUMO	01
ABSTRACT	03
CONSIDERAÇÕES INICIAIS E OBJETIVOS	05
INTRODUÇÃO GERAL	08
Transcritos diferencialmente expressos	08
A espécie Leporinus macrocephalus	12
A espécie <i>Danio rerio</i>	15
Referências	17
RESULTADOS E DISCUSSÃO	
CAPÍTULO 1	31
Molecular organization and expression analysis of skeletal	actin genes from
white and red fish muscle tissues: insights into the musc	culature energetic
demands	
Abstract	
Introduction	
Material and Methods	
Results and Discussion	42
Acknowledgements	53
References	53
Table, Figures and Legends	60
CAPÍTULO 2	68
Identification of sexually dimorphic gene expression in brain	n tissue of the fish
Leporinus macrocephalus through mRNA differential disp	
PCR analyses	play and real time
	lay and real time 69
Abstract	blay and real time 69 69

Introduction70
Material and Methods73
Results and Discussion77
Acknowledgements
References
Table, Figures and Legends97
CAPÍTULO 3
Spatio-temporal distribution of fatty acid binding protein 6 (fabp6) gene
transcripts in the developing and adult zebrafish (Danio rerio)106
Abstract
Introduction107
Material and Methods110
Results and Discussion112
Acknowledgements120
References
Table, Figures and Legends124
CAPÍTULO 4
A comparative expression analysis of gene transcripts in brain tissue of
wild-type and GH-transgenic zebrafish (Danio rerio) using DDRT-PCR
approach133
Abstract
Introduction134
Material and Methods136
Results and Discussion139
Acknowledgements146
References
Table, Figures and Legends
CONSIDERAÇÕES FINAIS E CONCLUSÕES161
SÚMULA CURRICULAR

RESUMO

O presente estudo teve o intuito de isolar e caracterizar transcritos diferencialmente expressos em espécies de peixes, visando contribuir com novos dados genéticos acerca deste grupo basal de vertebrados. Desta forma, diferentes análises foram realizadas em Leporinus macrocephalus (piauçu) (Characiformes, Anostomidae), espécie de peixe que representa um importante recurso alimentar que tem sido intensamente explorado na aqüicultura subtropical e que apresenta um sistema de determinação sexual ZZ/ZW, e também em Danio rerio (zebrafish) (Cypriniformes, Cyprinidae), um organismo modelo já bem estabelecido em diferentes áreas da pesquisa científica. Os resultados indicaram que: (1) As α-actinas esqueléticas de tecido muscular branco e vermelho parecem ser genética e funcionalmente distintas não somente em L. macrocephalus como também em outras espécies de peixes, característica que, até o momento, não foi encontrada em outro grupo de vertebrados. Além disso, as características de cada tipo de actina podem ser relacionadas às demandas energéticas dos tecidos musculares branco e vermelho; (2) Um expressão dimórfica entre machos e fêmeas ocorre em tecido cerebral de L. macrocephalus, relacionada ao gene que codifica a proteína APBA2 (amyloid beta (A4) precursor protein-binding, family A, *member 2*), que está associada à produção da proteína β -amilóide a qual, por sua vez, é relacionada à doença de Alzheimer; (3) A estrutura do gene fabp6 (fatty acid-binding protein 6) de D. rerio é similar à maioria dos outros membros da família multigênica fabp de outros vertebrados, e sua distribuição tecidual em embriões, larvas e adultos da espécie também é similar à encontrada em outros organismos, especialmente mamíferos; (4) Os genes que codificam a proteína p300 (E1A binding protein p300) e a proteína ADCY2 (adenilato ciclase 2) representam dois genes candidatos diferencialmente expressos entre amostras não-transgêncas e homozigotas e hemizigotas transgênicas para o gene do hormônio do crescimento (GH), reforcando dados prévios que evidenciam que estas proteínas estão envolvidas na formação/crescimento muscular e liberação de GH. Desta forma, os dados obtidos contribuem para a compreensão da evolução, função e organização molecular de diversos genes e podem servir de base a futuros estudos. Além disso, os resultados obtidos confirmam a utilidade de Danio rerio como espécie modelo em estudos genéticos e indicam que

Leporinus macrocephalus representa um potencial modelo experimental em análises direcionadas a genes envolvidos no desenvolvimento de doenças humanas, como alterações neurodegenerativas, distrofias musculares e câncer.

ABSTRACT

The present study intent to isolate and characterize differentially expressed transcripts in fish species in order to contribute to new genetic data on this basal vertebrate group. Therefore, different analyses were performed in Leporinus macrocephalus (piauçu) (Characiformes, Anostomidae), a fish species that represents an important food resource that has been intensively exploited in the subtropical aquaculture and that presents a ZZ/ZW sex determination system, and also in Danio rerio (zebrafish) (Cypriniformes, Cyprinidae), a well-established model organism in different areas of scientific research. The results indicated that: (1) The α -skeletal actins from white and red muscle tissues appear to be somewhat genetically and functionally distinguishable not only in L. macrocephalus but also in other fish species, a feature that was not found in other vertebrate groups so far. Moreover, the characteristics of each actin type could be related to the energetic demands of white and red muscle tissues; (2) A sexually dimorphic gene expression occurs in brain tissue of males and females of L. macrocephalus related to the gene that codes for the protein APBA2 (amyloid beta (A4) precursor protein-binding, family A, member 2), that is associated to the production of the β amyloid protein which, in turns, is related to the Alzheimer disease; (3) The fabp6 (fatty acid-binding protein 6) gene structure of D. rerio is similar to most of the other members of the multigenic fabp family of other vertebrates, and its tissue distribution in embryos, larvae, and adults of the species is also similar to the one found in other organisms, specially mammals; (4) The genes that code for the protein p300 (E1A binding protein p300) and for the ADCY2 protein (adenylate cyclase 2) represent two candidate differentially expressed genes among nontransgenic and growth hormone (GH)-transgenic homozygote and hemizygote samples of Danio rerio, reinforcing previous data that evidenced that these proteins are involved in muscle formation/growth and in GH release. Therefore, the obtained data contributes to the comprehension of the evolution, function, and molecular organization of several genes and can be used as basal information on future studies. Moreover, the results confirm the utility of Danio rerio as a model species in genetic analyses and indicate that Leporinus macrocephalus represents a potential experimental model in analyses of genes involved in the development

of human diseases, as neurodegenerative alterations, muscle dystrophies and cancer.

Considerações Iniciais e Objetivos

Nos últimos anos, vários estudos genéticos têm dedicado especial importância a análises de expressão gênica diferencial. Estes estudos permitem a caracterização de genes e identificação e avaliação do nível de expressão de determinados transcritos em relação a diversas variáveis. Desta forma, é possível analisar e comparar, por exemplo, amostras relacionadas a diferentes períodos de desenvolvimento ou a tecidos distintos, submetidas a diferentes tratamentos, presentes em distintos ambientes, ou mesmo amostras de diferentes sexos e de organismos transgênicos e não transgênicos.

Entre as técnicas mais utilizadas para isolamento e identificação de transcritos diferencialmente expressos (DETs - *Differentially Expressed Transcripts*), destacam-se: RT-PCR (*Reverse Transcription - Polymerase Chain Reaction*), DD (*Differential Display*), SH (*Suppression Hybridization*), SSH (*Suppression Subtractive Hybridization*), RAP-PCR (*RNA Arbitrarily Primed PCR*), RDA (*Representational Difference Analysis*), SAGE (*Serial Analysis of Gene Expression*), *microarrays* de DNA, PCR em tempo real, hibridação *in situ* e *Northern blotting*. Das metodologias citadas acima, RT-PCR, DD, PCR em tempo real e hibridação *in situ* foram utilizadas no presente trabalho. Os métodos de RT-PCR e DD, além de exibirem considerável eficiência e ampla aplicação em estudos de expressão diferencial, caracterizam-se pela simplicidade, custo relativamente reduzido e, em especial para DD, não requerem um conhecimento prévio do genoma que se deseja estudar.

Os trabalhos relacionados à identificação e caracterização de DETs abrangem grande diversidade de organismos e incluem análises em vários modelos experimentais como bactérias, fungos, plantas, invertebrados e vertebrados. Entretanto, esta área de pesquisa é ainda recente e, entre os vertebrados, embora os peixes representem um dos grupos mais diversificados, até o momento, poucas espécies apresentam seus genomas caracterizados.

Entre os peixes mais amplamente utilizados em estudos genômicos, destaca-se a espécie Danio rerio, popularmente conhecida como zebrafish ou paulistinha e de grande importância econômica para a área de aquariofilia. Apesar de nativa do continente asiático, apresenta ampla distribuição geográfica, podendo ser encontrada em quase todos os países do mundo e representa um dos mais importantes modelos animais utilizados em estudos de desenvolvimento embrionário, comportamentais, fisiológicos e genéticos. A facilidade de manipulação genética nesta espécie permitiu o desenvolvimento de várias linhagens transgênicas, o que tem contribuído para o desenvolvimento de novos estudos relacionados à expressão gênica. Atualmente, o genoma de Danio rerio apresenta-se quase totalmente caracterizado (~ 75%) devido ao projeto "Danio rerio Sequencing", iniciado em 2001 pelo Wellcome Trust Sanger Institute para pesquisas genômicas. Atualmente, inúmeras pesquisas, em diferentes áreas, vêm sendo desenvolvidas em D. rerio, especialmente direcionadas à criação de novas linhagens transgênicas e à compreensão do surgimento e desenvolvimento de doenças humanas, como câncer e processos degenerativos.

Por outro lado, trabalhos genéticos direcionados para diversos outros peixes são extremamente escassos e, para um grande número de espécies, são ainda inexistentes. Nesta perspectiva, o território brasileiro representa uma fonte imensa de recursos ainda inexplorados, especialmente levando-se em consideração que várias espécies apresentam estimável valor comercial e ecológico, sendo muitas delas endêmicas. Entre estas espécies, pode-se citar

6

Leporinus macrocephalus, popularmente conhecida como piauçu ou piavuçu, nativa do Pantanal e que representa uma importante fonte alimentar que, nos últimos anos, tem sido intensivamente explorada como recurso natural e de piscicultura no Brasil. Apesar de sua importância econômica, os dados genéticos acerca desta espécie são ainda extremamente reduzidos.

Dada a importância de *Danio rerio* e *Leporinus macrocephalus*, o presente trabalho teve o intuito de isolar e caracterizar transcritos diferencialmente expressos nestas duas espécies, visando ampliar os dados genéticos em peixes, grupo que ocupa uma posição basal na história evolutiva dos vertebrados. Desta forma, a aquisição de um maior número de informações genéticas em espécies de peixes pode contribuir à compreensão da evolução, função e organização molecular de diversos genes. Além deste objetivo geral, foram traçados objetivos específicos, relacionados a diferentes sub-propostas de pesquisa, detalhados abaixo.

- Identificar e caracterizar genes de actina de tecido muscular esquelético branco e vermelho de *Leporinus macrocephalus*.
- Identificar e caracterizar transcritos diferencialmente expressos em tecido cerebral de machos e fêmeas de *Leporinus macrocephalus*.
- Determinar a estrutura, a organização e a distribuição espaço-temporal do gene fabp6 (fatty acid binding protein type 6) em Danio rerio.
- Identificar e caracterizar transcritos diferencialmente expressos em tecido cerebral de linhagens de *Danio rerio* transgênicas e não-transgênicas para o gene do hormônio do crescimento (GH).

Introdução Geral

Transcritos diferencialmente expressos

Nas últimas décadas, pôde-se presenciar um considerável crescimento da genômica, área que se refere à aquisição de dados do genoma, especialmente à obtenção da seqüência completa do material genético dos organismos. Embora a obtenção desta informação não seja uma tarefa simples, a ampliação desta área de pesquisa está intimamente relacionada aos conhecimentos genéticos gerados com o advento dos denominados Projetos Genomas. Atualmente, vários projetos que visam desvendar o código genético vêm sendo desenvolvidos em organismos diversos como bactérias, fungos, plantas, invertebrados e vertebrados e os resultados gerados podem representar dados básicos para o desenvolvimento de trabalhos posteriores que visem compreender os mecanismos de expressão de diversos genes (Carneiro et al. 2006). Além dos dados de projetos de seqüenciamento total do genoma de várias espécies, resultados de projetos que objetivam a construção de bibliotecas de ESTs (Expressed Sequence Tags) são diretamente utilizados em pesquisas voltadas à elucidação dos mecanismos biológicos envolvidos na expressão gênica e em sua regulação (Carneiro et al. 2006).

Embora abordagens mais amplas de análise de expressão gênica em nível genômico (*Genome-Wide Expression Profiling*) constituam o maior desafio atual para identificação e análise simultânea de um grande número de genes envolvidos nos mais diversos processos biológicos - desde o desenvolvimento dos organismos até suas interações com fatores ambientais (Donson *et al.* 2002) - projetos pontuais, direcionados à identificação e caracterização de genes

específicos, também vêm sendo realizados em diversas espécies. Através da aplicação de diferentes tecnologias, análises deste tipo podem gerar dados primordiais para futuros projetos genômicos de larga escala (Matz & Lukyanov 1998, Ponsuksili *et al.* 2001).

Nas últimas décadas, inúmeras técnicas em biologia molecular foram desenvolvidas visando facilitar o isolamento e a caracterização de transcritos diferencialmente expressos (DETs - *Differentially Expressed Transcripts*), ou seja, genes expressos como RNAs mensageiros (RNAm) para síntese de proteínas que diferem em abundância entre amostras de tipos celulares ou tecidos específicos e cuja regulação pode estar relacionada a mecanismos ambientais, fisiológicos e/ou químicos. Entre estas diversas metodologias, podem ser citadas RT-PCR (*Reverse Transcription-Polymerase Chain Reaction*) (Kawasaki 1990), hibridação subtrativa (Duguid *et al.* 1988), DD (*Differential Display*) (Liang & Pardee 1992), RAP-PCR (*RNA Arbitrarily Primed PCR*) (Welsh *et al.* 1992), RDA (*Representational Difference Analysis*) (Lisitsyn *et al.* 1993), SAGE (*Serial Analysis of Gene Expression*) (Velculescu *et al.* 1995), *microarrays* de DNA (Shena *et al.* 1995), SSH (*Suppression Subtractive Hybridization*) (Diatchenko *et al.* 1996), PCR em tempo real (Freeman *et al.* 1999), hibridação *in situ* (Wilkinson 1998) e *Northern blotting* (Alwine *et al.* 1979).

Embora esta vasta gama de técnicas para isolamento e caracterização de transcritos diferencialmente expressos permita sua aplicação em estudos de diversas áreas, a escolha de uma destas metodologias deve considerar as características dos experimentos, os objetivos propostos e o organismo em estudo (Matz & Lukyanov 1998). Além disso, apesar de suas inúmeras vantagens, normalmente, a aplicação de metodologias experimentais para prospecção de

transcritos diferencialmente expressos nem sempre representa uma tarefa fácil, pois, na maioria das vezes, estes apresentam níveis de expressão extremamente baixos. Entre os milhares de genes que compõem o genoma dos organismos vivos, somente cerca de 10-15% são expressos em um determinado tipo de tecido durante um dado período, o que torna relativamente difícil sua identificação e caracterização (Liao & Zhang 2006). Além disso, as tecnologias disponíveis para o isolamento de DETs são geralmente laboriosas e de alto custo.

O crescente interesse por estudos de transcritos diferencialmente expressos pode ser comprovado pela publicação de mais de 30.000 trabalhos nas mais diversas áreas de aplicação e pelas inúmeras revisões sobre o tema (Medline, Julho de 2008 PubMed. NCBI Web site http://www.ncbi.nlm.nih.gov/entrez/guery.fcgi?db=PubMed). Grande parte das pesquisas baseadas em genes que apresentam expressão diferencial entre amostras distintas vem sendo direcionada à biologia do desenvolvimento (Kanamori 2000, Xie et al. 2001, Ando & Sakai 2002, Anway et al. 2003, Dann et al. 2003, Kim et al. 2004, Sunagawa & Magae 2005, Jianzhen et al. 2007, Williams et al. 2007, Chu et al. 2008, Elis et al. 2008, Sugimoto & Endoh 2008), biologia celular e fisiologia de tecidos (Eriksson et al. 1999, Ahmed et al. 2000, Boag et al. 2000, Kim et al. 2001, Blásquez & Piferrer 2004, Saunders & Magor 2004, Azumi et al. 2005, Venkatesh et al. 2005, Jin et al. 2006, Koury et al. 2007, Bevan et al. 2008, Cervigni et al. 2008, Kim et al. 2008, Smartt & Erickson 2008, Wong et al. 2008, Zhang et al. 2008) e a processos tumorais e diagnóstico de diversas doenças (An et al. 2000, Miller et al. 2001, Lee & Cicila 2002, Hoei-Hansen et al. 2004, Li & Chen 2004, Clinton et al. 2005, Kim et al. 2005, Poulin & Labelle 2005, Sharma et al. 2005, Zhang et al. 2005, Blaes et al. 2007, Collins et *al.* 2007, Nagai *et al.* 2007, Kamalian *et al.* 2008, Shin *et al.* 2008), visando gerar informações sobre a função gênica e suas relações com processos metabólicos e fisiológicos.

Embora análises nas áreas de biologia do desenvolvimento, fisiologia e carcinogênese representem a maior gama de dados relativos a transcritos diferencialmente expressos, outras áreas de pesquisa também têm merecido destaque. Uma destas áreas refere-se à caracterização de DETs em relação a alterações ambientais - como contaminação, variação drástica de temperatura e/ou disponibilidade de água (Llado *et al.* 1998, Casagrande *et al.* 2001, D'Cotta *et al.* 2001, Carginale *et al.* 2002, Chang *et al.* 2004, Basile *et al.* 2005, Meyer *et al.* 2005, Markovskaya *et al.* 2007, Mishra *et al.* 2007, Li *et al.* 2008, Smith-Keune & Dove 2008, Xu *et al.* 2008) - e a tratamentos químicos (Denslow *et al.* 2001a, Denslow *et al.* 2001b, Miller *et al.* 2001, Wong *et al.* 2002, Akbar *et al.* 2007, Zhang *et al.* 2007, Caamal-Velázquez *et al.* 2007, Zhang *et al.* 2008, Karakurt & Huber 2008, Kavar *et al.* 2008, Torelli *et al.* 2008). Estudos com este enfoque vêm também sendo, ao longo dos anos, ampliados, especialmente devido ao crescente aumento dos problemas gerados por alterações ambientais antrópicas.

Apesar do grande número de trabalhos direcionados a análises de expressão gênica diferencial, que abrangem uma grande diversidade de organismos, esta área de pesquisa é ainda recente. Particularmente em peixes, até o momento, existem poucos dados, especialmente se considerarmos que estes representam um dos grupos de vertebrados mais diversificados (Nelson 1994). Grande parte dos trabalhos de expressão gênica diferencial restringe-se a espécies nativas da América do Norte e Europa, especialmente de peixes de interesse comercial, como Salmoniformes, visando gerar benefícios para a piscicultura (Parrington & Coward 2002). Estes estudos incluem o isolamento e a caracterização de genes relacionados à determinação sexual (D'Cotta *et al.* 2001, Denslow *et al.* 2001a, Denslow *et al.* 2001b), ao crescimento e desenvolvimento (Kanamori 2000, Xie *et al.* 2001, Dann *et al.* 2003), à resistência a patógenos (Collins *et al.* 2007) e a condições ambientais adversas (Carginale *et al.* 2002, Larkin *et al.* 2003, Meyer *et al.* 2005). Além disso, espécies de peixes consideradas modelos animais também têm sido alvo de estudos em expressão gênica diferencial. Devido ao genoma compacto (390 Mb) e ao conteúdo de genes similar ao de humanos, *Fugu rubripes* (baiacu ou *pufferfish*) tem sido explorado como um modelo do genoma de vertebrados para identificação de genes e outros elementos funcionais nos genomas de outros animais (Brenner *et al.* 1993).

Desta forma, a ampliação de análises genéticas em peixes, que compreendem um grupo vertebrado basal, poderá levar à identificação e caracterização de diversos genes, assim como fornecer maiores informações sobre sua organização, função e evolução.

A espécie Leporinus macrocephalus

A família Anostomidae, grupo de peixes da ordem Characiformes relativamente rico em espécies distribuídas por toda a América do Sul e América Central, compreende 12 gêneros: *Abramites, Anostomus, Gnathodolus, Laemolyta, Leporellus, Leporinus, Rhytiodus, Schizodon, Synaptolaemus, Anostomoides, Pseudanos e Sator.* Entre estes, o gênero *Leporinus* é o mais diverso, possuindo 87 espécies válidas (Garavello & Britski 2003). Popularmente conhecida como piauçu ou piavuçu e nativa da região do Pantanal, a espécie *Leporinus macrocephalus* (Figura 1) representa um dos principais peixes de pesca de subsistência e de pesca esportiva no Brasil (Garavello & Britski 1988, Martins & Yoshitoshi 2003). Além disso, seu cultivo em estações de piscicultura tem se mostrado um empreendimento promissor, devido a seu hábito alimentar omnívoro, facilidade de adaptação às condições de cativeiro, bom desempenho reprodutivo e excelente taxa de crescimento (Corrêa Filho 2000, Takahashi *et al.* 2004). A espécie é gonocorística, ou seja, os indivíduos se desenvolvem somente como machos ou somente como fêmeas e permanecem, assim, durante todo seu ciclo de vida. Embora machos e fêmeas da espécie atinjam, na maturidade, cerca de 60 cm de comprimento, diferenças de peso e tamanho podem ser observadas entre os sexos em indivíduos adultos - machos da espécie apresentam um maior tamanho corporal (J. Senhorini - CEPTA/IBAMA, comunicação pessoal).

Dada a importância de *L. macrocephalus* como recurso alimentar, diversos estudos voltados à criação da espécie em cativeiro vêm sendo realizados, especialmente análises de crescimento, alimentação e reprodução (Soares *et al.* 2000, Pinto *et al.* 2001, Nagae *et al.* 2002, Ribeiro & Godinho 2003, Minucci *et al.* 2005, Normandes *et al.* 2006, Godinho 2007, Albinati *et al.* 2007). Além disso, *L. macrocephalus* tem sido, nos últimos anos, uma das espécies de destaque para pesca esportiva, especialmente nos estados do sul e sudeste do Brasil (Queiroz *et al.* 2002).

13



Figura 1: Exemplar adulto de Leporinus macrocephalus (piauçu ou piavuçu).

Por outro lado, os estudos genéticos referentes a *Leporinus macrocephalus* são escassos e a maioria das análises refere-se a dados citogenéticos. Embora grande parte das espécies que compõem o gênero *Leporinus* não apresente cromossomos sexuais morfologicamente diferenciados entre machos e fêmeas (Galetti *et al.* 1995, Venere *et al.* 2004), *L. macrocephalus* é caracterizado por apresentar um cariótipo composto por 54 cromossomos e, em especial, por apresentar um sistema de determinação sexual do tipo ZZ/ZW, o que caracteriza uma heterogametia feminina. Assim, as fêmeas possuem um par cromossômico heteromórfico consistindo de um cromossomo submetacêntrico (cromossomo Z), equivalente ao cromossomo nº 2 dos machos, e um cromossomo subtelocêntrico (cromossomo W), o maior do complemento cariotípico e ausente nos machos (Galetti & Foresti 1987). Tal característica torna a espécie extremamente interessante para estudos relacionados à determinação e diferenciação sexual e para estudos sobre a evolução dos caracteres envolvidos nestes processos.

As demais análises genéticas em *L. macrocephalus* referem-se à caracterização do DNA ribossomal 5S para análises filogenéticas (Ferreira *et al.* 2007) e isolamento e caracterização de locos microssatélites para utilização em

estudos de diversidade genética populacional (Morelli *et al.* 2007). Embora os dados genéticos em *L. macrocephalus* sejam ainda extremamente restritos, este cenário tende a mudar nos próximos anos. O crescente desenvolvimento do setor de piscicultura no Brasil - devido ao grande potencial hídrico, à enorme riqueza de espécies nativas e às áreas de clima propício à atividade encontradas no país -, associado à disponibilidade de diversas técnicas de biologia molecular, deverão levar à implementação de novas análises genéticas em espécies de peixes de interesse comercial, incluindo *Leporinus macrocephalus*.

A espécie Danio rerio

A espécie *Danio rerio* (Figura 2) (Ordem Cypriniformes, Família Cyprinidae), popularmente conhecida como paulistinha ou *zebrafish*, embora nativa do Sul e Sudeste da Ásia, tornou-se amplamente distribuída, podendo ser encontrada em todos continentes. Características como a fácil adaptação ao ambiente de cativeiro, o espaço relativamente pequeno necessário para sua manutenção, a prole numerosa, a produção contínua de ovos e o período curto de desenvolvimento embrionário e larval representam aspectos interessantes para os produtores, especialmente da área de aquariofilia (Westerfield 2000). Além destas características, *Danio rerio* é considerada uma espécie modelo em estudos de desenvolvimento embrionário, comportamentais, fisiológicos e genéticos de vertebrados, especialmente devido à transparência de estruturas como ovos e larvas, que permitem análises diretas dos estágios embrionários e da expressão espaço-temporal de determinados genes de interesse, através da aplicação de métodos de hibridação *in situ* (Detrich *et al.* 1998, Dahm & Geisler 2006, Dahm *et al.* 2007, Bass & Gerlai 2008, Fadool & Dowling 2008).



Figura 2: Exemplar adulto de *Danio rerio* (paulistinha ou *zebrafish*).

Ao contrário de diversas outras espécies de peixes, entre estas *Leporinus macrocephalus*, *D. rerio* apresenta inúmeros dados referentes à caracterização de seu genoma. A espécie apresenta um cariótipo composto por 50 cromossomos e não apresenta cromossomos sexuais morfologicamente diferenciados, sendo seu mecanismo de determinação sexual desconhecido e, até o momento, nenhum gene relacionado à determinação sexual foi identificado (Amores & Postlethwaite 1999, Sola & Gornung 2001). Acredita-se que fatores ambientais e ecológicos, como temperatura e densidade populacional, podem influenciar na determinação do sexo de *zebrafish* (Lawrence *et al.* 2008).

Atualmente, aproximadamente 75% do genoma de *Danio rerio* já se encontra caracterizado, devido ao projeto de seqüenciamento iniciado em 2001 pelo *Wellcome Trust Sanger Institute* (Cambridge, Inglaterra) para pesquisas genômicas, e uma rede de dados (*Zebrafish Information Network* - ZFIN: <u>http://www.zfin.org</u>) foi criada com o intuito de integrar diversos dados referentes ao genoma desta espécie de peixe (Sprague et al. 2003). Além disso, nos últimos cinco anos, com o advento de novas técnicas biotecnológicas e experimentais de análises genômicas, o número de estudos genéticos neste modelo animal cresceu

significativamente - mais de 640 mil ESTs (*Expressed Sequence Tags*) que representam transcritos de *zebrafish* foram seqüenciados e mais de 6.800 RNAs mensageiros foram identificados e caracterizados (Zon & Peterson 2005, Dahm & Geisler 2006). A facilidade de manipulação genética nesta espécie ainda levou ao desenvolvimento de várias linhagens transgênicas, o que tem contribuído para o desenvolvimento de novos estudos relacionados à expressão gênica e já permitiu um grande avanço na elucidação das relações existentes entre diversos genes de *zebrafish* e de outros vertebrados (Sprague *et al.* 2003).

Atualmente, diversas pesquisas, em diferentes áreas, vêm sendo desenvolvidas em *D. rerio*, especialmente direcionadas à criação de novas linhagens transgênicas (Linney & Udvadia 2004, Ashworth & Brennan 2005, Huang *et al.* 2005, Burket *et al.* 2008, Wu *et al.* 2008) e à compreensão do surgimento e desenvolvimento de doenças humanas, como câncer e processos degenerativos (Patton & Zon 2005, Lam & Gong 2006, Panulo *et al.* 2006, Goessling *et al.* 2007, Merlino & Khanna 2007, Beckman 2007, Feitsma & Cuppen 2008). Como perspectiva, pode-se prever a continuação e ampliação das análises genéticas em *Danio rerio*, tanto na área biológica como na área de medicina.

Referências Bibliográficas

- Ahmed N, Siddiqui AA, Ahmed A (2000) DDRT-PCR: use of agarose gels for detection of amplified products. *Molecular Vision* **6**: 144-147.
- Akbar MA, Chatterjee NS, Sen P, Debnath A, Pal A, Bera T, Das P (2004) Genes induced by a high-oxygen environment in *Entamoeba histolytica*. *Molecular and Biochemical Parasitology* **133**: 187-196.

17

- Albinati ACL, Moreira ELT, Albinati RCB, Carvalho JV, Santos GB, Lira AD (2007)
 Toxidade aguda do herbicida roundup para piauçu (*Leporinus macrocephalus*).
 Revista Brasileira de Saúde e Produção Animal 8: 184-192.
- Alwine JC, Kemp DJ, Parker BA, Reiser J, Renart J, Stark GR, Wahl GM (1979) Detection of specific RNAs or specific fragments of DNA by fractionation in gels and transfer to diazobenzyloxymethyl paper. *Methods in Enzymology* **68**: 220-242.
- Amores A & Postlethwaite JH (1999) Banded chromosomes and the zebrafish karyotype. *Methods in Cell Biology* **60**: 323-338.
- An G, Meka CSR, Luo GZ, Bright SP, Cazares L, Wright GL, Veltri RW (2000) Cloning and characterization of UROC28, a novel gene overexpressed in prostate, breast, and bladder cancers. *Cancer Research* **60**: 7014-7020.
- Anderson TD, Jin-Clark Y, Begum K, Starkey SR, Zhu KY (2008) Gene expression profiling reveals decreased expression of two hemoglobin genes associated with increased consumption of oxygen in *Chironomus tentans* exposed to atrazine: A possible mechanism for adapting to oxygen deficiency. *Aquatic Toxicology* **86**: 148-156.
- Ando S & Sakai S (2002) Isolation of an ethylene-responsive gene (ERAF16) for a putative methyltransferase and correlation of ERAF16 gene expression with female flower formation in cucumber plants (*Cucumis sativus*). *Physiologia Plantarum* **116**: 213-222.
- Anway MD, Li Y, Ravindranath N, Dym M, Griswold MD (2003) Expression of testicular germ cell genes identified by differential display analysis. *Journal of Andrology* 24: 173-184.
- Ashworth R & Brennan C (2005) Use of transgenic zebrafish reporter lines to study calcium signalling in development. *Briefly Function on Genomics and Proteomics* **4**: 186-193.
- Azumi K, Sasaki T, Okochi K, Yamasaki S, Saito T, Takayama H, Yokosawa H (2005) Differential display analysis reveals the expression of glutathione Stransferase ω and novel genes throughan ITAM- containing receptor in ascidian immunocytes. *Immunogenetics* **57**:444-452.

- Badonnel K, Denis JB, Caillo M, Monnerie R, Piumi F, Potier MC, Salesse R, Baly C (2007) Transcription profile analysis reveals that OBP-1F Mrna is downregulated in the olfactory mucosa following food deprivation. *Chemical Senses* 32: 697-710.
- Basile A, di Nuzzo RA, Capasso C, Sorbo S, Capasso A, Carginale V (2005) Effect of cadmium on gene expression in the liverwort *Lunularia cruciata*. *Gene* 356: 153-159.
- Bass SLS & Gerlai R (2008) Zebrafish (*Danio rerio*) responds differentially to stimulus fish: The effects of sympatric and allopatric predators and harmless fish. *Behavioural Brain Research* **186**: 107-117.
- Beckman M (2007) Zebrafish take the stage in cancer research. *Journal of the Natural Cancer Institute* **99**: 500-501.
- Bevan S, Vakharia V, Parker D (2008) Changes in gene expression and integrinmediated structural changes are associated with long-term plasticity of a spinal cord locomotor network. *Neuroscience* **152**: 160-168.
- Blaes N, Elbaz M, Heitz F, Caussé E, Glock Y, Puel J, Bayard F (2007) Differential display fingerprints: new approach to characterize smooth muscle cells and human coronary atherectomy tissues. *Pathologie Biologie* **55**: 328-335.
- Blázquez M & Piferrer F (2004) Cloning, sequence analysis, tissue distribution, and sex-specific expression of the neural form of P450 aromatase in juvenile sea bass (*Dicentrarchus labrax*). *Molecular and Cellular Endocrinology* **219**: 83-94.
- Boag PR, Newton SE, Hansen NP, Christensen CM, Nansen P, Gasser RB (2000)
 Isolation and characterization of sex-specific transcripts from
 Oesophagostomum dentatum by RNA arbitrarily-primed PCR. *Molecular and Biochemical Parasitology* 108: 217-224.
- Brenner S, Elgar G, Sandford R, Macrae A, Venkatesh B, Aparício S (1993) Characterization of the Pufferfish (*Fugu*) genome as a compact model vertebrate genome. *Nature* **366**: 265-268.
- Burket CT, Montgomery JE, Thummel R, Kassen SC, Lafave MC, Langenau DM, Zon LI, Hyde DR (2008) Generation and characterization of transgenic zebrafish lines using different ubiquitous promoters. *Transgenic Research* **17**: 265-279.

- Caamal-Velázquez JH, Chi-Manzanero BH, Canche-Yam JJ, Castaño E, Rodríguez-Zapata LC (2007) Low temperature induce differential expression genes in banana fruits. *Scientia Horticulturae* **114**: 83-89.
- Carginale V, Capasso C, Scudiero R, Parisi E (2002) Identification of cadmiumsensitive genes in the antarctic fish *Chionodraco hamatus* by messenger RNA differential display. *Gene* **299**: 117-124.
- Carneiro NP, Carneiro AA, Guimarães CT, Paiva E (2006) Desvendando o código genético. *Pesquisa e Biotecnologia* **6**: 1-11.
- Casagrande EC, Farias JRB, Neumaier N, Oya T, Pedroso J, Martins PK, Breton MC, Nepomuceno AL (2001) Expressão gênica diferencial durante déficit hídrico em soja. *Revista Brasileira de Fisiologia Vegetal* **13**: 168-184.
- Cervigni GDL, Paniego N, Pessino S, Selva JP, Díaz M, Spangenberg G, Echenique V (2008) Gene expression in diplosporous and sexual *Eragrostis curvula* genetypes with differing ploidy levels. *Plant Molecular Biology* 67: 11-23.
- Chang IS, Groh JL, Ramsey MM, Ballard MM, Krumholz, LR (2004) Differential expression of *Desulfovibrio vulgaris* genes in response to Cu(II) and Hg(II) toxicity. *Applied and Environmental Microbiology* **70**: 1847-1851.
- Chu F-H, Lee Y-R, Chou S-J, Chang T-T, Shaw J-F (2008) Isolation and analysis of genes specifically expressed during basidiomatal development in *Antrodia cinnamomea* by subtractive PCR and cDNA microarray. *FEMS Microbiology Letters* **280**: 150-159.
- Clinton M, Miele G, Nandi S, McBride D (2005) Identification of disease markers by differential display: prion disease. *Methods in Molecular Biology* **317**: 157-178.
- Collins CM, Olstad K, Sterud E, Jones CS, Noble LR, Mo TA, Cunningham CO (2007) Isolation of a novel fish thymidylate kinase gene, upregulated in Altantic salmon (*Salmo salar* L.) following infection with the monogenean parasite *Gyrodactylus salaries*. *Fish and Shellfish Immunology* **23**: 793-807.
- Corrêa Filho RAC (2000) Métodos de comparação do crescimento entre grupos genéticos de piauçu (*Leporinus macrocephalus*). Tese de Doutorado. Universidade Estadual Paulista.
- Daham R & Geisler R (2006) Learning from small fry: The zebrafish as a genetic model organism for aquaculture fish species. *Marine Biology* **8**:329-345.

- Dahm R, Schonthaler HB, Soehn AS, Marle J, Vrensen GFJM (2007) Development and adult of the eye lens in the zebrafish. *Experimental Eye Research* **85**: 74-89.
- Dann SG, Allison WT, Levin DB, Hawryshyn, CW (2003) Identification of a unique transcript down-regulated in the retina of rainbow trout (*Oncorhynchus mykiss*) at somltification. *Comparative Biochemistry and Physiology* Part B **136**: 849-860.
- D'Cotta H, Fostier A, Guiguen Y, Govoroun M, Baroiller JF (2001) Search for genes involved in the temperature-induced gonadal sex differentiation in the tilapia, *Oreochromis niloticus*. *Journal of Experimental Zoology* **290**: 574-585.
- Denslow ND, Bowman CJ, Ferguson RJ, Lee HS, Hemmer MJ, Folmar LC (2001a) Induction of gene expression in sheepshead minnows (*Cyprinodon variegatus*) treated with 17 beta-estradiol, diethylstilbestrol, or ethinylestradiol: The use of mRNA fingerprints as an indicator of gene regulation. *General and Comparative Endocrinology* **121**: 250-260.
- Denslow ND, Lee HS, Bowman CJ, Hemmer MJ, Folmar LC (2001b) Multiple responses in gene expression in fish treated with estrogen. *Comparative Biochemistry and Physiology B-Biochemistry and Molecular Biology* **129**: 277-282.
- Detrich HW, Westerfield M, Zon LI (1998) *The zebrafish: Biology.* Academic Press, New York.
- Diatchenko L, Lau YFC, Campbell AP, Chenchik A, Moqadam F, Huang B (1996) Supression subtractive hybridization: a method for generating differentially regulated or tissue-specific cDNA probes and libraries. *Proceedings of the Natural Academy of Sciences USA* **93**: 6025-6030.
- Donson J, Fang Y, Espiritu-Santo G, Xing W, Salazar A, Miyamoto S, Armendarez V, Volkmuth W (2002) Comprehensive gene expression analysis by transcript profiling. *Plant Molecular Biology* **48**: 75-97.
- Duguid JR, Rohwer RG, Seed B (1988) Isolation of cDNAs of scrapie-modulated RNAs by subtractive hybridization of a cDNA library. *Proceedings of the Natural Academy of Science USA* **85**: 5738-5742.

- Elis S, Batellier F, Couty I, Balzergue S, Martin-Magniette M-L, Monget P, BlesboisE, Govoroun MS (2008). Search for the genes involved in oocyte maturation and early embryo development in the hen. *BMC Genomics* 9: 110.
- Eriksson A, Wahlestedt C, Nordqvist K (1999) Isolation of sex-specific cDNAs from fetal mouse brain using mRNA differential display and representational difference analysis. *Molecular Brain Research* **74**: 91-97.
- Fadool JM & Dowling JE (2008) Zebafish: a model system for the study of eye genetics. *Progress in Retinal and Eye Research* **27**: 89-110.
- Feitsma H & Cuppen E (2008) Zebrafish as a cancer model. *Molecular and Cancer Research* **6**: 685-694.
- Ferreira IA, Oliveira C, Venere PC, Galetti PM, Martins C (2007) 5S rDNA variation and its phylogenetic inference in the genus *Leporinus* (Characiformes: Anostomidae). *Genetica* **129**: 253-257.
- Freeman WM, Walker SJ, Vrana SJ (1999) Quantitative RT-PCR: Pitfalls and potential. *Biotechniques* **26**: 112-125.
- Galetti PM, Lima NRW, Venere PC (1995) A monophyletic ZW sex chromosome system in *Leporinus* (Anostomidae, Characiformes). *Cytologia* **60**: 375-382.
- Galetti PM & Foresti F (1987) Two new cases of ZZ/ZW heterogamety in *Leporinus* (Anostomidae, Characiformes) and their relationships in the phylogeny of the group. *Revista Brasileira de Genética* **10**: 135-140.
- Garavello JC & Britski HA (1988) *Leporinus macrocephalus* sp. n. da Bacia do Rio Paraguai (Ostariophysi, Anostomidae). *Naturalia* **13**: 67-74.
- Garavello JC & Britski HA (2003) Family Anostomidade (Head-standers). In: Reis RE, Kullander SO, Ferraris CJ (eds.). *Check list of the freshwater fishes of South and Central America*. EDIPUCRS. pp.75-84.
- Godinho HP (2007) Estratégias reprodutivas de peixes aplicadas à aqüicultura: bases para o desenvolvimento de tecnologias de produção. *Revista Brasileira de Reprodução Animal* **31**: 351-360.
- Goessling W, North TE, Zon LI (2007) New waves of discovery: modeling cancer in zebrafish. *Journal of Clinical Oncology* **25**: 2473-2479.
- Hoei-Hansen CE, Nielsen JE, Almstrup K (2004) Identification of genes differentially expressed in testes containing carcinoma in situ. *Molecular Human Reproduction* **10**: 423-431.

- Hosoi-Tanabe T, Tomishima S, Nagai S, Sako Y (2005) Identification of a gene induced in conjugation-promoted cells of toxicmarine dinoflagellates Alexandrium tamarense and Alexandrium catenella using differential display analysis. *FEMS Microbiology Letters* **251**:161-168.
- Huang CJ, Jou TS, Ho YL, Lee WH, Jeng YT, Hsieh FJ, Tsai HJ (2005) Conditional expression of a myocardium-specific transgene in zebrafish transgenic lines. *Developmental Dynamics* **233**: 1294-1303.
- Jianzhen H, Haitian M, Liming Y, Sixiang Z (2007). Developmental changes of protein profiles in the embryonic sanhuang chicken liver. *Journal of Veterinary Medicine* **54**: 464-469.
- Jin EJ, Park JH, Lee SY, Chun JS, Bang OS, Kang SS (2006) Wnt-5 is involved in TGF-β3-stimulated chondrogenic differentiation of chick wing bud mesenchymal cells. *The International Journal of Biochemistry & Cell Biology* **38**: 186-195.
- Kamalian L, Gosney JR, Forootan SS, Foster CS, Bao ZZ, Beesley C, Ke YQ (2008) Increased expression of Id family proteins in small cell lung cancer and its prognostic significance. *Clinical Cancer Research* 14: 2318-2325.
- Kanamori, A. (2000) Systematic identification of genes expressed during early oogenesis in Medaka. *Molecular Reproduction and Development* **55**: 31-36.
- Karakurt Y & Huber DJ (2008) Cloning and characterization of differentially expressed genes in ethylene-treated water-melon fruit. *Postharvest Biology and Technology* **48**: 372-377.
- Kavar T, Maras M, Kidric M, Sustar J, Meglic V (2008) Identification of genes involved in the responsive of leaves of *Phaseolus vulgaris* to drought stress. *Molecular Breeding* **21**: 159-172.
- Kawasaki ES (1990) Amplification of RNA. In: *PCR Protocols: A guide to methods and applications*. Innis *et al.* (eds.). Academic Press, New York, pp. 21-27.
- Kim DI, Eo HS, Joh JH (2005) Identification of differentially expressed genes in primary varicose veins. *Journal of Surgical Research* **123**: 222-226.
- Kim H, You S, Farris J, Foster LK, Choi YJ, Foster DN (2001) Gonad-specific expression of two novel chicken complementary DNA isoforms. *Biology of Reproduction* 64: 1473-1480.

- Kim TW, Kim HJ, Lee C, Kim HY, Baek S-H, Kim JH, Kwon K-S, Kim J-R (2008) Identification of replicative senescence-associated genes in human umbilical vein endothelial cells by an annealing control primer system. *Experimental Gerontology* **43**: 286-295.
- Kim Y-J, Kwak CI, Gu YY, Hwang IT, Chun JY (2004) Annealing control primer system for identification of differentially expressed genes on agarose gels. *Biotechniques* 36: 424-428.
- Koury S, Yarlagadda S, Moskalik-Liermo K, Popli N, Kim N, Apolito C, Peterson A,
 Zhang X, Zu P, Tamburlin J, Bofinger D (2007) Differential gene expression
 during terminal erythroid differentiation. *Genomics* **90**: 574-582.
- Lam SH & Gong Z (2006) Modeling liver cancer using zebrafish: a comparative oncogenomics approach. *Cell Cycle* **5**: 573-577.
- Larkin P, Knoebl I, Denslow, ND (2003) Differential gene expression analysis in fish exposed to endocrine disrupting compounds. *Comparative Biochemistry and Physiology Part B* **136**: 149-161.
- Lawrence C, Ebersole JP, Kesseli RV (2008) Rapid growth and out-crossing promote female development in zebrafish (*Danio rerio*). *Environmental Biology of Fishes* **81**:239-246.
- Lee SJ & Cicila GT (2002) Functional genomics in rat models of hypertension: using differential expression and congenic strains to identify and evaluate candidate genes. *Critical Reviews in Eukaryotic Gene Expression* **12**: 297-316.
- Li S, Xiao X, Sun P, Wang F (2008) Screening of genes regulated by cold shock in *Shewanella piezotolerans* WP3 and time course expression of cold-regulated genes. *Archives of Microbiology* **189**: 549-556.
- Li YJ & Chen ZC (2004) Molecular cloning and characterization of LCRG1, a novel gene localized to the tumor suppressor locus D17S800-D17S930. *Cancer Letters* **209**: 75-85.
- Liang P & Pardee AB (1992) Differential display of eukaryotic messenger RNA by means of the polymerase chain reaction. *Science* **257**: 967-971.
- Liao B-Y & Zhang J (2006) Low rates of expression profile divergence in highly expressed genes and tissue-specific genes during mammalian evolution. *Molecular and Biology Evolution* **23**: 1119-1128.
- Linney E & Udvadia AJ (2004) Construction and detection of fluorescent germline transgenic zebrafish. *Methods in Molecular Biology* **254**: 271-288.

- Lisitsyn N, Lisitsyn N, Wigler M (1993) Cloning the differences between two complex genomes. *Science* **259**: 946-951.
- Llado RR, Urwin P, Atkinson H, Gilmartin PM (1998) Hgm1, a novel male-specific gene from *Heterodera glycines* identified by differential mRNA display. *Molecular and Biochemical Parasitology* **95**: 45-52.
- Markovskaya EF, Sysoeva MI, Sherudilo EG, Topchieva LV (2007) Differential gene expression in cucumber plants in response to brief daily cold treatments. *Russian Journal of Plants Physiology* **54**: 607-611
- Martins ML & Yoshitoshi ER (2003) A new nematode species *Goezia leporinus* n. sp. (Anisakidae) from cultured freshwater fish *Leporinus macrocephalus* (Anostomidae) in Brazil. *Brazilian Journal of Biology* **63**: 497-506.
- Matz MV & Lukyanov SA (1998) Different strategies of differential display: areas of application. *Nucleic Acids Research* **26**: 5537-5543.
- Merlino G & Khanna C (2007) Fishing for the origins of cancer. *Genes Development* **21**: 1275-1279.
- Meyer JN, Volz DC, Freedman JH, Giulio RT (2005) Differential display of hepatic mRNA from killifish (*Fundulus heteroclitus*) inhabiting a superfund estuary. *Aquatic Toxicology* **73**: 327-341.
- Miller AN, Raghavan N, Fitz-Gerald P, Lewis FA, Knight M (2001) Differential gene expression in haemocytes of the snail *Biomphalaria glabrata*: effects of *Shistosoma mansoni* infection. *International Journal for Parasitology* **31**: 687-696.
- Minucci LV, Pinese JF, Espíndola ELG (2005) Análise limnológica de sistema semi-intensivo de criação de *Leporinus macrocephalus* (Pisces, Anostomidae). *Biosciences Journal* 21: 123-131.
- Mishra, RN, Reddy PS, Nair S, Markandeya G, Reddy AR, Sopory SK, Reddy MK (2007) Isolation and characterization of expressed sequence tags (ESTs) from subtracted cDNA libraries of *Pennisetum glaucum* seedlings. *Plant Molecular Biology* 64: 713-732.
- Morelli KA, Revaldaves E, Oliveira C, Foresti F (2007) Isolation and characterization of eight microsatellite loci in *Leporinus macrecephalus* (Characiformes: Anostomidae) and cross-species amplification. *Molecular Ecology Notes* 7: 32-34.

Nagae MY, Hayashi C, Soares CM, Furuya WM (2002) Inclusão do milheto (*Pennisetum americanum*) em rações para alevinos de piavuçu (*Leporinus macrocephalus*). *Revista Brasileira de Zootecnia* **31**: 1875-1880.

Nelson JS (1994) *Fishes of the world*. 3rd ed, John Wiley and Sons Inc., New York.

- Normandes EB, Barreto RE, Carvalho RF, Delicio HC (2006) Effects of Betaine on the growth of the fish piauçu, *Leporinus macrocephalus*. *Brazilian Archives of Biology and Technology* **49**: 757-762.
- Panula P, Sallinen V, Sundvik M, Kolehmainen J, Torkko V, Tiittula A, Moshnyakov M, Podlasz P (2006) Modulatory neurotransmitter systems and behavior: towards zebrafish models of neurodegenerative diseases. *Zebrafish* 3: 235-247.
- Parrington J & Coward K (2002) Use of emerging genomic and proteomic technologies in fish physiology. *Aquatic Living Resources* **15**: 193-196.
- Patton EE & Zon LI (2005) Taking human cancer genes to the fish: a transgenic model of melanoma in zebrafish. *Zebrafish* **1**: 363-368.
- Pinto LGQ, Pezzato LE, Miranda EC, Barros MM (2001) Desempenho do piauçu (*Leporinus macrocephalus*) arroçoado com dietas contendo diferentes teores de tanino. *Revista Brasileira de Zootecnia* **30**: 1164-1171.
- Ponsuksili S, Wimmers K, Schmoll F, Schellander K (1999) Application of the differential display technique for the detection of candidate genes. *Archives of Animal Breeding* **42**: 147-149.
- Poulin H & Labelle Y (2005) The PLAGL1 gene is down-regulated in human exrtaskeletal myxoid chondrosarcoma tumors. *Cancer Letters* **227**: 185-191.
- Queiroz JF, Lourenço JNP, Kitamura PC (2002) *A Embrapa e a aqüicultura. Demandas e prioridades de pesquisa*.Embrapa, Brasília.
- Ribeiro RIMA & Godinho HP (2003) Criopreservação do sêmen testicular do teleósteo piau-açu *Leporinus macrocephalus*. *Arquivo Brasileiro de Medicina Veterinária e Zootecnia* **55**: 1-7.
- Saunders HL & Magor BG (2004) Cloning and expression of the AID gene in the channel catfish. *Developmental and Comparative Immunology* **28**: 657-663.
- Sharma R, Sud N, Chattopadhyay TK, Ralhan R (2005) TC21/R-Ras2 upregulation in esophageal tumorigenesis: potential diagnostic implications. *Oncology* **69**: 10-18.

- Shena M, Shalon D, Davis R, Brown PO (1995) Quantitative monitoring of gene expression patterns with a complementary DNA microarray. *Science* **270**: 467-470.
- Shin G-T, Kim DR, Lim J-E, Yim H, Kim H (2008) Upregulation and function of GADD45^y in unilateral ureteral obstruction. *Kidney International* **73**: 1251-1265.
- Smartt CT & Erickson JS (2008) Bloodmeal-induced differential gene expression in the disease vector *Culex nigripalpus* (Diptera:Culicidae). *Journal of Medical Entomology* **45**: 326-330.
- Smith-Keume C & Dove S (2008) Gene expression of a green fluorescent protein homolog as a host-specific biomarker of heat stress within a reef-building coral. *Marine Biotechnology* **10**: 166-180.
- Soares CM, Hayashi C, Furuya VRB, Furuya WM, Galdioli EM (2000) Substituição parcial e total da proteína do farelo de soja pela proteína do farelo de canola na alimentação de alevinos de piavuçu (*Leporinus macrocephalus*, L.). *Revista Brasileira de Zootecnia* **29**: 15-22.
- Sola L & Gornung E (2001) Classical and molecular cytogenetics of the zebrafish, *Danio rerio* (Cyprinidae, Cypriniformes): an overview. *Genetica* **111**: 397-412.
- Sprague J, Clements D, Conlin T, Edwards P, Frazer K, Schaper K, Segerdell E, Song P, Sprunger B, Westerfield M (2003) The zebrafish information network (ZFIN): the zebrafish model organism database. *Nucleic Acids Research* **31**: 241-243.
- Sugimoto H & Endoh H (2008) Differentially expressed genes during fruiting body development in the aggregative ciliate *Sorogena stoianovitchae* (Ciliophora: Colpodea). *Journal of Eukaryotic Microbiology* **55**: 110-116.
- Sunagawa M & Magae Y (2005) Isolation of genes differentially expressed during the fruit body development of *Pleurotus ostreatus* by differential display of RAPD. *FEMS Microbiology Letters* **246**:279-284.
- Takahashi LS, Gonçalves FD, de Abreu JS, Martins MIEG, Ferreira ACM (2004)
 Economic viability of the piauçu *Leporinus macrocephalus* (Garavello & Britski, 1988) production. *Scientia Agricola* 61: 228-233.
- Torelli A, Marieschi M, Castagnoli B, Zanni C, Gorbi G, Corradi MG (2008) Identification of S₂-T A63: a cDNA fragment corresponding to a gene differentially expressed in a Cr-tolerant strain of the unicellular green alga *Scenedesmus acutus. Aquatic Toxicology* **86**: 495-507.

- Velculescu VE, Zhang L, Vogelstein B, Kinzler KW (1995) Serial analysis of gene expression. *Science* **270**: 484-487.
- Venere PC, Ferreira IA, Martins C, Galetti PM (2004) A novel ZZ/ZW sex chromosome system for the genus *Leporinus* (Pisces, Anostomidae, Characiformes). *Genetica* **121**: 75-80.
- Venkatesh B, Hettwer U, Koopmann, Karlovsky P (2005) Conversion of cDNA differential display results (DDRT-PCR) into quantitative transcription profiles. *BMC Genomics* **6**: 51-63.
- Welsh J, Chada K, Dalal SS, Cheng R, Ralph D, McClelland M (1992) Arbitrarily primed PCR fingerprinting of RNA. *Nucleic Acids Research* **20**: 4965-4970.
- Westerfield M (2000) The Zebrafish Book. A Guide for the Laboratory Use of Zebrafish Danio (Brachydanio) rerio. University of Oregon Press, Eugene.
- Wilkinson D (1998) *In situ hybridization: a practical approach*. 2nd edn. IRL Press, Oxford.
- Williams, DL, Sayed AA, Bernier J, Birkeland SR, Cipriano MJ, Papa AR, McArthur AG, Taft A, Vermeire JJ, Yoshino TP (2007) Profiling Schistosoma mansoni development using serial analysis of gene expression (SAGE). Experimental Parasitology 117: 246-258.
- Wong JS, Ye XQ, Muhlenkamp CR, Gill SS (2002) Effect of a peroxisome proliferator on 3 beta-hydroxysteroid dehydrogenase. *Biochemical and Biophysical Research Communications* **293**: 549-553.
- Wong QWL, Mak WY, Chu KH (2008) Differential gene expression in Hepatopancreas of the shrimp *Metapenaeus ensis* during ovarian maturation. *Marine Biotechnology* **10**: 91-98.
- Wu YL, Pan X, Mudumana SP, Wang H, Kee PW, Gong Z (2008) Development of a heat shock inducible *gfp* transgenic zebrafish line by using the zebrafish *hsp27* promoter. *Gene* **408**: 85-94.
- Xie J, Wen JJ, Chen B, Gui JF (2001) Differential gene expression in fully-grown oocytes between gynogenetic and gonochoristic crucian carps. *Gene* **271**: 109-116.
- Xu J, Belanger F, Huang B (2008) Differential gene expression in shoots and roots under heat stress for a geothermal and non-thermal *Agrostis grass* species contrasting in heat tolerance. *Environmental and Experimental Botany* **63**: 240-247.
- Zhang C, Wang JG, Liu GS, Chen QX (2005) Effect of the Chinese traditional medicine "Bushen Yinao Pian" on the cerebral gene expression of the senescence-accelerated mouse prone 8/Ta. *American Journal of Chinese Medicine* **33**: 639-650.
- Zhang J, Turley RB, Stewart JM (2008) Comparative analysis of gene expression between CMS-D8 restored plants and normal non-restoring fertile plants in cotton by differential display. *Plant Cell Reprodution* **27**: 553-561.
- Zhang JJ, He ZH, Tian H, Zhu GH, Peng XX (2007) Identification of aluminiumresponsive genes in rice cultivars with different aluminium sesitivities. *Journal of Experimental Botany* **58**: 2269-2278.
- Zon LI & Peterson RT (2005) In vivo drug discovery in the zebrafish. *Nature Reviews* **4**: 35-44.

Resultados e Discussão

Os resultados e a discussão dos dados obtidos encontram-se apresentados na forma de capítulos, referentes a trabalhos científicos. Visando facilitar a apresentação e compreensão de cada um destes capítulos, as figuras e tabelas foram numeradas em relação a cada trabalho específico. Da mesma forma, as referências bibliográficas citadas em cada capítulo não se encontram ao final do item "Introdução Geral" da tese e sim discriminadas no final de cada trabalho.

Capítulo 1: Molecular organization and expression analyses of skeletal actin genes from white and red fish muscle tissues: insights into the musculature energetic demands;

Capítulo 2: Identification of sexually dimorphic gene expression in brain tissue of the fish *Leporinus macrocephalus* through mRNA differential display and real time PCR analyses;

- **Capítulo 3:** Spatio-temporal distribution of fatty acid-binding protein 6 (*fabp6*) gene transcripts in the developing and adult zebrafish (*Danio rerio*);
- **Capítulo 4:** A comparative expression analysis of gene transcripts in brain tissue of wild-type and GH-transgenic zebrafish (*Danio rerio*) using a DDRT-PCR approach.

CAPÍTULO 1

Organização molecular e análises de expressão de genes de actina esquelética de tecidos musculares branco e vermelho de peixes: compreensão sobre a demanda energética muscular Molecular organization and expression analyses of skeletal actin genes from white and red fish muscle tissues: insights into the musculature energetic demands

Abstract

Two α -skeletal actin genes were isolated from the neotropical fish *Leporinus* macrocephalus, referring to white and red muscle tissues. Similar actin isoforms were also identified in other fish species, mainly differing by a Ser/Ala¹⁵⁵ substitution, which can have a functional significance related to actin-ATP interaction. A Ser¹⁵⁵, identified in the α -skeletal white muscle actin, seems to lead to a higher affinity to ATP molecules which, in turns, could be related to the fast contraction fibers of this tissue. However, an Ala¹⁵⁵, as observed in the α -skeletal actin from red muscle might result in a decrease in actin's affinity for ATP, which could also be associated to its tissue slow contractile performance. FISH mapped the skeletal muscle actin genes to a pair of submetacentric chromosomes. Comparisons on the predicted secondary protein structure evidenced no differences between the two analyzed actin isoforms, although a Phe/Ile²⁶² substitution at the red muscle actin leads to a hydrophobicity variation at the Dplug of the protein which could alter its stability. Data on gRT-PCR evidenced a significant gene expression variation between white and red muscle actin genes (T=105 Mann Whitney; p<0.001). Moreover, although no significant variation was observed between adult males and females (white muscle: females 0.01175±0.00247 vs. males 0.01082±0.004447; red muscle: females 0.000847±0.000613 vs. males 0.000754±0.000374), there was a great variation on gene expression within individuals of each sex. Available data on muscle actins

lead to the proposal that white and red α -skeletal actins are genetically and functionally distinguishable in fish species, a feature that is not found in other vertebrate groups.

Introduction

Muscles are a major source of energy consumption and heat generation and there has been a longstanding interest in the mechanisms that underlie muscle energetic properties. Skeletal muscle is found throughout the body and, in higher vertebrates, form an integrated network with a prominent skeletal system via tendons, resisting gravity and facilitating mobility. Lower vertebrates, particularly those that are aquatic, have a proportionally greater muscle mass (e.g. Sambasivan & Tajbakhsh 2007). In fish, body mass is mainly composed by skeletal musculature that constitutes 40-75% of the total weight of the animal. In most fish species, this tissue is composed by different muscle fibers that occupy distinct axial regions - a large portion of a deeper layer of white muscle (with fast contraction fibers and glycolytic metabolism, for energy supply), a superficial thin layer of red muscle (with slow contraction fibers and oxidative metabolism), and an intermediate layer between the red and white musculatures (with fibers of fast contraction and oxidative/glycolytic metabolism). While white muscle fibers are associated to fast swimming in fish, as predation and escape behaviors, the red muscle fibers are associated to slow movements, as migration and foraging habits (Driedzic & Hochachka 1976, Luther et al. 1995, Sänger & Stoiber 2001).

One of the major components of muscle tissues is actin that, together with myosin, tropomyosin, and troponin, represents a protein that composes the parallel

fibrils (miofibrils) of the striate musculature. Actin plays a crucial role in maintaining cytoskeletal structure, contractile processes, cell motility and division, and intracellular movements, and is essential to eukaryotic cells (Clarke & Spudich 1977, Lazarides & Revel 1979). More recently, it was evidenced that actin is also involved in diverse nuclear functions, including transcription, ribonucleoprotein packaging and transport, chromatin remodeling, and the formation of karyoskeletal elements (Pederson & Aebi 2002, Bettinger *et al.* 2004, Kiseleva *et al.* 2004). Reflecting the importance of its biological functions, the actin structure is highly conserved among different organisms.

Although the genes that code for actin proteins can occur in single copies, as observed in Sacharomyces cerevisae, some protozoans, and several green algae, actin proteins are encoded by a multigene family in animals, plants, and in many protozoans examined to date (Vandeckerckhove & Weber 1984, Hightower & Meagher 1986, Reece et al. 1992, Bhattacharya et al. 2000). Mammals have four muscle actin isoforms - two in striated muscles (α -skeletal and α -cardiac) and two in smooth muscles (γ -enteric and α -vascular) - and two nonmuscle or cytoplasmic types, the β - and γ -isoforms (e.g. Vandekerckhove & Weber 1984, Kusakabe et al. 1997, Mounier & Sparrow 1997). However, teleost fish seems to contain a higher number of distinct actin isoforms, as revealed by the most extent research on actin's diversity and tissue expression profile in fish genome performed to date on Takifugu rubripes, revealing nine isoforms (two askeletal, three α -cardiac, an α -anomalous testis type actin, two β -cytoplasmic, and one β -cytoplasmic vascular type-actin) (Venkatesh *et al.* 1996). Similarly, insects have at least six actin genes (Fyrberg et al. 1980). The actin gene family of plants is much larger, comprising 8-44 genes, depending on the taxa (Reece et al. 1992). Thus, the actin multigene family provides an excellent model for the study of the molecular mechanisms of gene regulation and expression and for insights into its evolution and function.

The multiple actin isoforms are expressed in different stages and/or tissues and seem to be encoded by a set of structurally related genes that were originated by gene duplication followed by functional divergence (Hightower & Meagher 1986, Miwa *et al.* 1991, Venkatesh *et al.* 1996). The controversy between the actin high structure conservation and several biophysical properties suggests that its several isoforms result from a very few amino acid substitutions in key functional positions. Small differences related to nucleotide or amino acid substitutions can constitute the structural base for changes in the conformation of the actin protein which, in turns, could modify its affinity to numerous bindings found in distinct cellular types (Ooi & Soematsu 2007).

Although there are several data on mammalian actins, the evolutionary origin, pattern of organization, and their diversity in lower vertebrates, especially fishes, remain to be better investigated. Moreover, data on the organization and expression of the main components of the muscle tissues can give new insights on the molecular events underlying muscle contraction and/or growth. Therefore, the purpose of the present study was the isolation and characterization of skeletal muscle actin genes of *Leporinus macrocephalus*, a fish species that represents an important food resource that has been intensively exploited in the subtropical aquaculture (Takahashi *et al.* 2004). The analyses resulted in the first description of white and red skeletal muscle actin genes that may be genetically and functionally distinguishable in fish species.

35

Material and Methods

Animal samples

Ten adult specimens (five males and five females) of *Leporinus macrocephalus* (piauçu) (Characiformes, Anostomidae) were obtained from a private fishery station (Piscicultura Água Milagrosa, Itajobi, São Paulo State, Brazil), where they were maintained under the same conditions. Animals with at least 40 days were selected since they already present gonadal differentiation (Toriyama 2001) which permits the sex identification under stereomicroscopy. White and red muscle samples from the dorsal region of the animals were collected and immediately stored at -80°C until RNA extraction. It was not possible to obtain muscle samples from the intermediate layer between the red and white musculature due to the difficulty to correctly isolate this tissue. Kidney tissue was also collected and immediately used for chromosome preparations.

RNA isolation

White and red muscle frozen samples were mechanically homogenized with TRizol Reagent (Invitrogen) and total RNA extraction followed the manufacturer's protocol. RNA samples were eluted in RNase-free water and quantified (NanoDrop 1000 Spectrophotometer) by measuring the optical density (OD) at 260nm. RNA purity was ensured by obtaining a 260/280nm OD ratio ≥1.80.

RT-PCR and cDNA amplification

Each RNA sample was reverse transcribed with the commercial kit SuperScript First-Strand Synthesis System for RT-PCR (Invitrogen) using an oligonucleotide (dT)₁₂₋₁₈ as an anchor primer, according to the manufacturer's instructions. cDNA (2 μ L) amplified forward AactF (5'was using AGGCCAACAGGGAGAAGATT-3') (5'and reverse AactR TCCATACCGATGAAGGAAGG-3') primers, designed based on alpha-actin gene sequence GenBank databases of the fish Takifugu rubripes (http://www.ncbi.nlm.nih.gov), in order to amplify a segment of this gene. Each PCR consisted of 2µL of cDNA, 0.2mM of each primer, 1x 25mM MgCl₂ PCR buffer, 0.2mM of dNTPs, and 1 unit of Platinum Tag DNA polimerase (Invitrogen), in a final volume of 25µL. PCR amplifications were carried out with an initial denaturation step at 95°C for 2 minutes, followed by 34 cycles at 95°C for 30 seconds, 55°C for 30 seconds, and 72°C for 1 minute, with an additional extension step at 72°C for 5 minutes. PCR products were fractionated on 1% agarose gel, stained with ethidium bromide, and visualized under UV light using Eagle Eye II image documentation system (Strategene).

Semi quantitative RT-PCR and statistical analysis

Four pools of RNA samples were prepared in order to minimize the detection of simple variations among *Leporinus macrocephalus* individuals (pool of RNA samples from white muscles of males; pool of RNA samples from white muscles of females; pool of RNA samples from red muscles of females; pool of RNA samples from red muscles of females; pool of RNA samples from red muscles of females; pool of RNA samples from red muscles of females; pool of RNA samples from red muscles of females; pool of RNA samples from red muscles of females; pool of RNA samples from red muscles of females; pool of RNA samples from red muscles of females; pool of RNA samples from red muscles of females; p

described in Marone et al. (2001) using 2µL of the obtained cDNA, and PCR was carried out at 95°C for 2 minutes, followed by 30 cycles with denaturation at 95°C (30 seconds), annealing at 55°C (30 seconds), and extension at 72°C (1 minute and 15 seconds), followed by an additional 5 minute extension step at 72°C. The PCR signals were normalized to the housekeeping 18S rRNA gene. A 245 bp segment of the 18S rRNA gene was amplified using the four RNA pools individually and the primers 18S1 (5'-TACCACATCCAAAGAAGGCAG-3') and 18S2 (5'-TCGATCCCGAGATCCAACTAC-3'), designed based on consensus sequences of this gene described for several fish species. Preliminary PCR experiments, using 28-30-32-34 cycles, were conducted to determine the number of cycles that represented the linear amplification range for analysis (Tom et al. 2004). PCR products were fractionated on 1% agarose gel, stained with ethidium bromide, and visualized under UV light using Eagle Eye II image documentation system (Strategene). The bands corresponding to each gene were quantified by densitometry as Integrated Optical Density (IOD) using Imaging Master VDS Software version 3.2 (GE Life Sciences).

Quantitative PCR and statistical analysis

Quantitative RT-PCR was performed using Power SYBER Green PCR Master Mix Kit (Applied Biosystems), according to the manufacturer's instructions. Standard reactions (25µL) were assembled using 12.5µL of Power SYBER Green PCR Master Mix 2x, 2µL of forward primer AactF2 (5µM) (5'-CCATCTATGAGGGTTACGCTCTTC-3'), 2µL of reverse primer AactR2 (5µM), (5'-CGGTTGTGACGAAAGAGTAGCC-3'), 2µL of template (10ng/µL), and 6.5µL of ultrapure water. Primers were designed with the software Primer Express v.2.0 (Applied Biosystems), based on the actin gene sequences previously obtained from white and red muscles of L. macrocephalus. Templates cDNA were 1:10 diluted, and cDNA samples were replaced by DEPC water in the negative controls. All real time assays were carried out in duplicate using an Applied Biosystem 7300 HT Real-Time PCR system (Applied Biosystems). Forty amplification cycles were performed and each cycle consisted of 94°C for 15 seconds followed by 60°C for 1 Amplification and dissociation curves generated by the 7300 minute. System/Sequence Detection Software version 4.0 (Applied Biosystems) were used for gene expression analysis. As performed for the semi-quantitative analysis, the qRT-PCR signals were normalized to a segment of the 18S rRNA housekeeping gene using the primers 18S3 (5'-CGGAATGAGCGTATCCTAAACC-3') and 18S4 (5'-GCTGCTGGCACC AGACTTG-3'), also designed based on consensus sequences of this gene described for several fish species. Following the removal of outliers, raw fluorescence data were exported to the online program QPCR Standard Curve Slope Efficiency Calculator to (http://www.stratagene.com/techtoolbox/calc/qpcr_slope_eff.aspx) to determine the PCR amplification efficiency. The Ct values were used to calculate a relative gene expression value for each transcript, according to 2^{-ΔCt} method (Livak & Schimittgen 2001). The relative gene expression values were submitted to statistical analysis using Student's unpaired t-test (p<0.05) to normal distribution, or Mann-Whitney T value (Mann-Wh) (p<0.05) to data which did not display a normal distribution (Zar 1999). Differences were considered significant when p<0.05 and a 95% confidence level of the difference was used.

Cloning, sequencing, and sequence analysis

PCR products were cloned into pGEM-T (Promega) vector and used to transform competent cells of the *E. coli* strain DH5 α (Invitrogen), following the manufacturers' instructions. Clones were submitted to automated sequencing on an ABI 377 Automated DNA Sequencer (Applied Biosystems) with a DYEnamic ET Terminator Cycle Sequencing kit (GE Healthcare Life Sciences), following the manufacturer's instructions, and using primers complementary to vector arms. Nucleic acid and amino acid sequence database searches were performed using BLAST/N (Altschul et al. 1990) at the National Center for Biotechnology (http://www.ncbi.nlm.nih.gov/blast). Information (NCBI) website Sequence alignments were obtained by Clustal-W function (Thompson et al. 1994) and the consensus sequences were manually determined. The deduced amino acid sequences were subjected to the bioinformatics computer program CLC Main Workbench 4.0.1 to generate protein secondary structure data and hydropathy plots using the Kyte-Doolittle scale with a window size default value of 9 (Kyte & Doolitle 1982).

Chromosome analyses

Mitotic chromosomes were obtained from a suspension of anterior kidney cells using direct preparations (Bertollo *et al.* 1978). A cDNA actin product (300-1.000ng), obtained using primers AactF and AactR, was isolated from agarose gels using a commercial kit (Sephaglas[™] BandPrep Kit, GE Healthcare Life Sciences). A SCOMP (single cell comparative genomic hybridization) methodology for DNA amplification was performed. Briefly, the cDNA was digested using 24U of *Msel* (New England Biolabs) at 37°C for 3 hours. In a separate tube, the adaptor-oligonucleotides LIB1 (5'-AGTGGGATTCCTGCTGTCAGT-3') and ddMsel (5'-

TAACTGACAGCdd-3') were incubated at 65°C to 15°C, using a slope of 1 min/°C. Further, 1µL of ATP (10mM) and 1µL of T4 DNA ligase (5U/µL) were added. The digested cDNA was mixed to this solution and incubated at 15°C overnight. The kit Elongase Amplification System (Invitrogen) was used for the primary DNA amplification, following the manufacturer's instructions. The amplified product was used as a template for a second amplification reaction, using the QIAquick PCR Amplification Kit (QIAGEN), following its instructions. The oligonucleotide LIB1 was removed by a Trul treatment at 65°C for 3 hours. The DNA probe (8µg) was further precipitated in the presence of the competitor Human Cot-1 DNA (50x) and salmon sperm DNA (50x), and labeled with biotin-11-dUTP by nick translation using the kit Bionick[™] Labeling System (Gibco), according to the manufacturer's instructions. After chromosome treatment with 10% pepsine for 5 minutes at 37°C, FISH (Fluorescent in situ hybridization) was performed as detailed in Wasko & Galetti (2000). Metaphases were examined with an Olympus **BX50** epifluorescence microscope.

Results and Discussion

The alpha-skeletal actin plays a key role in biological muscle movement and represents the major protein in muscles along with myosin. This protein has been intensively studied in mammals and its functions include polymerization in neutral salt and binding to Ca²⁺, Mg²⁺, adenine nucleotides, tropomyosin, and myosin (Estes *et al.* 1992, Reisler 1993). These multiple functions make it one of the structurally best-conserved cell proteins. Although muscles of most vertebrates contain one α -skeletal actin, the presence of two isoforms in the muscles of fish species has been reported. Venkatesh *et al.* (1996) identified two α -skeletal actins in the pufferfish *Takifugu rubripes*, which have 98.7% of identity at the amino acid sequence level. Two other fish species - *Coryphaenoides acrolepis* and *C. cinereus* - also present two distinct α -skeletal muscle actin isoforms (Morita 2000). However, the functional difference of these two actins in fish muscle was not elucidated.

Distinct muscle actin isoforms, with tissue and/or stage-specific patterns, could be related to the different structure of the muscle tissues of fish species. As so, the aim of the present study exploits the isolation and characterization of actin genes of white and red skeletal muscles of the neotropical fish species *Leporinus macrocephalus*, through the idea that it should be feasible to also find distinct α -skeketal actin genes, as already identified in other fish species. Moreover, it should be possible to identify the genetic bases that could be beyond the specific functions of different skeletal actin isoforms.

The amplification of the cDNA samples of *L. macrocephalus* from white and red muscle, using the two selected actin primers, revealed fragments of approximate sizes of 450 bp. Cloning and sequencing of the PCR products lead to the characterization of segments composed by 472 bp (Figure 1). Database

DDBJ/EMBL/GenBank searches for nucleotide and amino acid similarity index indicated that the isolated fragments of *L. macrocephalus* correspond to partial α -skeletal muscle actin genes, containing the complete regions of exons 2, 3, and 4, and a fragment of exon 5 (Figure 1). These results were confirmed through comparisons with data on vertebrate skeletal muscle actin genes that are composed by 6 exons, whose proteins present 375 amino acid residues (Venkatesh *et al.* 1996).

The results evidenced no nucleotide differences between the isolated actin genes from white muscles of males and females of *L. macrocephalus*. In the same manner, no differences could be detected between sexes when comparing the actin gene from red muscle. However, several base substitutions were identified when comparing the isolated α -skeletal muscle actin genes from white and red muscle tissues - 7 and 12 nucleotide substitutions were found in the exons 3 and 4, respectively (Figure 1). Despite these differences between the white and red muscle actin gene sequences of *L. macrocephalus*, their deduced amino acid sequences, referring to the residues 113 to 269, are very conserved (Figure 2), since most nucleotide variations correspond to synonymous substitutions and just three nucleotide differences are related to amino acid substitutions.

Available data on fish actins indicate the occurrence of two different skeletal muscle actin genes, firstly denominated α -*Sk1* and α -*Sk2* by Venkatesh *et al.* (1996). Comparisons with amino acid sequences of database banks indicated that the α -skeletal muscle actin gene isolated from white muscle of *L. macrocephalus* presents a higher identity with the actin gene type 2 identified in the fish species *Takifugu rubripes* (Venkatesh *et al.* 1996), *Coryphaenoides acrolepis*, and *C. cinereus* (Morita 2000). On the other hand, the characterized α -skeletal muscle

43

actin gene from red muscle of *L. macrocephalus* is more similar to the actin gene type 1 also characterized in these same other fish species.

The precursors of vertebrate muscle actins (muscle-like actins) are found only in urochordates and represent a transition from non-muscle like actins to vertebrate skeletal muscle actins (Kovilur *et al.* 1993). Actins have undergone a rapid diversification during the emergence of vertebrates to give rise to several muscle-type actins, as seen in fish species. Distinct α -skeletal muscle actin isoforms were not evidenced in other vertebrate groups. Therefore, it is possible that some of the actins that have been characterized in fish species are either specific to the teleost lineage or are yet to be identified in higher vertebrates.

The actin monomer structure, already characterized in several species, presents two domains, originally termed large and small, although they are now known to be nearly identical in size. The small domain comprehends the subdomain 1 (a.a. residues 1-32, 70-144, and 338-375) and the subdomain 2 (a.a residues 33-69), while the large domain comprises the subdomains 3 (a.a. residues 145-180 and 270-337) and 4 (a.a. residues 181-269) (Kabsch *et al.* 1990) (Figure 3a). The structure and rotations of these domains with respect to one another are important in many biological processes, such as enzyme catalysis, ligand binding, and oligomerization (Page *et al.* 1998). Amino acid substitutions could lead to conformational changes of the protein which in turn would modify chemical affinities for numerous ligands found inside the cells (Mounier & Sparrow 1997). The strong conservation of actin isoforms indicates a great restriction in the rate and nature of amino acid changes in this protein during evolution. Only substitutions compatible with the folded structure and protein function will have

been retained. As so, no extreme change on protein conformation is thus expected, but, rather, small and local modifications (Mounier & Sparrow 1997).

One of the actin ligands is the ATP molecule (Figure 3a), which is hydrolysed to ADP.Pi, playing a role in force generation by the actomyosin complex and accompanies assembly of actin monomers (G-actin) in to filaments (F-actin) (Chen *et al.* 2000). Amino acid residues 12-17 and 154-161 comprehend two protruding loops that form a hydrogen bond with the phosphate of ATP (Kabsch *et al.* 1990, Schüler *et al.* 1999, Morita 2000) since their amine nitrogen can be hydrogen bounded to oxygen atoms of the nucleotide phosphate tail (Schüler *et al.* 1999). This binding bridges the two loops between the small and the large domains, and prevents the unfolding of the protein and stabilizes its structure (Kabsch *et al.* 1990, Schüler *et al.* 1990). Any disturbance in the interdomain coupling is expected to result in a decreased stability.

An amino acid difference observed between the α -skeletal muscle actins from white muscle and from red muscle of *L. macrocephalus* corresponds to a substitution of a Serine (white muscle) for an Alanine (red muscle) at the residue 155 (Figure 2) that lies inside one of the ATP-binding site to the actin molecule, found at the subdomain 3. This amino acid substitution can influence the actin's affinity for ATP, a feature that has been described in actin isoforms of some vertebrate species, including fish, and in *Sacharomyces cervisae* (Table 1), an yeast that has one essential actin gene that encodes a protein that shares a high identity degree (87%) to vertebrate skeletal muscle actins (Geeves *et al.* 2005). Similarly, it has been evidenced that a Ser/Ala substitution at the amino acid 14, at the actin subdomain 1, results in a 40 to 60-fold decrease in actin's affinity for ATP (Chen *et al.* 1995) (Table 1), since Alanine has a lower hydrogen bonding capacity. Ser¹⁴ hydroxyl forms a polar bridge between the γ -phosphate ATP and the amide nitrogen of Gly⁷⁴, thus conferring additional stability on the actin small domain (Morita 2000, Ooi & Soematsu 2007). Therefore, the Ser/Ala¹⁵⁵ substitution found in the α -skeletal actin isoform of red muscle of *L. macrocephalus* could also influence the actin-ATP interaction and, consequently, alter this protein stability, structure, or function.

In fish, the presence of a Serine¹⁵⁵ has been also identified in an α -skeletal muscle actin type of some species, as Takifugu rubripes (Venkatesh et al. 1996), Coryphaenoides acrolepis and C. cinereus (Morita 2000), and in α -cardiac muscle actins, as identified in T. rubripes (Venkatesh et al. 1996), Oreochromis niloticus and O. mossambicus (Wasko et al. 2007), and Salmo trutta (Mudalige et al. 2007). On the other hand, Alanine was also identified as the 155th amino acid in a second a-skeletal muscle actin isoform in the fish Takifugu rubripes (Venkatesh et al. 1996), Coryphaenoides acrelepis, C. cinereus, and Cyprinus carpio (Morita 2000). Therefore, it is reasonable to suppose that the occurrence of a Serine or an Alanine in the vicinity of the actin ATP-binding site, at the amino acid residue 155, could be related to the type of contraction of the muscle fibers (fast vs. slow contraction), since force generation during muscle contraction can be understood in terms of cyclical length changes in segments of actin thin filaments moving trough the tree-dimensional lattice of myosin filaments (Schutt & Lindberg 1998, Sieck & Regnier 2001) and any disturbance in these proteins could alter the chemomechanical transduction (work, force, and shortening) in muscle.

Therefore, the residue Ser¹⁵⁵, that seems to lead to a higher affinity to ATP molecules, could be related to the contractile performance of white skeletal and cardiac muscle tissues (fast contraction fibers). On the other hand, the presence of

an Ala¹⁵⁵ might result in a decrease in actin's affinity for ATP, which could also be related to the slow contraction fibers of the red muscle tissue. Moreover, it is possible that the genetic base differences between white and red skeletal muscle actins of fish species may have been hidden since previous analyses probably have used both muscle tissues together and not individually, as performed in the present study. Therefore, some of the α -skeletal muscle actin isoforms already described for other fish species, as the denominated Sk1 actin of Takifuqu rubripes (Venkatesh et al. 1996), actin-1 of Coryphaenoides acrolepis and C. cinereus (Morita 2000), and the actin-2b of Coryphaenoides yaquinae and C. armatus (Morita 2003), that also present an Ala¹⁵⁵ may represent a red skeletal muscle actin type, while the other skeletal actin isoforms with a Ser¹⁵⁵ identified in these same fish species may be related to a white muscle type. As so, the amino acid 155 might be a diagnostic position in α -skeletal actins of red and white muscle tissues of fish species. In contrast, other vertebrates, as rabbit, chicken, mouse, human, and frog, synthesize a single α -skeletal actin isoform with Ser¹⁵⁵. This different pattern may reflect that fish skeletal musculature is layered organized and largely confined to locomotion demands rather than bundled as in the other vertebrates (Luther et al. 1995).

Comparisons between the two distinct α -skeletal muscle actin isoforms of *L. macrocephalus* also evidenced another amino acid substitution (Phe/IIe²⁶²) (Figure 2). Although there is no available data regarding to actin amino acid substitutions at this residue, analyses of the protein structure show that residues Phe²⁶² to Gly²⁷³ form an outer loop which buttresses both subdomains 3 and 4, and stabilizes the extensive intersubdomains interface (Page *et al.* 1998). Therefore, the identified Phe/IIe²⁶² amino acid variation could also influence the F-actin formation, nucleotide binding and exchange, and interactions with various actinbinding proteins.

The proposal that the different α -skeletal actin isoforms already described in some fish species may be correlated to white and red muscle tissues led us to predict their protein secondary structure, since amino acid differences can lead to a weak aggregation of these residues which could strongly influence the chain's motifs - as the alpha helices (coiled conformation in which every backbone N-H group donates a hydrogen bond to the backbone C=O group of the amino acid four residues earlier) and the beta strands (typically 5-10 amino acid long whose peptide backbones are almost fully extended) - and its folding (Kurgan 2008). As the present study characterized a partial region of two different α -skeletal actins of Leporinus macrocephalus, a protein diagram representation (Figure 3b) was achieved based on the complete amino acid sequence of the skeletal actin-1 and skeletal actin-2 described for Coryphaenoides acrolepis and C. cinereus (Morita 2000), since they present a identity level of 99.6% and 100% with the partial deduced actin amino acid sequences (residues 113-269) of the actin isoforms from red and white muscles of L. macrocephalus, respectively. Comparisons between the predicted secondary protein structures evidenced no differences between the two analyzed actin isoforms, since their alpha helices and beta strands configurations were identical, even at the 155th and 262th amino acid residues that evidence a Ser/Ala and a Phe/Ile substitution, respectively.

However, a comparison of the amino acids hydropathy of the two actin isoforms evidenced that a Serine instead of an Alanine as the 155th residue results in a small decrease in the protein hydrophobicity. Similarly, the presence of an Isoleucine instead of a Phenylalanine as the 262th amino acid also leads to a reduction in the actin hydrophobicity. Although the Ser¹⁵⁵ found in the white muscle actin isoform of *L. macrocephalus* probably not alter the protein stability or function, the lle²⁶² identified in the red muscle actin isoform of this fish species can alter the protein since amino acids 262 to 274 form an actin hydrophobic plug (H-plug) that is important for the globular actin monomer to polymerize reversibly into double-stranded actin filaments and also important for the protein stability (Shvetsov *et al.* 2008). Actin stability problems due to hydrophobicity variations have been shown to be related to amino acid substitutions along the 262-274 residues, even when these substitutions caused no significant changes in overall secondary structure (Kuang & Rubenstein 1997).

Genetic analyses of actins in fish species are mainly restricted to nucleotide and amino acid sequences descriptions and there is no data concerning the chromosome localization of these genes throughout *in situ* hybridization, probably due to the difficulty in obtaining a good resolution level, specially using PCR products directly or in plasmid vectors as hybridization probes. However, FISH mapping of actin genes could be a reasonably efficient and straightforward way of improving the resolution of comparative maps. A cDNA amplification product of *L. macrocephalus*, related to actin genes, was used as a probe in chromosome fluorescence in situ hybridization analyses. Preliminary results evidenced that the characterized α skeletal muscle actin genes were localized at the pericentromeric region of the short arm of a large submetacentric chromosome pair (Figure 4). No additional fluorescent signals were detected and generaly two signals were observed in interphase cells.

Data on vertebrate chromosome in situ localization of muscle actin genes were mostly obtained for human cells, indicating that the α -cardiac, α -skeletal, α vascular, β -cytoplasmic, and γ -cytoplasmic genes are located on chromosomes 15q14, 1q42.1, 10q23.3, 7p22, and 17q25, respectively (Ueyama et al. 1995, 1996). This distribution pattern of each actin gene to a particular chromosome pair seems to be common in other mammal species. However, actin loci in the genus Drosophila are dispersed over all chromosomal elements (Bondinas et al. 2002). Linkage mapping data on fish species, Danio rerio as (http://www.ncbi.nlm.nih.gov/genome/guide/zebrafish/index.html), also suggest that in this vertebrate group actin loci can be spread through several chromosomes. Therefore, even though the two different isolated α -skeletal actin genes of L. macrocephalus seems to be localized in a major group along a chromosome pair, other small actin cistrons might not be identified due to the FISH technical procedure.

Although no variations were identified in the nucleotide sequences of the actin genes isolated from males and females of *L. macrocephalus*, quantitative PCR analyses were performed in the α -skeletal actin genes of white and red muscle tissues from each sex in order to verify the eventual possibility of finding a gender-associated gene expression in this fish species. The occurrence of actin genes associated to a sex was already observed in some eukaryotes, as *Aedes aegypti* (Muñoz *et al.* 2004, Vyazunova & Lan 2004), *Schistosoma mansoni* (Davis *et al.* 1986), and *Schistosoma japonicum* (Fitzpatrick *et al.* 2004), a feature that seems to be associated to muscles that are required to sustain specific activities in each sex, as feeding, reproduction, or particular movements (Fitzpatrick *et al.* 2004).

The semi-quantitative analysis performed using RNA sample pools from white and red muscles of males and females of *L. macrocephalus* separately, evidenced an estimated actin mRNA level decrease in females compared to males (IOD ratio values of white muscle: males 0.675 vs. females 0.492; IOD ratio values

50

of red muscle: males 0.873 vs. females 0.642) (Figure 5). Further gene expression analyses were also performed, throughout Real-Time PCR, using RNA samples of white and red muscles of each male and each female separately, in order to confirm the previous semi-quantitative gene expression results. Surprisingly, individual results indicated that the characterized α -skeletal muscle actin genes had a higher relative expression in most females than in males (white muscle: females 0.01175±0.00247 vs. males 0.01082±0.004447; red muscle: females 0.000847±0.000613 vs. males 0.000754±0.000374) (Figure 6a, b). However, statistical analysis revealed no significant difference in gene expression between sexes, as values of t=-0.479, p=0.641, and t=-0.343, p=0.738 were obtained when comparing the α -skeletal actin genes from white and red muscles of males and females, respectively.

The apparently different results obtained through semi-quantitative and quantitative gene expression analyses could be related to the use of a mixture of RNA samples and the use of individual RNA samples in each of these two analyses, respectively. Moreover, although there was no significant difference when comparing all males and all females together, there was a relevant variation on gene expression when comparing individuals within each sex (Figure 6a, b). Although the biological significance of these differences is unknown, it may be related to the analysis of animals in different developmental stages, even these individuals were in adult phase and presented gonadal differentiation. As so, future gene expression analyses should be done in order to verify the occurrence of a variation in α -skeletal muscle actin gene expression throughout the developmental stages.

Gene expression comparisons between the two characterized α -skeletal muscle actin genes of *L. macrocephalus*, through Real-Time PCR, also revealed a significant higher actin mRNA level in white muscle when compared to red muscle (T=105 Mann-Wh; p<0.001) (Figure 6c). This finding could be related to the energetic demands of the white muscle tissue, with fast contraction fibers and glycolytic metabolism for energy supply (Driedzic & Hochachka 1976, Luther *et al.* 1995, Sänger & Stoiber 2001).

Due to its conserved and ubiquitous nature, actin genes are often used as internal controls in gene expression analyses based on data that they are expressed at a constant level across all samples. However, the present results and also data on β -actin genes of fish species (e.g. Tang *et al.* 2007) indicate that these constitutive genes can exhibit variable expression and should be used with careful as internal standards in order to normalize each particular set of experimental RNA samples.

Taken together, the present data propose that α -skeletal actins from white and red muscle tissues appear to be somewhat genetically and functionally distinguishable in fish species and should be under different evolutionary selective pressures, a feature that was not found in other vertebrate groups so far. Further analyses comparing the organization and expression pattern of distinct actin isoforms in several vertebrate species would be useful to better understand the molecular evolution and the function of these genes and to verify if the variable muscle actin isoforms of fish that are not found in other animal groups are either specific to this lineage or are yet to be identified in higher vertebrates.

52

Acknowledgements

The authors thank Mr. Odval S. Stradioti (Piscicultura Água Milagrosa, Itajobi, SP) for fish supply samples, and Dr. Maeli Dal Pai Silva, Dr. João Pessoa and Dr. Débora Colombi for technical assistance on the Real Time analyses. This work was supported by grants from FAPESP (Fundação da Amparo à Pesquisa do Estado de São Paulo) and CNPq (Conselho Nacional de Desenvolvimento Científico e Tecnológico).

References

- Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ (1990) Basic local alignment search tool. *Journal of Molecular Biology* **215**: 403-410.
- Aspenström P, Schutt CE, Lindberg U, Karlsson R (1993) Mutation in β-actin: influence on polymer formation and on interactions with myosin and profilin. *FEBS-Federation of European Biochemical Societies* **329**: 163-170.
- Bertollo LAC, Takahashi CS, Moreira Filho O (1978) Cytotaxonomic consideration on *Hoplias lacerdae* (Pisces, Erythrinidae). *Revista Brasileira de Genética* **1**: 103-120.
- Bettinger BT, Gilbert DM, Amberg DC (2004) Opinion Actin up in the nucleus. *Nature Review - Molecular Cell Biology* **5**: 410-415.
- Bhattacharya D, Aubry J, Twait EC, Jurk S (2000) Actin gene duplication and the evolution of morphological complexity in land plants. *Journal of Phycology* **36**: 813-820.
- Bondinas GP, Loukas MG, Goulielmos GN, Sperlich D (2002) The actin loci in the genus *Drosophila*: establishment of chromosomal homologies among five neartic species of the *Drosophila obscura* group by *in situ* hybridization. *Chromosoma* **111**: 256-266.

- Chen H, Bernstein BW, Bamburg JR (2000) Regulating actin-filament dynamics in vivo. *Trends in Biochemical Sciences* **25**: 19-23.
- Chen X & Rubenstein PA (1995) A mutation in an ATP-binding loop of *Saccharomyces cerevisiae* actin (S14A) causes a temperature-sensitive phenotype in-vivo and in-vitro. *Journal of Biological Chemistry* **270**: 11406-11414.
- Chen X, Peng J, Pedram M, Swenson CA, Rubenstein A (1995) The effect of the S14A mutation on the conformation and thermostability of Saccharomyces cerevisiae G-actin and its interactions with adenine nucleotides. The Journal of Biological Chemistry 270: 11415-11423.
- Clarke M & Spudich JA (1977) Non-muscle contractile proteins role of actin and myosin in cell motility and shape determination. *Annual Review of Biochemistry* **46**: 797-822.
- Davis AH, Blanton R, Rottman F, Maurer R, Mahmoud A (1986) Isolation of cDNA clones for differentially expressed genes of the human parasite *Schistosoma mansoni*. *Proceedings of the National Academy of Sciences of the United States of America* **83**: 5534-5538.
- Driedzic WR & Hochachka PW (1976) Control of energy metabolism in fish white muscle. *American Journal of Physiology* **230**: 579-582.
- Estes JE, Selden LA, Kinosian HJ, Gershman LC (1992) Tightly-bound divalent cation of actin. *Journal of Muscle Research and Cell Motility* **13**: 272-284.
- Fitzpatrick JM, Johansen MV, Johnston DA, Dunee DW, Hoffmann KF (2004) Gender-associated gene expression in two related strains of *Schistosoma japonicum*. *Molecular and Biochemical Parasitology* **136**: 191-210.
- Fyrberg EA, Kindle KL, Davidson N, Sodja A (1980) Actin genes of *Drosophila* dispersed multigene family. *Cell* **19**: 365-378.
- Geeves MA, Fedorov R, Manstein DJ (2005) Review Molecular mechanism of actomyosin-based motility. *Cell and Molecular Life Science* **62**: 1462-1477.
- Hightower RC & Meagher RB (1986) The molecular evolution of actin. *Genetics* **114**: 315-332.

- Kabsch W, Mannehrz HG, Suck D, Pai EF, Holmes KC (1990) Atomic structure of the actin DNase I complex. *Nature* 347: 37-44.
- Kisileva E, Drummond SP, Goldberg MW (2004) Actin- and protein-4.1 containing filaments link nuclear pore complexes to subnuclear organelles in *Xenopus* oocyte nuclei. *Journal of Cell Science* **117**: 2481-2490.
- Klein CA, Schmidt-Kittler O, Schardt JA, Pantel K, Speicher MR, Riethmuller G (1999) Comparative genomic hybridization, loss of heterozygosity, and DNA sequence analysis of single cells. *Proceedings of the Natural Academy of Sciences USA*: 4494-4499.
- Kovilur S, Jacobson JW, Beach RL, Jefferey WR, Tomlinson CR (1993) Evolution of the chordate muscle actin gene. *Journal of Molecular Evolution* **36**: 361-368.
- Kuang B & Rubenstein PA (1997) Beryllium fluoride and phalloidin restore polymerizability of a mutant yeast actin (V266G, L267G) with severely decreased hydrophobicity in a subdomain 3/4 loop. The American Society for Biochemistry and Molecular Biology 272: 1237-1247.
- Kurgan L (2008) On the relation between the predicted secondary structure and the protein size. *Protein Journal* **27**: 234-239.
- Kusakabe T, Araki I, Satoh N, Jeffery WR (1997) Evolution of chordate actin genes: evidence from genomic organization and amino acid sequences. *Journal of Molecular Evolution* **44**: 289-298.
- Kyte J & Doolittle R (1982) A simple method for displaying the hydropathic character of a protein. *Journal of Molecular Biology* **157**: 105-132.
- Lazarides D & Revel JP (1979) Molecular basis of cell movement. *Scientific American* **240**: 100-113.
- Livak KJ & Schmittgen TD (2001) Analysis of relative expression data using realtime quantitative PCR and the $2^{-\Delta\Delta C}$ Method. *Methods* **25**: 402-408.
- Luther PK, Munro PMG, Squire JM (1995) Muscle ultrastructure in the teleost fish. *Micron* **26**: 431-459.

- Marone M, Mozzetti S, Ritis D, Pierelli L, Scambia G (2001) Semiquantitative RT-PCR analysis to assess the expression levels of multiple transcripts from the same sample. *Biological Procedures Online* **3**: 19-25.
- Miwa T, Manabe Y, Kurokowa K, Kamada S, Kanda N, Bruns G, Ueyama H, Kakunaga T (1991) Structure, chromosome location, and expression of the human smooth muscle (enteric type) gamma-actin gene: evolution of six actin genes. *Molecular Cell Biology* **11**: 3296-3306.
- Morita T (2000) Amino acid sequences of α-skeletal muscle actin isoforms in two species of rattail fish, *Coryphaenoides acrolepis* and *Coryphaenoides cinereus*. *Fisheries Science* **66**: 1150-1157.
- Morita T (2003) Structure-based analysis of high pressure adaptation of α-actin. *The Journal of Biological Chemistry* **278**:28060-28066.
- Mounier N & Sparrow JC (1997) Structural comparisons of muscle and nonmuscle actins give insights into the evolution of their functional differences. *Journal of Molecular Evolution* **44**: 89-97.
- Mudalige WAKA, Jackman DM, Waddleton DM, Heeley DH (2007) A vertebrate slow skeletal muscle actin isoform. *FEBS-Federation of European Biochemical Societies* **274**: 3452-3461.
- Muñoz D, Jimenez A, Marinotti O, James AA (2004) The *AeAct-e* gene is expressed in the developing flight muscles of female *Aedes aegypti*. *Insect Molecular Biology* **13**: 563-568.
- Ooi A & Soematsu H (2007) Insight into nucleotide and Ca²⁺ binding to carp αactin. *Fisheries Science* **73**: 684-693.
- Page R, Lindberg U, Schutt CE (1998) Domain motions in actin. *Journal of Molecular Biology* **280**: 463-474.
- Pederson T & Aebi U (2002) Actin in the nucleus: what form and what for? *Journal* of Structural Biology **140**: 3-9.
- Reece KS, McElroy D, Wu R (1992) Function and evolution of actins. *Evolutionary Biology* **26**: 1-34.

- Reisler E (1993) Actin molecular structure and function. *Current Opinion in Cell Biology* **5**: 41-47.
- Sambasivan R & Tajbakhsh S (2007) Skeletal muscle stem cell birth and properties. *Seminars in Cell and Developmental Biology* **18**: 870-882.
- Sänger AM & Stoiber W (2001) Muscle fiber diversity and plasticity. In: Muscle development and growth. Johnston IA (ed.). Academic Press, San Diego, USA. p. 187-250.
- Schüler H, Korenbaum E, Schutt CE, Lindberg U, Karlsson R (1999) Mutational analysis of Ser14 and Asp157 in the nucleotide-binding site of β-actin. *European Journal of Biochemistry* **265**: 210-220.
- Schüler H, Schutt CE, Lindberg U, Karlsson R (2000) Covalent binding of ATP γ S to the nucleotide-binding site in S14C-actin. *FEBS Letters* **476**: 155-159.
- Schutt CE & Lindberg U (1998) Muscle contraction as a Markov process I: energetics of the process. *Acta Physiologica Scandinavia***163**: 307-323.
- Shvetsov A, Galkin VE, Orlova A, Phillips M, Bergeron SE, Rubenstein PA, Egelman EH, Reisler E (2008) Actin hydrophobic loop 262-274 and filament nucleation and elongation. *Journal of Molecular Biology* **375**: 793-801.
- Sieck GC & Regnier M (2001) Plasticity and energetic demands of contraction in skeletal and cardiac muscle. *Journal of Applied Physiology* **90**: 1158-1164.
- Strzelecka-Golaszewska H, Venyaminov SY, Zmorzynski S, Mossakowska M (1985) Effects of various amino acid replacements on the conformational stability of G-actin. *European Journal of Biochemistry* **147**: 331-342.
- Takahashi LS, Gonçalves FD, de Abreu JS, Martins MIEG, Ferreira ACM (2004) Economic viability of the piauçu *Leporinus macrocephalus* (Garavello & Britski, 1988) production. *Scientia Agricola* **61**: 228-233.
- Tang R, Dodd A, Lai D, McNabb W, Love DR (2007) Validation of zebrafish (*Danio rerio*) reference genes for quantitative Real-time RT-PCR normalization. *Acta Biochimica et Biophysica Sinica* **39**: 384-390.
- Thompson JD, Higgins DG, Gibson TJ (1994) Clustal W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting,

position-specific gap penalties and weight matrix choice. *Nucleic Acids Research* **22**: 4673-4680.

- Tom M, Chen Nir, Segev M, Herut B, Rinkevich B (2004) Quantifying fish metallothionein transcript by real time PCR for its utilization as an environmental biomarker. *Marine Pollution Bulletin* **48**: 705-710.
- Toriyama E (2001) Reversão de sexo e análise do crescimento em piauçu (*Leporinus macrocephalus*). Master Thesis. Universidade Estadual Paulista Júlio de Mesquita Filho, Botucatu, SP.
- Ueyama H, Inazawa J, Ariyama T, Nishino H, Ochiai Y, Ohkubo I, Miwa T (1995) Reexamination of chromosomal loci of human muscle actin genes by fluorescence in-situ hybridization. *Japanese Journal of Human Genetics* **40**: 145-148.
- Ueyama H, Inazawa J, Nishino H, Ohkubo I, Miwa T (1996) FISH localization of human cytoplasmic actin genes ACTB to 7p22 and ACTG 1 to 17q25 and characterization of related pseudogenes. *Cytogenetic and Cell Genetics* **74**: 221-224.
- Vandekerckhove J & Weber K (1984) Chordate muscle actins differ distinctly from invertebrate muscle actins: the evolution of the different vertebrate muscle actins. *Journal of Molecular Biology* **179**: 391-413.
- Venkatesh B, Tay BH, Elgar G, Brenner S (1996) Isolation, characterization and evolution of nine pufferfish (*Fugu rubripes*) actin genes. *Journal of Molecular Biology* 259:655-665.
- Vyazunova I & Lan Q (2004) Stage-specific expression of two actin genes in the yellow fever mosquito, *Aedes aegypti*. *Insect Molecular Biology* **13**: 241-249.
- Wasko AP & Galetti PM (2000) Mapping 18S ribosomal genes in fish of the genus Brycon (Characidae) by fluorescence in situ hybridization (FISH). Genetics and Molecular Biology 23: 135-138.
- Wasko AP, Severino FE, Presti FT, Poletto AB, Martins C (2007) Partial molecular characterization of the Nile tilapia (*Oreochromis niloticus*) alpha-cardiac muscle

actin gene and its relationship to actin isoforms of other fish species. *Genetics and Molecular Biology* **30**: 1089-1092.

- Wen KK & Rubenstein PA (2003) Biochemical consequences of the cardiofunk (R177H) mutation in yeast actin. *Journal of Biological Chemistry* 278: 48386-48394.
- Wong WW, Doyle TC, Choung P, Olson TM, Reisler E (2001) Functional studies of yeast actin mutants corresponding to human cardiomyopathy mutations. *Journal of Muscle Research and Cell Motility* **22**: 665-674.

Zar JH (1999) *Biostatistical analysis*. Prentice-Hall, New Jersey, pp.663.

Table, Figures and Legends

Table 1: Actin amino acid residues as adenosine triphosphate binding sites and actin's affinity for ATP related to some actin isoforms of distinct organisms. The vertebrate data were based on comparisons with the chicken α -skeletal muscle actin, and the yeast data were based on *Saccharomyces cerevisae* wild type actin, which were used as controls.

Actin isoform	Amino acid residue	ATP binding	Reference
carp (<i>Cyprinus carpio</i>) α-skeletal muscle actin *	a.a. 155 Ser \rightarrow Ala	weak	Ooi & Soematsu (2007)
brown trout (Salmo trutta) α-skeletal muscle actin	a.a. 155 Ser \rightarrow Ala	weak	Mudalige et al. (2007)
Atlantic salmon (Salmo salar) α-skeletal muscle actin	a.a. 155 Ser	same as control	Mudalige et al. (2007)
Pacific grenadier (<i>Coryphaenoides acrolepis</i>) α-skeletal muscle actin-1	a.a. 155 Ser \rightarrow Ala	weak	Morita (2003)
Pacific grenadier (<i>Coryphaenoides. acrolepis</i>) α-skeletal muscle actin-2a	a.a. 155 Ser	same as control	Morita (2003)
Loosescale grenadier (Coryphaenoides yaquinae) α-skeletal muscle	a.a. 155 Ser \rightarrow Ala	weak	Morita (2003)
actin-2b			
Abyssal grenadier (Coryphaenoides armatus) α-skeletal muscle actin-2b	a.a. 155 Ser \rightarrow Ala	weak	Morita (2003)
Saccharomyces cerevisae actin *	a.a. 14 Ser \rightarrow Ala	weak	Chen <i>et al.</i> (1995); Chen & Rubenstein (1995)
Saccharomyces cerevisae actin *	a.a. 157 Asp \rightarrow Ala	same as control	Schüler <i>et al.</i> (1999)
Saccharomyces cerevisae actin *	a.a 177 Arg → Asp	same as control	Schüler <i>et al.</i> (2000)
Saccharomyces cerevisae actin * [†]	a.a. 177 Arg \rightarrow His	same as control	Wen & Rubenstein (2003)
Saccharomyces cerevisae actin *	a.a. 38 Pro \rightarrow Ala	same as control	Aspenström <i>et al.</i> (1993)
Saccharomyces cerevisae actin *	a.a. 374 Cys → Ser	same as control	Aspenström <i>et al.</i> (1993)
chicken smooth actin	a.a. 17 Val \rightarrow Cys	weak	Strezelecka-Gotlaszewska <i>et al.</i> (1985)
	a.a. 89 Thr \rightarrow Ser		
bovine smooth actin	a.a. 17 Val \rightarrow Cys	weak	Strezelecka-Gotlaszewska <i>et al.</i> (1985)
	a.a. 89 Thr \rightarrow Ser		

* refer to induced actin amino acid substitutions

[†] equivalent to the not induced zebrafish (*Danio rerio*) cardiofunk actin mutation and to the not induced human cardiomyopathy mutation that cause abnormal heart development (Wong *et al.* 2001)

	r exon 2
WM	AAGGCCAACAGGGAGAAGATTACCCAGATCATGTTCGAGACCTTCAACGTCCCTGCCATG
RM	AAGGCCAACAGGGAGAAGATTACCCAGATCATGTTCGAGACCTTCAACGTCCCTGCCATG
	exon 2 - exon 3
WM	TATGTGGCCATCCAGGCTGTCCTGTCCCTGTACGCCTCCGGCCGTACCACTGGTATTGTG
RM	TATGTGGCCATCCAGGCTGTCCTGTCCCTGTACGCCTCCGGCCGTACCACTGGTATTGTG
WM	CTGAGGTCCGGTGATGGTGTGACCCACAACGTCCCCATCTATGAGGGTTACGCTCTTCCC
RM	CTGCGGTCCGGTGATGGTGTGACCCACAACGTCCCCATCTATGAGGGTTACGCCCTTCCC
WM	CACGCCATCATGCGTCTGGATCTGGCCGGTCGCGATCTGACTGA
RM	CATGCCATCATGCGTCTGGATCTGGCTGGTCGCGATCTGACTGA
	exon 3 ++> exon 4
WM	CTGACTGAGAGAGGCTACTCTTTCGTCACAACCGCTGAGCGTGAGATCGTGCGTG
RM	CTGACTGAGCGTGGCTACTCTTTCGTCACAACTCCCGAGCGTGAGATTGTGCGCGACATC
WM	AAGGAGAAGCTGTGCTATGTGGCTCTGGACTTTGAGAACGAGATGGCCACCGCTGCCTCC
RM	AAGGAGAAGCTGTGCTATGTGGCTCTGGACTTTGAGAACGAGATGGCCACCGCTGCCTCC
WM	TCCTCCTCCCTGGAGAAGAGCTACGAGCTTCCCGATGGTCAGGTCATCACCATTGGTAAC
RM	TCCTCCTCCCTGGAGAAGAGCTACGAGTTGCCTGACGGCCAGGTCATCACAATTGGTAAC
WM	
RM	GAGCGTTTCCGCTGCCCCAGAGACCCCTGAACCAGCCTTCCTT
1.110	

Figure 1: Consensus nucleotide sequences of partial á-skeletal muscle actin genes isolated from white muscle (WM) and red muscle (RM) of *Leporinus macrocephalus*. Red letters indicate nucleotide variations. Regions of beginning and ending of the exons are indicated by arrows.

 113
 155
 167

 KANREKMTQIMFETFNVPAMYVAIQAVLSLYASGRTTGIVLDSGDGVTHNVPIYE
 222

 WM
 GYALPHAIMRLDLAGRDLTDYLMKILTERGYSFVTTAEREIVRDIKEKLCYVALD

 WM
 EENEMATAASSSSLEKSYELPDGQVITIGNERFRCPETLFQPSFIGM

 RM
I....

Figure 2: Deduced amino acid sequences (residues 113-269) of the partial α -actin skeletal muscle genes isolated from white muscle (WM) and red muscle (RM) of *Leporinus macrocephalus*. Dots indicate identical amino acid residues. The 155th and the 262th residues are indicated by bold letters.



Figure 3: Actin monomer (G-actin) tridimensional structure. **(a)** the small subdomain (composed of subdomains 1 and 2) and the large subdomain (composed of subdomains 3 and 4), and the ATP-binding site at the subdomain 3. The N and C termini are located in subdomain 1. **(b)** predicted secondary structure of the α -skeletal muscle actin-2 of *Coryphaenoides acrolepis* and *C. cinereus* (based on amino acid data described by Morita 2000), evidencing the alpha helices and the beta strands configurations.


Figure 4: *In situ* hybridization results of the actin genes in metaphase chromosomes of *Leporinus macrocephalus*. Arrows indicate the actin loci. Scale bar = 3μ m.



Figure 5: Alpha-skeletal muscle actin representative semi-quantitative RT-PCR results from white muscle (WM) and red muscle (RM) of males (M) and females (F) of *Leporinus macrocephalus*. 1% agarose gel evidencing the expression of the constitutive 18S rRNA gene used as control (a), and the expression of the α -skeletal muscle actin genes of white and red muscle tissues (b), statistical data on gene expression levels of the α -skeletal muscle actin genes of white and red muscle tissues (c). Quantification of PCR signals was obtained by densitometric analysis of the product as Integrated Optical Density (IOD). L, 1Kb DNA molecular marker.



Figure 6: Relative gene expression analysis (Real-Time PCR) of the α -skeletal muscle actin genes of white and red (c) muscle tissues of males and females (a, b) of *Leporinus macrocephalus* relative to the constitutive 18S rRNA gene (18S) used as control. Data were run in duplicate. Normalized data were expressed as means \pm SD.

CAPÍTULO 2:

Identificação de expressão gênica dimórfica relacionada ao sexo em tecido cerebral da espécie de peixe *Leporinus macrocephalus* através de análises de *display* diferencial e PCR em tempo real Identification of sexually dimorphic gene expression in brain tissue of the fish *Leporinus macrocephalus* through mRNA differential display and real time PCR analyses

Abstract

Differentially expressed genes in males and females of vertebrate species generally have been investigated in gonads and, to a lesser extent, in other tissues. Therefore, we attempted to identify sexually dimorphic gene expression in the brains of adult males and females of Leporinus macrocephalus, a gonochoristic fish species that presents a ZZ/ZW sex determination system, throughout a comparative analysis using differential display reverse transcriptase-PCR and real-time PCR. Four cDNA fragments were characterized, representing candidate genes with differential expression between the samples. Two of these fragments presented no significant identity with previously reported gene sequences. The other two fragments, isolated from male specimens, were associated to the gene that codes for the protein APBA2 (amyloid beta (A4) precursor proteinbinding, family A, member 2) and to the Rab 37 gene, a member of the Ras oncogene family. The overexpression of these genes has been associated to a greater production of the β -amyloid protein which, in turns, is the major factor that leads to Alzheimer's disease, and to the development of brain-tumors, respectively. Quantitative RT-PCR analyses revealed a higher Apba2 gene expression in males, thus validating the previous data on differential display. L. macrocephalus may represent an interesting animal model to the understanding of the function of several vertebrate genes, including those involved in neurodegenerative and cancer diseases.

Introduction

The development of the sexual phenotype in several vertebrates involves a complex cascade of events, initiated at conception by the fusion of two gametes and involving the action of regulatory sex determination genes - often in a response to a karyotype signal - and ending in the production of sex-specific molecules, anatomy, and behaviors (Marin & Baker 1999, Eriksson *et al.* 1999). Sex determination pathways are highly variable among different vertebrate groups, although they often include the presence or absence of a testis determining factor (SRY), leading to the formation of ovaries or testis, which in turns determine several sex-specific expression of gene networks in somatic tissues (Zarkower 2001, Wilhelm *et al.* 2007). This lack of conservation is intriguing itself, but means that the pathways of gender-specific genes must be dissected at a species level.

Mammals and birds present a relative stability in the sexual determination process, due to the presence of a XX/XY or a ZZ/ZW sex chromosome system, respectively. However, the presence and role of sex chromosomes in sex determination and differentiation might not be always clear in all vertebrates, particularly in fish species. Teleost fish can have different sex determination systems, including chromosomes that are not morphologically differentiated between males and females and sex chromosomes that are completely different between the two sexes. Moreover, males (XY) or females (ZW) can represent the heterogametic sex in different species. It has been also reported that some teleost fish have sex determination systems composed by multiple chromosomes (Tave 1993, Moreira-Filho *et al.* 1993, Almeida-Toledo & Foresti 2001). Despite this, the sexual differentiation in fish can also be determined by environmental factors and,

depending on the species, different hermaphroditic and gonochoristic modes of reproduction can occur (Yamamoto 1969, Conover *et al.* 1992).

The chromosomal, gonads, and behavioral differences observed between males and females of vertebrate species represent the major factors that lead to several efforts in the isolation and characterization of genes that are differentially expressed, especially in gonads. In fish, sex-specific genes have been isolated from male and females gonads generally in order to understand the sexual determination, and also sexual reversion, hermaphroditism, and unisexuality processes (Liu *et al.* 1996, Kanamori 2000, Guan *et al.* 2000, D'Cotta *et al.* 2001, Denslow *et al.* 2001a, b, Xie *et al.* 2001, Matsuda *et al.* 2002, Kondo *et al.* 2003, Lutfalla *et al.* 2003).

Despite the relative extensive data on genes with differential expression in vertebrate's testis and ovaries, data on sexually differential expressed genes in other organs are not so frequent, probably because these transcripts seem to be less common in somatic tissues (Santos *et al.* 2008). Although gender-associated genes have been described in heart, liver, kidney, adipose tissue, and muscles of some vertebrate species (Hutchison 1997, Mayer *et al.* 1998, Eriksson *et al.* 2008), Grossmann *et al.* 2004, Rinn & Snyder 2005, Yang *et al.* 2006, Isensee *et al.* 2008), data on sexually dimorphic genes in brain are still scarce - a well characterization of these transcripts was only performed in mouse (Eriksson *et al.* 1999, Xu *et al.* 2001, Yang *et al.* 2006, Yuge *et al.* 2007), specially due to the brain structure dimorphism between males and females. However, brain represents an extremely interesting tissue in order to analyze sexually dimorphic gene expression. Hundreds of sex differences have been identified in human and animal brains, ranging from structure and behaviors to molecules. Many of these differences have been

proven to be due to hormones and its metabolites, acting in the developing brain and permanently wiring it in a sex specific fashion (McCarthy & Konkle 2005, Becker *et al.* 2005, Morris *et al.* 2004, Xu & Disteche 2006).

Gene expression differences in brain associated to males and females have only been described in few fish species, as *Onchorhynchos nerka* (Jadhao *et al.* 2001), *Danio rerio* (Goto-Kazeto *et al.* 2004, Kallivretaki *et al.* 2007, Santos *et al.* 2008), *Fundulus heteroclitus* (Lauer *et al.* 2006), *Pimephales promelas* (Filby & Tyler 2007), *Oreochromis niloticus* (Davis *et al.* 2008), and *Dicentrarchus labrax* (Moles *et al.* 2007). There is no data on differentially expressed genes on brain tissues of neotropical fish species. Analyses in the brain of lower vertebrates represent an important source for the identification and characterization of target genes that could be related to differences in neurophysiology between males and females and for the understanding of the molecular pathways that are responsible for these differences.

In an attempt to isolate sexually dimorphic genes expressed in adult fish brain we used a DDRT-PCR approach for mRNA screening in *Leporinus macrocephalus* (piauçu), a gonochoristic fish species that represents an important food resource that has been intensively exploited in the subtropical aquaculture (Takahashi *et al.* 2004) and that presents a *ZZ/ZW* sex determination system (Galetti & Foresti 1987). Further quantitative PCR analysis, through real-time PCR, permitted to identify the expression levels of the isolated transcripts.

Material and Methods

Animal samples

Adult specimens of *Leporinus macrocephalus* (Characiformes, Anostomidae) were obtained from a private fishery station in São Paulo State, Brazil (Piscicultura Kabeya - municipality of Glicério), where they were maintained under the same conditions. Animals with at least 40 days were selected since they already present gonadal differentiation (Toriyama 2001) which permits the sex identification under stereomicroscopy. Brain tissue samples (including the medulla, cerebellum, optic lobes, pineal gland, pituitary gland, cerebral hemispheres, and olfactory lobes) of the animals were collected and immediately stored at -80°C until RNA extraction.

RNA isolation

Approximately 100mg of the brain samples were mechanically homogenized with 1mL of TRizol Reagent (Invitrogen) and total RNA extraction followed the manufacturer's protocol. RNA samples were eluted in RNase-free water and quantified (NanoDrop 1000 Spectrophotometer) by measuring the optical density (OD) at 260nm. RNA purity was ensured by obtaining a 260/280nm OD ratio ≥1.80.

DDRT-PCR

Approximately 2µg of five RNA samples from males and females were used to prepare RNA pools of each sex which were reverse transcribed (RT-PCR), separately, with the commercial kit SuperScript First-Strand Synthesis System for RT-PCR (Invitrogen) using an oligonucleotide (dT)₁₂₋₁₈ as an anchor primer, according to the manufacturer's instructions. The cDNA samples were amplified using as single primers some oligonucleotides with 15-24 bp that were designed based on VNTR core sequences (Table 1), and also RAPD (Randomly Amplified Polymorphic DNA) decamer primers (OPP series - Operon Technologies Inc.) (Table 2). Each cDNA amplification reaction consisted of $2\mu L$ of cDNA (10%), 0.2mM of primer, 1x 25mM MgCl₂ PCR buffer, 0.2mM of dNTPs, and 0.2 unit of Platinum Tag DNA polymerase (Invitrogen), in a final volume of 50µL. Reactions using the 15-24 bp primers were carried out with an initial denaturation step at 94°C for 2 minutes, followed by 40 cycles at 94°C for 50 seconds, 55°C for 2 minutes, and 72°C for 50 seconds, with an additional extension step at 72°C for 5 minutes. Reactions using the 10-mer primers were carried out with an initial denaturation step at 95°C for 5 minutes, followed by 45 cycles at 94°C for 1 minute, 36°C for 1 minute, and 72°C for 2 minutes, following the manufacturer's instructions of the Kit Ready-To-Go RAPD Analysis Beads (GE Healthcare Biosciences). DDRT-PCR products (10µL) were fractionated on 6% polyacrilamide gel, stained with 0.17% silver nitrate (AgNO₃) (Sambrook & Russel 2001), and visualized under white light. The molecular weigh of the amplified fragments was assigned through comparison with 1Kb DNA ladder (Invitrogen).

DNA isolation and reamplification

Selected DNA fragments that seemed to correspond to genes with differential expression between the analyzed samples were reamplified in order to obtain a higher DNA content to be used on cloning and nucleotide sequencing. The fragments were cut from the polyacrilamide matrix and purified using a QIAquick® Gel Extraction Kit (Qiagen), following the manufacturer's instructions. The purified fragments were used on reamplification reactions, as described in Dakis & Kouretas (2002). The amplification reactions followed the conditions described above. PCR products were analyzed through 6% polyacrilamide gel electrophoresis.

Cloning, sequencing, and sequence analysis

Products of reamplification were cloned into pGEM-T (Promega) vector and used to transform competent cells of the *E. coli* strain DH5α (Invitrogen). Clones were submitted to automated sequencing on an ABI 377 Automated DNA Sequencer (Applied Biosystems) with a DYEnamic ET Terminator Cycle Sequencing kit (GE Healthcare Life Sciences), following the manufacturer's instructions, and using primers complementary to vector arms. Nucleic acid sequence database searches were performed using BLAST/N (Altschul *et al.* 1990) at the National Center for Biotechnology Information (NCBI) website (http://www.ncbi.nlm.nih.gov/blast). Sequence alignments were obtained by Clustal-W function (Thompson *et al.* 1994) and the consensus sequences were manually determined.

Real-time PCR and statistical analysis

Quantitative RT-PCR was performed using Power SYBER Green PCR Master Mix Kit (Applied Biosystems), according to the manufacturer's instructions. Standard reactions (25µL) were assembled using 12.5µL of Power SYBER Green PCR Master Mix 2x, 2µL of each primer (5µM), 2µL of template cDNA treated with DNase I (Invitrogen) (10ng/µL), and 6.5µL of ultrapure water. Primers were designed with the software Primer Express v.2.0 (Applied Biosystems), based on the DNA sequences previously obtained (Table 3). Templates cDNA were 1:10 diluted, and cDNA samples were replaced by DEPC water in the negative controls. All real time assays were carried out in duplicate using an Applied Biosystem 7300 HT Real-Time PCR system (Applied Biosystems). Forty amplification cycles were performed and each cycle consisted of 94°C for 15 seconds followed by 60°C for 1 minute. Amplification and dissociation curves generated by the 7300 System/Sequence Detection Software version 4.0 (Applied Biosystems) were used for gene expression analysis. The qRT-PCR signals were normalized to a segment of the 18S rRNA housekeeping gene using the primers 18S3 (5'-CGGAATGAGCGTATCCTAAACC-3') 18S4 (5'and GCTGCTGGCACCAGACTTG-3'), also designed based on consensus sequences of this gene described for several fish species. Following the removal of outliers, raw fluorescence data were exported to the online program QPCR Standard Curve Slope to Efficiency Calculator (http://www.stratagene.com/techtoolbox/calc/qpcr slope eff.aspx) to determine the PCR amplification efficiency. The Ct values were used to calculate a relative gene expression value for each transcript, according to 2-4Ct method (Livak & Schimittgen 2001). The relative gene expression values were submitted to statistical analysis using Student's unpaired t-test (p<0.05) to normal distribution, or Mann-Whitney T value (Mann-Wh) (p<0.05) to data which did not display a normal distribution (Zar 1999). Differences were considered significant when p<0.05 and a 95% confidence level of the difference was used.

76

Results and Discussion

Polymerase chain reaction (PCR)-based mRNA differential display has been widely used for identifying differentially expressed transcripts in a variety of species (e.g. Liang 2002). The choice of applying a DDRT-PCR approach in Leporinus macrocephalus in order to identify genes with different expression levels in males and females was mainly due to the principle that this methodology does not need a previous knowledge on the mRNA sequences of the target biological samples. Moreover, the differential display (DD) technique can simultaneously visualize increased or decreased expression of numerous mRNAs from many samples, and requires relatively small amounts of starting material (Pardee & McClelland 1999). Although DD also has some disadvantages, as the possibility of isolation "false-positive" transcripts (PCR products that appear to be differentially expressed on the gel, but that cannot be verified when subsequent expression analyses are performed) (McClelland et al. 1995, Liang 1998, Pardee & McClelland 1999), technical improvements have been increasing its successful application. Modifications which have been proposed to improve differential display include the use of primers tailored to amplify members of a particular gene family (Jurecic 1996), the optimization of annealing temperature (Malhotra et al. 1998), and optimization of primer design (Graf et al. 1997).

In order to improve the isolation of differentially expressed genes in *L. macrocephalus*, total RNA samples were incubated with DNase I to remove DNA contamination which represents a further source of false-positive bands that can be observed on differential display gels. After DNase treatment, small aliquots of each sample were loaded onto an agarose gel to check for the RNA integrity. Subsequently, good quality RNA samples were used to prepare RNA pools of males and females, separately, in order to avoid the detection of interindividual variations not related to differences between sexes.

Further, preliminary analyses were performed to determine primers that give good and reproducible amplification results. As so, cDNA amplification was achieved using decamer oligonucleotides commonly used in RAPD analysis, and also oligonucleotides that were designed based on VNTR (Variable Number of Tandem Repeats) core sequences that correspond to minisatellite short and highly conserved regions (Jeffreys et al. 1985). The use of the 10-mer primers usually resulted on several faint and diffuse bands, and the results were generally not reproducible (data not shown). These faint bands often make interpretation of the data difficult. Moreover, just a reduced number of candidate differentially expressed transcripts could be identified when comparing males and females of L. macrocephalus through the use of these RAPD primers. The use of longer primers on differential display strategies can also lead to detection of DNA polymorphism through RAPD-like results. Some studies evidenced that primers with 13 bases or longer have better effects in the DD efficiency (Zhao et al. 1995, Liang 1998, Motlik et al. 1998, Huang et al. 2001). As the oligonucleotides of VNTR core sequences (14-24 bp), used as single primers to amplify the cDNA samples of L. macrocephalus, are longer than RAPD primers, the methodology could be effectively carried out at a relatively high stringency, thus yielding better results - the generated bands were more distinct and the results of cDNA amplification were consistently repeatable.

The DD methodology generally leads to the visualization of 5 to 100 amplification products on polyacrilamide gel (McClelland *et al.* 1995). The obtained results indicated that, even using longer primers, it was not possible to achieve

complex amplification patterns with a high number of bands, a fact that could be due to the high stringency of the PCR conditions, trough an annealing temperature of 55° C, and due to the use of RNA sample pools that can avoid the detection of individual variations. Despite the relative reduced number of amplified fragments (around 5 to 15 bands) visualized on polyacrilamide gel, some of them were identified only in males or females, which indicate that they correspond to presumptive differentially expressed transcripts between the two samples. Thirty one cDNAs appeared to be differentially expressed between males and females of *L. macrocephalus*.

During the course of the experiments, we experienced a failure to reconfirm differences in mRNA expression, due to difficulties in re-amplifying and/or cloning the selected transcripts, a problem that is relatively common in differential display strategies (Liang *et al.* 1993, Pardee & McClelland 1999). Therefore, just three of the detected candidate fragments (Figure 1) were re-amplified and cloned into a plasmid vector. The primer INS leaded to the identification of a conspicuous fragment of approximately 400 bp only in male samples of *L. macrocephalus* (Figure 1a). A fragment of around 300 bp was also identified only in male samples through the use of the primer HBV5 (Figure 1b). The primer OPP-11 permitted the visualization of a band of approximately 400 bp, unique to females of the species (Figure 1c). Since different cDNA species can be contained theoretically within one differentially expressed fragment (Liang & Pardee 1995), at least five white colonies were selected and analyzed from each cloned fragment.

Real-time PCR was used to validate the differential display results. The indication that some of the isolated genes do not present a sexually dimorphic expression between males and females of *L. macrocephalus*, as previously

deduced through the differential display data, can be associated to several factors that can affect results. Not only the experimental designs and the amplification reaction conditions, but also the systems being compared can alter the obtained data. Comparisons between the same type of cells, as performed in the present study, leads to a higher rate of false-positives simply due to the fewer differences in gene expression (Liang 1998).

DDRT-PCR results using primer INS

Cloning and sequencing of the selected band from males of *L. macrocephalus*, obtained with the primer INS, permitted the characterization of two distinct fragments with 265 bp and 231 bp (Figure 2a, b).

Database searches for nucleotide similarity evidenced that the 265 bp fragment has a high identity level with a partial region of the gene that codes for the protein APBA2 or amyloid beta (A4) precursor protein-binding, family A, member 2, of several vertebrates. A high identity level was evidenced with the *apba2* gene described for the species *Danio rerio* (85%) and *Xenopus tropicalis* (77%), in relation to a 95% query coverage of the analyzed sequence. Gene expression analyses were performed throughout Real-Time PCR using primers denominated Bami (F) and Bami (R) (Table 3), leading to the amplification of a fragment of 126 bp. Further cloning and sequencing of this fragment confirmed that it represents a segment of the *apba2* gene. The qRT-PCR analysis (reaction efficiency above 99%; R2 correlation coefficients \geq 0.97) revealed a significant difference between males and females, as a higher gene expression was detected in males of *L. macrocephalus* (Figure 3a, d; Table 4) thus validating the isolated

fragment as a differentially expressed transcript. The mRNA level of males was almost three-fold higher than in females.

The APBA2 corresponds to a transmembrane cell surface glycoprotein that is mainly expressed in brain (Sutcliffe et al. 2003, Taru & Suzuki 2004, Sano et al. 2006) and that can modulate the A β PP (amyloid β -protein precursor) expression, leading to its decrease or increase. Consecutive cleavages of the ABPP by secretases generate beta-amyloid proteins (βA) (Taru & Suzuki 2004, Sano et al. 2006) that, at normal physiological levels, appear to be involved in neurotrophic events, i.e. the growth and survival of developing neurons and the maintenance of mature neurons (Deister & Schimidt 2006). However, an overexpression of the $a\beta pp$ gene can lead to an increased production of the beta-amyloid protein (e.g. Mattson et al. 1993), a feature that has been related to Alzheimer's disease, a frequent neurodegenerative disorder of the central nervous system that causes mental deterioration and progressive dementia due to neuropathologic lesions including the presence of senile plaques mainly composed by the beta-amyloid protein (e.g. Selkoe 1994). Furthermore, it also seems that mutations along the aßpp gene or mutations in the enzymes that cleave the ABPP into beta-amyloid proteins can also lead to the development of Alzheimer's disease (Harrison et al. 1991, Hendriks & VanBroeckhoven 1996, Hartmann et al. 1996, Panegyres 1997, Cole et al. 2001, Yamamoto & Behl 2001, Xie et al. 2003).

Data on Alzheimer's disease indicate a prevalence of this neurodegenerative disorder in females of some human populations, a result that seem to be correlated not only to hormones, as estrogen and gonadotropins (Casadesus *et al.* 2008), but also to age and education level (Letenneur *et al.* 2000). The identification of a higher *apba2* gene expression in males of *L*.

macrocephalus can be correlated to an increase in amyloid β-protein precursor which, in turns, would lead to a higher content of beta-amyloid protein in the brain of animals of this sex. Even though the functional significance of the increased *apba2* expression in males than in females of *L. macrocephalus* is unclear, this fish species, similarly to other vertebrates as mouse (Fisher *et al.* 1991), *Xenopus laevis* (Van den Hurk *et al.* 2001) and chicken (Carrodeguas *et al.* 2005), can also be used as a model in order to studying Alzheimer's disease development or even disease therapeutic agents.

The second characterized 231 bp fragment of *L. macrocephalus*, isolated through the use of the primer INS, evidenced a high similarity with a partial region of the *Rab* 37 gene, a member of the *Ras* oncogene family. This isolated transcript of *L. macrocephalus* has 87% and 85% identity, at nucleotide level, to the *Rab* 37 gene described for *Danio rerio* and with the variant transcript 3 of this gene of *Homo sapiens*, respectively. The quantitative gene expression analysis (reaction efficiency above 99%; R2 correlation coefficients \geq 0.97) was performed using designed primers denominated Rab37 (F) and Rab37 (R) (Table 3), leading to the amplification of a fragment of 170 bp. Further cloning and sequencing of this fragment confirmed that it represents a segment of the *Rab37* gene. Although the statistical analyses revealed a higher expression level in males than in females of *L. macrocephalus*, the differences were not statistically significant (Figure 3b, d; Table 4).

The *Rab37* gene is a member of the most common oncogenes related to human tumors - the *Ras* oncogene family. The Ras proteins correspond to small GTP-binding proteins that are involved in signal transduction pathway of several normal cellular functions, as cell proliferation, differentiation, adhesion, apoptosis,

82

and migration (Masuda *et al.* 2000, Stenmark & Olkkonen 2001). In brain, the *Rab37* gene is mainly related to the differentiation of neuronal cells (Bar-Sagi & Feramisco 1985, Noda *et al.* 1985, Hagag *et al.* 1986). Cancer induction seems to be related to overexpression of *Ras* genes or alterations in their proteins, which lead to an abnormal cellular proliferation (Capon *et al.* 1983, Brown *et al.* 1984, Feramisco *et al.* 1984, Stacey & Kung 1984).

Higher expression levels of *Ras* oncogenes have been described in human brain-tumors (Zimmer *et al.* 1987, Lee *et al.* 2008). However, there is no conclusive data on a sexually differential expression of these genes, since some genes of the *Ras* family present a higher expression in women while others are highly expressed in men, a feature that can also be related to the type of the cancer and age (Stenmark & Olkkonen 2001). Therefore, the obtained results that suggest a higher expression of the *Rab37* gene in males of *L. macrocepalus* remain to be elucidated. Comprehending the ways in which the regulatory actions of Rab proteins intertwine with cell-signaling cascades and developmental processes remains to be clarified.

DDRT-PCR results using primer HBV5

After cloning and sequencing the cDNA fragment that was amplified through the use of primer HBV5, observed in male samples of *L. macrocephalus*, it was possible to characterize a 194 bp segment (Figure 2c). Data base searches revealed no significant similarity with any known gene, despite a ~30 bp region that showed an identity level of 93% with a region of the gene encoding protein C (PROC) of *Gallus gallus* that functions as an anti-inflammatory and also as an inhibitor of the coagulation factors Va and VIIIa, subsequently blocking the generation of thrombin (Jijo *et al.* 2007). Although the isolated transcript did not reveal a significant similarity to the *proc* gene (database searches evidenced a 15% query coverage of the analyzed sequence), it is also possible that the characterized segment represents an untranslated region of the mRNA which is commonly less conserved than the coding regions between homologous genes in different organisms (Colona-Romano *et al.* 1998).

A new set of primers - HBV5 (F) and HBV5 (R) - was designed in order to be used on qRT-PCR (reaction efficiency above 99%; R2 correlation coefficients \geq 0.96), leading to the amplification of a fragment of 108 bp. Cloning and sequencing of this fragment confirmed a high identity with the previous isolated fragment trough the differential display strategy. Although most males presented a higher expression level of this transcript, the observed difference between both sexes was not statistically significant (Figure 3c, d; Table 4).

DDRT-PCR results using primer OPP-11

The cDNA amplification using primer OPP-11 generated a female-specific band of approximately 400 bp (Figure 1c). Cloning and sequencing of this fragment permitted the characterization of a 414 bp segment that revealed no significant identity level with previously reported gene sequences. An 89% identity level could be observed between a 36 bp of the isolated fragment of *L. macrocephalus* and the BAC clone CH251-315P9 of *Pan troglodytes* and the BAC clone RP11-54F22 of *Homo sapiens* that were mapped to the X chromosome of these species. Despite no information on the sequences contained in these BACs, the database searches evidenced extremely reduced query coverage (9%) of the analyzed sequence.

84

The characterized nucleotide sequence of the isolated transcript was used to design a set of primers – primers 11P(F) and 11P (R) (Table 3)– to be used on the real time PCR assay. However, there was no success in obtaining a single amplification product. Primer design and PCR conditions could have lead to the observed results, a feature that seem to be common in quantitative RT-PCR analyses (Roling *et al.* 2004). Although the obtained data do not provide relevant information about the nature of the isolated fragment, its putative dimorphic expression in males and females of *L. macrocephalus* indicates that this transcript should also be better investigated.

Final considerations

Sexually dimorphic genes have been described for different vertebrates in somatic tissues, a feature that has been proven to be due to sexual hormones, specially estrogens and androgens and its metabolites (Dewing *et al.* 2003, Morris *et al.* 2004, McCarthy & Konkle 2005, Becker *et al.* 2005, Tomassini & Pozzilli 2006, Verma & Shapiro 2006). Recent evidences indicate that differences in the expression of some genes in brain also arise under the influence of the sex chromosome complement, independent of gonadal sex steroids levels (Arnold 2004). However, there is still several lacks in the comprehension of the regulation of these genes in brain, especially in lower vertebrates. Improving data on the isolation and characterization of these transcripts in fish species, that comprise the basal vertebrate group, could give new insights into genes structure, function and evolution.

To get better insights into several genes that can present a different expression level depending on the sex, the availability of a proper model system may be crucial. The zebrafish *Danio rerio* is actually one of the few lower

85

vertebrate models in which genome-wide methods can be employed effectively. However, the fundamental process controlling sex assignment in this species has yet to be clarified, since no heteromorphic sex chromosomes have been detected (Sola & Gornung 2001). *Leporinus macrocephalus* is a gonochoristic species that presents a ZZ/ZW sex determination system (Galetti & Foresti 1987). Moreover, it has been intensively exploited in the subtropical aquaculture due to characteristics as the facility to adaptation to captive breeding conditions, good reproductive performance and excellent growth rate (Takahashi *et al.* 2004). Therefore, it may represent an interesting fish model to the understanding of the function of several vertebrate genes associated to sex determination and differentiation. In addition, as evidenced by the present results on the *apba2* and *Rab37* genes, it could also be used as a natural model for the assay of drugs that can regulate several processes related to genes involved in neurodegenerative and cancer diseases.

Acknowledgements

The authors thank Dr. Maeli Dal Pai Silva, Dr. João Pessoa, and Dr. Débora Colombi for technical assistance on the Real Time analyses. This work was supported by grants from FAPESP (Fundação da Amparo à Pesquisa do Estado de São Paulo) and CNPq (Conselho Nacional de Desenvolvimento Científico e Tecnológico).

References

- Almeida-Toledo LF & Foresti F (2001) Morphologically differentiated sex chromosomes in neotropical freshwater fish. *Genetica* **111**: 91-100.
- Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ (1990) Basic local alignment search tool. *Journal of Molecular Biology* **215**: 403-410.
- Arnold AP (2004) Sex, chromosomes and brain gender. *Natural Review in Neuroscience* **5**: 701-708.
- Bar-Sagi D & Feramisco JR (1985) Microinjection of the ras oncogene protein into PC12 cells induces morphological differentiation. *Cell* **42**: 841-848.
- Becker JB, Arnold AP, Berkley KJ, Blaustein JD, Eckel LA, Hampson E, Herman JP, Marts S, Sadee W, Steiner M, Taylor J, Young E (2005) Strategies and methods for research on sex differences in brain and behavior. *Endocrinology* **146**: 1650-1673.
- Brown R, Marshall CJ, Pennie SG, Hall A (1984) Mechanism of activation of an N*ras* gene in the human fibrosarcoma cell line. *EMBO Journal* **3**: 1321-1326.
- Capon DJ, Seeburg PH, McGrath JP, Hatflick JS, Edman U, Levinson AD, Goeddel DV (1983). Activation of K1-ras2 gene in the human colon and lung carcinomas by two different point mutations. *Nature* **304**: 507-513.
- Carrodeguas JA, Rodolosse A, Garza MV, Sanz-Clemente A, Pérez-Pé R, Lacosta AM, Dominguez L, Monleón I, Sánchez-Díaz R, Sorribas V, Sarasa M (2005) The chick embryo appears as a natural model for research in beta-amyloid precursor protein processing. *Neuroscience* **134**: 1285-1300.
- Casadesus G, Rolston RK, Webber KM, Atwood CS, Bowen RL, Perry G, Smith MA (2008) Menopause, estrogen, and gonadotropins in Alzheimer's disease. *Advances in Clinical Chemistry* **45**: 139-153.
- Cole GM, Masliah E, Shelton ER (1991) Accumulation of amyloid precursor fragment in Alzheimer plaques. *Neurobiology of Aging* **12**: 85-91.
- Colona-Romano S, Porta A, Franco A, Kobayashi GS, Maresca B (1998) Identification and isolation by DDRT-PCR of genes differentially expressed by

Histoplasma capsulatum during macrophages infection. *Microbial Pathogenesis* **25**: 55-66.

- Conover DO, Van Voorhes DA, Ehtisham A (1992) Sex-ratio selection and the evolution of environmental sex determination in laboratory populations of *Menidia menidia. Evolution* **46**: 1722-1730.
- Dakis D & Kouretas D (2002) Differential display of m-RNAs in fish gonads by modified DD-PCR. *In vivo* **16**: 103-105.
- Davis LK, Pierce AL, Hiramatsu N, Sullivan CV, Hirano T, Grau EG (2008) Genderspecific expression of multiple estrogen receptors, growth hormone receptors, insulin-like growth factors and vitelogenins, and effects of 17 beta-estradiol in the male tilapia (Oreochromis mossambicus). General and Comparative Endocrinology **156**: 544-551.
- D'Cotta H, Fostier A, Guiguen Y, Govoroun M, Baroiller JF (2001) Search for genes involved in the temperature-induced gonadal sex differentiation in the tilapia, *Oreochromis niloticus*. *Journal of Experimental Zoology* **290**: 574-585.
- Deister C & Schmidt CE (2006) Optimizing neurotrophic factor combinations for neurite outgrowth. *Journal of Neural Engineering*. **3**: 172-179.
- Denslow ND, Bowman CJ, Ferguson RJ, Lee HS, Hemmer MJ, Folmar LC (2001a) Induction of gene expression in sheepshead minnows (*Cyprinodon variegatus*) treated with 17 beta-estradiol, diethylstilbestrol, or ethinylestradiol: The use of mRNA fingerprints as an indicator of gene regulation. *General and Comparative Endocrinology* **121**: 250-260.
- Denslow ND, Lee HS, Bowman CJ, Hemmer MJ, Folmar LC (2001b) Multiple responses in gene expression in fish treated with estrogen. *Comparative Biochemistry and Physiology B-Biochemistry and Molecular Biology* **129**: 277-282.
- Dewing P, Shi T, Horvath S (2003) Sexually dimorphic gene expression in mouse brain precedes gonadal differentiation. *Molecular Brain Research* **118**: 82-90.
- Eriksson A, Wahlestedt C, Nordqvist K (1999) Isolation of sex-specific cDNAs from fetal mouse brain using mRNA differential display and representational difference analysis. *Molecular Brain Research* **74**: 91-97.

- Feramisco JR, Gross M, Kamata T, Rosenberg M, Sweet RW (1984) Microinjection of the oncogene formo f the human H-ras (T-24) protein results in rapid proliferation of guiescent cells. *Cell* **38**: 109-117.
- Filby AL & Tyler CR (2007) Cloning and characterization of cDNAs for hormones and/or receptors of growth hormone, insulin-like growth factor-I, thyroid hormone, and corticosteroid and the gender-, tissue-, and developmental-specific expression of their mRNA transcripts in fathead minnow (*Pimephales promelas*). *Genetics and Comparative Endocrinology* **150**: 151-163.
- Fisher S, Gearhart JD, Oster-Granite ML (1991) Expression of the amyloid precursor protein gene in mouse oocytes and embryos. *Proceedings of the Natural Academy of Science USA* **88**: 1779-1782.
- Galetti PM & Foresti F (1987) Two new cases of ZZ/ZW heterogamety in *Leporinus* (Anostomidae, Characiformes) and their relationships in the phylogeny of the group. *Revista Brasileira de Genética* **10**: 135-140.
- Goto-Kazeto R, Kight KE, Zohar Y, Place AR, Trant JM (2004) Localization and expression of aromatase mRNA in adult zebrafish. *Genetics and Comparative Endocrinology* **139**: 72-84.
- Graf D, Fisher AG, Merkenschlager M (1997) Rational primer design greatly improves differential display PCR (DD–PCR). *Nucleic Acids Research* **25**: 2239-2240.
- Grossmann R, Jurkevich A, Klein S (2004) Gender-specific expression of argininevasotocin in the chicken brain. *Avian and Poultry Biology Reviews* **15**: 219-223.
- Guan G, Kobayashi T, Nagahama Y (2000) Sexually dimorphic expression of two types of DM (Doublesex/Mab-3)-domain genes in a teleost fish, the tilapia (*Oreochromis niloticus*). *Biochemistry and Biophysics Research Community* 272: 662-666.
- Hagag N, Halegoua S, Viola M (1986) Inhibition of growth factor-induced differentiation of PC12 cells by microinjection of antibody to ras p21. *Nature* **319**: 680-682.
- Harris AS & Wrigth JM (1995) Nucleotide sequence and genomic organization of cichlid fish minisatellites. *Genome* **38**: 177-184.

- Harrison PJ, Barton AJL, Pearson RCA (1991) Expression of amyloid beta-protein precursor messenger-RNAs in familial Alzheimer disease. *Neuroreport* **2**: 152-154.
- Hartmann T, Bergsdorf C, Sandbrink (1996) Alzheimer's disease beta A4 protein release and amyloid precursor protein sorting are regulated by alternative splicing. *Journal of Biological Chemistry* **271**: 13208-13214.
- Hendriks L & VanBroeckhoven C (1996) The beta A4 amyloid precursor protein gene and Alzheimer's disease. *European Journal of Biochemistry* **237**: 6-15.
- Huang YJ, Li Y, Shi M, Wang JA, Hu MR, Shen BF (2001) Isolation of genes related with IL-6 effect by a rapid and simple RT-PCR method. *Progress in Biochemistry and Biophysics* **28**: 563-567.
- Hutchison JB (1997) Gender-specific steroid metabolism in neural differentiation. *Cellular and Molecular Neurobiology* **17**: 603-626.
- Isensee J, Witt H, Pregla R, Hetzer R, Regitz-Zagrosek V, Noppinger PR (2008) Sexually dimorphic gene expression in the heart of mice and men. *Journal of Molecular Medicine* **86**: 61-74.
- Jadhao AG, D'Aniello B, Malz CR, Pinelli C, Meyer DL (2001) Intrasexual and intersexual dimorphisms of the red salmon prosencephalon. *Cell Tissue Research* **304**:121-140.
- Jeffreys, AJ, Wilson V, Thein SL (1985) Hypervariable minisatellite regions in human DNA. *Nature* **314**: 67-73.
- Jijo J, Awab A, Norman D, Dernaika T, Kinasewitz GT (2007) Activated protein C improves survival sepsis patients with elevated troponin. *Intensive Care Medicine* 33: 2122-2128.
- Jurecic R (1996) Differential mRNA display using anchored oligo-dT and long sequence-specific primers as arbitrary primers. *Trends in Genetics* **12**: 502-504.
- Kallivretaki E, Eggen RIL, Neuhauss SCF (2007) The zebrafish brain-specific aromatase cyo19a2 is neither expressed nor distributed in a sexually dimorphic manner during sexual differentiation. *Developmental Dynamics* **236**: 3155-3166.

- Kanamori A (2000) Systematic identification of genes expressed during early oogenesis in Medaka. *Molecular Reproduction and Development* **55**: 31-36.
- Kondo M, Nanda I, Hornung U, Asakawa S, Shimizu N, Mitani H, Schmid M, Shima A, Schartl M (2003) Absence of the candidate male sex-determining gene dmrt1b(Y) of medaka from other fish species. *Current Biology* **13**: 416-420.
- Lauer LE, McCarthy MM, Mong J, Kane AS (2006) Sex differences in neural morphology in the killifish hypothalamus. *Brain Research* **1070**:145-149.
- Lee JS, Gil JE, Kim JH (2008) Brain cancer stem-like genesis p53-deficient mouse astrocytes by oncogenic Ras. *Biochemical and Biophysical Research Communications* **365**: 496-502.
- Letenneur L, Launer LJ, Andersen K (2000) Education and the risk for Alzheimer's disease: sex makes a difference. EURODEM pooled analyses. *American Journal of Epidemiology* **151**: 1064-1071.
- Liang P (1998) Current progress in differential display methods and applications.
 In: *Methods. A Companion to Methods in Enzymology*. Vol 16. Academic Press, New York.
- Liang P (2002) A decade of differential display. *BioTechniques* **33**: 338-346.
- Liang P, Averboukh L, Pardee AB (1993) Distribution and cloning of eukaryotic mRNAs by means of differential display: refiniments and optimization. *Nucleic Acids Research* **21**: 3269-3275.
- Liang P & Pardee AB (1995) Recent advances in differential display. *Current Opinion in Immunology* **7**: 274-280.
- Liu O, Goudie CA, Simco BA, Davis KB (1996) Sex-linkage of glucose phosphate isomerase-B and mapping of the sex-determining genes in channel catfish. *Cytogenetics and Cell Genetics* **73**: 282-285.
- Livak KJ & Schmittgen TD (2001) Analysis of relative expression data using realtime quantitative PCR and the $2^{-\Delta\Delta C}$ Method. *Methods* **25**: 402-408.
- Lutfalla G, Crollius HR, Bruner FG, Laudet V, Robinson-Rechavi M (2003) Inventing a sex-specific gene: A conserved role of DMRT1 in teleost fishes plus

a recent duplication in the medaka *Oryzias latipes* resulted in DMY. *Journal of Molecular Evolution* **57**: 148-153.

- MacClelland M, Mathieu-Daude, F, welsh J (1995) RNA fingerprinting and differential display using arbitrary primed PCR. *Trends in Genetics* **11**: 242-246.
- Malhotra L, Foltz L, Mahoney WC, Schueler PA (1998) Interaction and effect of annealing temperature on primers used in differential display RT–PCR. *Nucleic Acids Research* **16**: 854-856.
- Marin I & Baker BS (1999) The evolution dynamics of sex determination. *Science* **281**: 1990-1994.
- Masuda ES, Luo Y, Young C, Sen M, Rossi AB, Huang BC, Yu S, Bennett MK, Payan DG, Scheller RH (2000) Rab37 is a novel mast cell specific GTPase localized to secretory granules. *Federation of European Biochemical Societies Letters* 470: 61-64.
- Matsuda M, Nagahama Y, Shinomiya A, Sato T, Matsuda C, Kobayashi T, Morrey CE, Shibata N, Asakawa S, Shimizu N, Hori H, Hamaguchi S, Sakaizumi M (2002) A Y-specific, DM-domain gene, *DMY*, is required for male development in the medaka (*Oryzias latipes*). *Nature* **417**: 559-563.
- Mattson MP, Cheng B, Culwell AR, Esch FS, Lieberburg I, Rydel RE (1993) Evidence for excitoprotective and intraneuronal calcium-regulating roles for secreted forms of the ß-amyloid precursor protein. *Neuron* **10**: 243-254.
- Mayer A, Lahr G, Swaab DF, Pilgrim C, Reisert I (1998) The Y-chromosomal genes SRY and ZFY are transcribed in adult human brain. *Neurogenetics* **1**: 281-288.
- McCarthy MM & Konkle AT (2005) When is a sex difference not a sex difference? *Frontiers in Neuroendocrinology* **26**: 85-102.
- Moles G, Carrillo M, Mananos E, Mylonas CC, Zanuy S (2007) Temporal profile of brain and pituitary GnHs, GnRH-R and gonadotropin mRNA expression and content during early development in European sea bass (*Dicentrarchus labrax* L.). *Genetics and Comparative Endocrinology* **150**: 75-86.

- Moreira-Filho O, Bertollo LAC, Galetti PM (1993) Distribution of sex-chromosome mechanisms in Neotropical fish and description of a ZZ/ZW system in *Parodon hilarii* (Parodontidae). *Caryologia* **46**: 115-125.
- Morris JA, Jordan CL, Breedlove SM (2004) Sexual differentiation of the vertebrate nervous system. *Natural Neurosciences* **7**: 1034-1039.
- Motlik J, Carnwath JW, Herrmann D, Terletski V, Anger M, Niemann H (1998) Automated recording of RNA differential display patterns from pig granulose cells. *Biotechniques* **24**: 148-153.
- Murray MJ, Haldeman BA, Grant FJ, O'Hara N (1988) Probing the human genome with minisatellite-like sequences from the human coagulation factor-VII gene. *Nucleic Acids Research* **16**: 4166.
- Nakamura Y, Leppert M, O'Connell P, Wolff R, Holm T, Culver M, Martin C, Fujimoto E, Hoff M, Kumlin E, White R (1987) Variable number of tandem repeat (VNTR) markers for human-gene mapping. *Science* **235**: 1616-1622.
- Noda M, Ko M, Ogura A, Liu DG, Takano T, Ikawa Y (1985) Sarcoma viruses carryinf ras oncogenes induce differentiatio-associated properties in a neuronal cell line. *Nature* **318**: 73-75.
- Panegyres PK (1997) The amyloid precursor protein gene: a neuropeptide gene with diverse functions in the central nervous system. *Neuropeptides* **31**: 523-535.
- Pardee AB & McClelland M (1999) Expression genetics: differential display. In: *Expression genetics: differential display*. Pardee AB & Mc Clelland M (eds.). Eaton Publishing, USA, pp. 1-7.
- Rinn JI & Snyder M (2005) Sexual dimorphism in mammalian gene expression. *Trends in Genetics* **21**: 298-305.
- Roling JA, Bain LJ, Baldwin WS (2004) Differential gene expression in mummichogs (*Fundulus heteroclitus*) followinh treatment with pyrene: comparison to a creosote contaminated site. *Marine Environmental Research* 57: 377-395.
- Sambrook J & Russel DW (2001) Molecular cloning. A Laboratory Manual, 3rd Edition. Cold Spring Harbor Laboratory Press, New York.

- Sano Y, Syuzo-Takabatake A, Nakaya T, Saeto Y, Tomita S, Itohara S, Suzuki T (2006) Enhanced amyloidogenic metabolism of the amyloid β-protein precursor in the X11L-deficient mouse brain. *The Journal of Biological Chemistry* 281:3785-37860.
- Santos EM, Kille P, Workman VL, Paull GC, Tyler CR (2008) Sexually dimorphic gene expression in the brains of mature zebrafish. *Comparative Biochemistry and Physiology* Part A **149**: 314-324.
- Selkoe DJ (1994) Cell biology of the amyloid ß-protein precursor and the mechanism of Alzheimer's disease. *Annual Review in Cell Biology* **10**: 373-403.
- Stacey DW & Kung HF (1984) Transformation of NIH 3T3 cells by microinjection of Ha-ras p21 protein. *Nature* **310**: 508-511.
- Stenmark H & Olkkonen VM (2001) The Rab GTPase family. *Genome Biology* **2**: 1-7.
- Sola L & Gornung E (2001) Classical and molecular cytogenetics of the zebrafish *Danio rerio* (Cyprinidae, Cypriniformes): an overview. *Genetica* **111**: 397-412.
- Sutcliffe JS, Han MK, Amin T, Kesterson RA, Nurmi EL (2003) Partial duplication of the APBA2 gene in chromosome 15q13 corresponds to duplicon structures. *BMC Genomics* **4**:15.
- Takahashi LS, Gonçalves FD, de Abreu JS, Martins MIEG, Ferreira ACM (2004) Economic viability of the piauçu *Leporinus macrocephalus* (Garavello & Britski, 1988) production. *Scientia Agricola* **61**: 228-233.
- Taru H & Suzuki T (2004) Facilitation of stress-induced phosphorylation of βamyloid precursor protein family members by X11-like/ Mint2 protein. *The Journal of Biological Chemistry* **279**: 21628-21636.
- Tave D (1993) Genetics of quantitative phenotypes. Genetics for Fish Hatchery Managers, 2nd edn. An AVI book, New York, pp. 117–264
- Thompson JD, Higgins DG, Gibson TJ (1994) Clustal W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Research* **22**: 4673-4680.

- Tomassini V & Pozzilli C (2006) Sex hormones: a role in the control of multiple sclerosis? *Expert Opinion on Pharmacotherapy* **7**: 857-868.
- Toriyama E (2001) Reversão de sexo e análise comparativa do crescimento em piuçú (*Leporinus macrocephalus*) Anostomidae, Teleostei. Master Thesis.
 Universidade Estadual Paulista Júlio de Mesquita Filho, Botucatu, SP.
- Van den Hurk W, Bloemen M, Martens GJM (2001) Expression of the gene encoding the β-amyloid precursor protein APP in *Xenopus laevis*. *Molecular Brain Research* **97**: 13-20.
- Verma AS & Shapiro BH (2006) Sex-dependent expression of seven housekeeping genes in rat liver. *Journal of Gastroenterology and Hepatology* 21: 1004-1008.
- Zhao A, Ooi SL, Pardee AB (1995) New primer strategy improves precision of differential display. *BioTechniques* **18**: 842-850.
- Wilhelm D, Palmer S, Koopman P (2007) Sex determination and gonadal development in mammals. *Physiology Review* **87**: 1-28.
- Xie J, Wen JJ, Chen B, Gui JF (2001) Differential gene expression in fully-grown oocytes between gynogenetic and gonochoristic crucian carps. *Gene* **271**: 109-116.
- Xie YY, Yao ZB, Chai H (2003) Potential roles of Alzheimer precursor protein A4 and beta-amyloid in survival and function of aged spinal motor neurons after axonal injury. *Journal of Neuroscience Research* **73**: 557-564.
- Xu J, Burgoyne P, Arnold AP (2001) Expression of sex chromosome genes in mouse brain. *American Journal of Human Genetics* **69**: 341-341.
- Xu J & Disteche CM (2006) Sex differences in brain expression of X- and Y-linked genes. *Brain Research* **1**: 50-55.
- Yamamoto T (1969) Sex differentiation. In: *Fish Physiology*. Hoar W & Randal DP (eds.). Academic Press, New York, pp.117-175.
- Yamamoto A & Behl C (2001) Human Nck-associated protein1 and its binding protein affect the metabolism of beta-amyloid precursor protein with Swedish mutation. *Neuroscience Leters* **316**: 50-54.

- Yang X, Schadt EE, Wang S, Wang H, Arnold AP, Ingram-Drake L, Drake TA, Lusis AJ (2006) Tissue-specific expression and regulation of sexually dimorphic genes in mice. *Genome Research* **16**: 995-1004.
- Yuge K, Ikeo K, Gojobori T (2007) Evolutionary origin of sex-related genes in the mouse brain. *Gene* **406**: 108-112.
- Zar JH (1999) Biostatistical analysis. Prentice-Hall, New Jersey, pp.663.
- Zarkower D (2001) Establishing sexual dimorphism: conservation amidst diversity? *Nature Review Genetics* **2**: 175-185.
- Zimmer SG, Fetherston J, Cotton J (1987) Role of activated Ras oncogene in progression of hamster brain-tumors. *Journal of Cellular Biochemistry* **11**: 103.

Tables, Figures and Legends

Primer	Sequence (5' - 3')	Reference
INS	ACAGGGGTGTGGGG	Nakamura <i>et al.</i> (1987)
YNZ22	CTCTGGGTGTCGTGC	Nakamura <i>et al.</i> (1987)
HBV5	GGTGTAGAGAGGGGT	Nakamura <i>et al.</i> (1987)
HBV3	GGTGAAGCACAGGTG	Nakamura <i>et al.</i> (1987)
FvIIex8	ATGCACACACAGG	Murray <i>et al.</i> (1988)
EMBL	AGAGCTTCAGGCTGGGCAGCTAAG	Harris & Wright (1995)

Table 1: Oligonucleotides that were designed based on VNTR core sequences and used as single primers for the amplification of the cDNA samples of *Leporinus macrocephalus*.

Primer	Sequence (5' - 3')
OPP-1	GTAGCACTCC
OPP-2	TCGGCACGCA
OPP-3	CTGATACGCC
OPP-4	GTGTCTCAGG
OPP-5	CCCCGGTAAC
OPP-6	GTGGGCTGAC
OPP-7	GTCCATGCCA
OPP-8	ACATCGCCCA
OPP-9	GTGGTCCGCA
OPP-10	TCCCGCCTAC
OPP-11	AACGCGTCGG
OPP-12	AAGGGCGAGT
OPP-13	GGAGTGCCTC
OPP-14	CCAGCCGAAC
OPP-15	GGAAGCCAAC
OPP-16	CCAAGCTGCC
OPP-17	TGACCCGCCT
OPP-18	GGCTTGGCCT
OPP-19	GGGAAGGACA

Table 2: Decamer oligonucleotides used as singleprimers for the amplification of the cDNA samples ofLeporinus macrocephalus.

Primer	(5' - 3')	
Bami (F) Bami (R)	CATTCTTACGCTCTGCCATG TAGTAGCCCGTGATAAGGAC	-
Rab37 (F) Rab37 (R)	GTGGGAGTTCAAATGCTGGG GGAAATTCATCCCTGGCTCC	
HBV5 (F) HBV5 (R)	CCCTGTTACTGCGCATGAG CACGCGGTATCAGGATCA	
P11 (F) P11 (R)	GCTACATTTCAGAGTATAATAG CATATGAAAGCTTAGAATCTCT	

Table 3: Set of primers that were designed based on the nucleotidesequences of the isolated DNA fragments from Leporinusmacrocephalus, in order to be used on Real-Time PCR assays.

F = forward; R = reverse
Designed set of primers	Mean* ± SD	"t" value	"P" values
Bami	M = 2,53x10 ⁻⁵ ±3,47x10 ⁻⁶ F = 8,16x10 ⁻⁶ ±1,53x10 ⁻⁶	-14,27	<0,001
Rab37	M = 8,44x10 ⁻⁷ ±4,89x10 ⁻⁷ F = 7,03x10 ⁻⁷ ±4,33x10 ⁻⁷	98**	0,623**
HBV5	M = 1,92×10 ⁻⁷ ±8,52×10 ⁻⁸ F = 1,4×10 ⁻⁷ ±5,85×10 ⁻⁸	-1,568	0,134

Table 4: Data on the relative expression of the isolated genes from brain tissue of Leporinus macrocephalus.

*Mean value refering to the values obtained for 2^{-ΔCt} ** Refers to "T" value obtained for these samples through the Mann-Whitney statistical test M, male samples; F, female samples



Figure 1: cDNA amplification patterns of *Leporinus macrocephalus*, visualized in 6% polyacrilamide gel, obtained through the use of primers INS (a), HBV5 (b), and OPP-11 (c). Sexassociated fragments are indicated by arrows. L, 1Kb DNA ladder; M, males; F, females.

GCTCCTGCCCCCCCCCCCCCCCCCCCCCCCCCCCCCCC	AGAATTCT ACGGGTAC
GCGTGTCACAGCGGAAAACGTGACATTTTAATATCCGAGTGTCCTTATCACG205	TGGTTCTC
GGCTACATACGCAAAATCCCTCTCTCTGCCGTTGTCCCGGCCTACG <u>CCCCAC</u> 257 ACCCCTGT 3'	(c)
(a)	
	5'AACGCO
	AGATGTNO
	CTCTACT
5' <u>ACAGGGGTGTGGGG</u> GGGTGGGAGTTCAAATGCTGGGTACTATAGTGAAA49	AATATTT
GATCAATCATGGAAAGGCAAAAAAAGATGAACTGTGTATTCTGCTCTACGCA101	GAACATT
ACGCTGCAAATAACATATTAAATGCTTGTTTTATTCGTGAAGCCGATGCCCA153	ACATCAAA
CAGTAGCAGAAAAGGAGCCAGGGATGAATTTCCCCTGATCAAACTGGACGAG205	CTCTGAAA
CAGGGACGTTTT <u>CCCCACACCCCTGT</u> 3'231	AGTAACA
(b)	(d)

5'GGTGTAGAGAGGGGTGCTGGAGCTGTGATACCACACGCGGTATCAGGATC50
AGAATTCTGGTATCACGACAGACGCACAGCTGCAGTAGCGACCCTGCTGATA102
ACGGGTACTCGGTTACAGCACTCATGCGCAGTAACAGGGTCTCCGGGCTTTC154
TGGTTCTCCTGCTGACCCGATGATG <u>ACCCCTCTCTACACC</u> 3'194
(c)
5' <u>AACGCGTCGG</u> GCCAGGAATCAAAAATAAATATATTTTAAGGATATATCTGA50
AGATGTNCAGTACCACCAAAGACAATCAAAGTACCATATGAAAGCTTAGAAT102
CTCTACTTTTCGGTCGTTTTGCCATTTTTTTTTTTTTTT
AATATTTTGTTCAAAAACTAGCGTTAGTATGACAAAATATTAGGTACACTAC206
GAACATTTTTTTTTTTTCGTTAAGCCATCTTTTGGCCAATCTGGACATATTAC258
ACATCAAACGAATGGGGGAGACCCTCAGGATTACGATGGTGTAAGCTATTATA310
CTCTGAAATGTAGCAGTATTTAAATAAAAAAAAAAAGAATGGCTTTTCCAAAC362
AGTAACACGCCTCCTAGCGCCGCCACTACTTCTGCGTTTCT <u>CCGACGCGTT</u> 3'.414
4-1

Figure 2: Consensus nucleotide sequences of cDNA fragments of *Leporinus macrocephalus*, corresponding to candidate differentially expressed genes between males and females, isolated from brain tissue using primers INS (**a** and **b**), HBV5 (**c**), and OPP-11 (**d**). The underlined regions are homologous to the primer sequences.



Figure 3: Relative gene expression analysis (Real-Time PCR) of the candidate sexually dimorphic expressed genes isolated through DDRT-PCR from brain tissue of *Leporinus macrocephalus*, relative to the constitutive 18S rRNA gene used as control. (a), (b), and (c) represent data on males and females, separately, using set of primers (denominated as Bami, Rab37, and HBV5) designed based on the previous nucleotide sequences of the isolated transcripts; (d) represents a comparative analysis of the relative expression of the three isolated genes – the females were assigned by a reference value = 1, and the males were represented by a ratio of their values in relation to the reference female value. Data were run in duplicate. Normalized data were expressed as means \pm SD.

CAPÍTULO 3*

Distribuição espaço temporal de transcritos do gene de *fatty acid-binding protein 6 (fabp6)* em *zebrafish (Danio rerio)* adulto e em desenvolvimento

* Artigo publicado: FEBS Journal 275: 3325-3334 (2008).

Spatio-temporal distribution of fatty acid-binding protein 6 (*fabp6*) gene transcripts in the developing and adult zebrafish (*Danio rerio*)

Abstract

We have determined the structure of the fatty acid-binding protein 6 (fabp6) gene and the tissue-specific distribution of its transcripts in embryos, larvae and adult zebrafish (Danio rerio). Like most members of the vertebrate FABP multigene family, the zebrafish fabp6 gene contains four exons separated by three introns. The coding region of the gene and expressed sequence tags code for a polypeptide of 131 amino acids (14 kDa, pl of 6.59). The putative zebrafish Fabp6 protein shared greatest sequence identity with human FABP6 (55.3%) compared to other orthologous mammalian FABPs and paralogous zebrafish Fabps. Phylogenetic analysis showed that the zebrafish Fabp6 formed a distinct clade with the mammalian FABP6s. The zebrafish fabp6 gene was assigned to linkage group (chromosome) 21 by radiation hybrid mapping. Conserved gene synteny was evident between the zebrafish fabp6 gene on chromosome 21 and the FABP6/Fabp6 genes on human chromosome 5, rat chromosome 10 and mouse chromosome 11. Zebrafish fabp6 transcripts were first detected in the distal region of the intestine of embryos at 72 h postfertilization. This spatial distribution remained constant to 7-day-old larvae, the last stage assayed during larval development. In adult zebrafish, fabp6 transcripts were detected by RT-PCR in RNA extracted from liver, heart, intestine, ovary and kidney (most likely adrenal tissue), but not in RNA from skin, brain, gill, eye or muscle. In situ hybridization of a *fabp6* riboprobe to adult zebrafish sections revealed intense hybridization signals in the adrenal homolog of the kidney and distal region of the intestine, and to a lesser extent in the ovary and liver, a transcript distribution that is similar, but not identical, to that seen for the mammalian *FABP6/Fabp6* gene.

Introduction

Intracellular lipid-binding proteins (iLBPs) are encoded by a highly conserved multigene family, and include fatty acid-binding proteins (FABP/Fabps), cellular retinol-binding proteins (CRBPs) and the cellular retinoic acid-binding proteins (CRABPs) [1, 2]. Currently, 16 paralogous iLBP genes have been identified in animals, but no member of this multigene family has thus far been identified in plants and fungi. Schaap *et al.* [2] have therefore suggested that the first iLBP gene emerged after the divergence of animals from plants and fungi approximately 930 million years ago. This ancestral iLBP gene presumably then underwent a series of duplication events followed by sequence divergence, giving rise to the extant iLBP multigene family.

To date, 11 isoforms of FABP/Fabps or their genes, or both, have been identified in vertebrate species [3]. Originally, these proteins were named according to the tissue from which they were initially isolated, *e.g.*, liver-type fatty acid-binding protein (L-FABP), brain-type fatty acid-binding protein (B-FABP), intestinal-type fatty acid-binding protein (I-FABP), *etc.* However, this nomenclature has become confusing because different types of FABPs have been isolated from the same tissue, and some orthologous FABPs from different species exhibit distinctly different tissue-specific patterns of distribution [4, 5]. Furthermore, two so-called liver-type *fabp* genes, *fabp1a* and *fabp1b* (based on phylogenetic analysis and conserved gene synteny), are not expressed in the liver of teleost

fishes [6]. In this paper, we have used an alternative nomenclature proposed by Hertzel and Bernlohr [4], which uses numerals to distinguish the FABP proteins and genes corresponding to the chronological order of their discovery, *e.g.*, FABP1 (liver-type FABP), FABP2 (intestinal-type FABP), FABP3 (heart-type FABP), *etc.* We have also followed the gene and protein designations for mammalian and teleost fish genes and proteins according to the recommendations of the Zebrafish Model Organism Database (http://zfin.org), in which zebrafish genes and proteins are represented as *fabp6* and Fabp6, respectively, human genes and proteins are given in upper-case letters, e.g., *FABP6* and FABP6, respectively, and the mouse gene is designated *Fabp6* and its protein FABP6.

Phylogenetic studies have identified three main groups for the FABPs: group 1 includes FABP1, FABP6 and Fabp10 (Fabp10 has only been found in non-mammalian vertebrates), group 2 consists of a single protein, FABP2, and group 3 consists of FABP4, FABP5, FABP8, FABP9 and Fabp11 (Fabp11 may be unique to teleost fishes [3]). Schaap *et al.* [2] estimate that the FABPs from group 1 diverged from the last common ancestral FABP gene approximately 679 million years ago.

Although the first FABP, FABP1, was described almost four decades ago [7], and extensive studies have focused on the tissue distribution and binding activities and regulation of *FABP* genes, including FABP gene knock-out experiments [8], our understanding of the physiological function(s) of these proteins remains limited or, in many cases, unknown. However, sufficient evidence exists to strongly suggest the following roles for FABPs: (a) uptake of fatty acids across the plasma membrane and transport to various subcellular organelles; (b) modulation of activity of enzymes involved in fatty acid metabolism; (c) protection

of enzymes and membranes from the detergent effects of excess fatty acids by sequestering them, and (d) modulation of cell growth and differentiation by transport of fatty acids to the nucleus where they activate specific gene transcription [4, 8, 9].

Here we report studies on the *fabp6* gene from zebrafish, the first *fabp6* gene described for non-mammalian vertebrates. Previous work has reported the cloning and sequencing of mammalian FABP6/Fabp6 genes and cDNAs, and their expression in mammalian species including human [10], rat [11-13], mouse [14, 15] and pig [16]. Over the years, FABP6 has been given a variety of names, such as the ileal lipid-binding protein, intestinal 15-kDa protein, ileal bile acid-binding protein and gastrotropin, reflecting the speculations of authors on its intracellular function(s). In vitro binding assays revealed a surprisingly low affinity of recombinant-derived human FABP6 and rat Fabp6 for long chain fatty acids, such as palmitate and oleate, despite these proteins having a common threedimensional structural motif with other FABP/Fabps known to bind long chain fatty acids [10, 17]. Work by Gong et al. [12] suggests that the ligands of FABP6 are bile salts and that FABP6 is involved in their uptake from the ileal epithelium. However, other studies have detected mammalian FABP6/Fabp6 gene transcripts and encoded protein in the ovary and steroid endocrine cells of the adrenal gland, leading to speculation that FABP6 may also function in steroid metabolism [11]. Comparative studies of FABP6/Fabp6/fabp6 gene expression in mammals and teleost fishes may provide additional evidence for the role of this protein in cellular physiology. In this paper, we describe the structure of the zebrafish fabp6 gene, its linkage group (chromosome) assignment, conserved gene synteny with

mammalian orthologs, and the tissue-specific distribution of the *fabp6* gene transcripts in embryos, larvae and adults.

Material and Methods

Husbandry of zebrafish

The AB strain of zebrafish was used throughout this work and maintained according to established procedures [27]. Experimental protocols were reviewed by the Animal Care Committee of Dalhousie University in accordance with guidelines set down by the Canadian Committee on Animal Care.

Nucleotide sequence of zebrafish fabp6 cDNA and gene

We retrieved а previously uncharacterized Ensembl gene (ENSDAR00000044566) by a BLASTn search of the zebrafish genome sequence database at the Wellcome Trust Sanger Institute (version Zv7, scaffold 296.3, http://www.ensembl.org/Danio rerio/index.html) using NM 001002076 (GenBank accession number) as the query sequence. This sequence was also used in BLASTn searches for other ESTs coding for zebrafish Fabp6. Based on the NM 001002076 sequence, primers were designed for RT-PCR amplification of this transcript from total RNA extracted from a whole adult zebrafish of strain AB (forward primer, 5'-CTC TTC TTC TCC GCT CAA-3'; reverse primer, 5'-ATC AGT TTA GCT CGT ACA-3'). The resulting product of expected size as estimated by agarose gel electrophoresis was cloned into the pGEM-T vector (Promega, Madison, WI, USA) and six clones sequenced. To identify SNPs, the cDNA sequences obtained by us were compared to the coding sequence of zebrafish fabp6 gene retrieved from the Zebrafish Genomic Sequence Database at the Wellcome Trust Sanger Institute by alignment using CLUSTALW [28]. The

molecular mass and isoelectric point of the Fabp6 polypeptide encoded by clone NM_001002076 was determined using the program at http://ca.expasy.org/tools/pi_tool.html.

Phylogenetic analysis

Sequence alignment and determination of percentage amino acid sequence identity of FABP/Fabp sequences from zebrafish and other vertebrates was performed using BIOEDIT (v.7.0.9) [29]. Phylogenetic analysis was performed using CLUSTALW [28] to generate a neighbour-joining tree. Bootstrap values were based on 100 replicates.

Linkage group (chromosome) assignment by radiation hybrid mapping of the zebrafish fabp6 gene

Radiation hybrids of the LN54 panel were used to assign the *fabp6* gene to a specific zebrafish linkage group according to the protocol described by Hukriede *et al.* [21]. Two primers were designed (forward primer, 5'-TAG GCA AAG AGA GCC ACA TGC AGA-3'; reverse primer, 5'-TGC TCA AAT CCT GAC ACC ATG GAC-3') to PCR-amplify a portion of the zebrafish *fabp6* gene from genomic DNA samples isolated from the LN54 hybrid panel using Platinum PCR Super Mix (Invitrogen, Burlington, Ontario, Canada).

Whole-mount in situ hybridization to zebrafish embryos and larvae

Whole-mount *in situ* hybridization using a cloned *fabp6* cDNA to generate an antisense riboprobe was performed according to the methods described previously [30].

Detection of fabp6 transcripts in adult zebrafish tissues by RT-PCR

RT-PCR was used to determine the tissue distribution of *fabp6* transcripts in RNA extracted from tissues of adult zebrafish. RNA was extracted from tissue using Trizol reagent (Invitrogen). Following synthesis of cDNA using the Omnicript RT kit (Qiagen, Mississauga, Ontario, Canada), the zebrafish *fabp6* transcripts were amplified by PCR from total RNA extracted from various tissues using the forward primer, 5'-TAG GCA AAG AGA GCC ACA TGG AGA-3', and the reverse primer, 5'-GCG GTT AAA CCT TCT TGC TTG TGC-3', according to the protocol described by Liu *et al.* [23]. The constitutively expressed gene for elongation factor 1 α (*ef1\alpha*) was used as a positive control to assay the integrity of RNA extracted from each tissue. The primers and RT-PCR conditions employed have been described previously [31].

Detection of fabp6 transcript in adult sections of zebrafish by in situ hybridization

A synthetic antisense probe, 5'-GTA CTG GGT CCA TGT GAA GTC ATC TCC GTT C-3', was used for *in situ* hybridizations to detect *fabp6* transcripts in sections of adult zebrafish according to the method described by Denovan-Wright *et al.* [32].

Results and Discussion

Identification of zebrafish cDNA and genomic fabp6 sequences

Following BLAST searches of GenBank at the National Center for Biotechnology Information (NCBI), we identified an expressed sequence tag (EST) (GenBank accession number: NM 001002076) [3] for which the deduced amino acid sequence showed highest percentage sequence identity to the amino acid sequences of mammalian FABP6/Fabp6s (see below). Using this EST sequence as a query, we retrieved numerous other ESTs coding for zebrafish Fabp6 from NCBI and the sequence for the zebrafish fabp6 gene from the genomic DNA assembly Zv7, scaffold 296.3 (ENSDARG00000044566), at the Wellcome Trust Sanger Institute (http://www.ensembl.org/Danio rerio/index.html). In order to generate a hybridization probe for further study of the tissue-specific distribution of fabp6 transcripts in zebrafish embryos, we amplified the fabp6 transcript by RT-PCR of total RNA extracted from a whole adult zebrafish. The resulting DNA of the expected size was cloned, and six independent clones were sequenced and found to be identical (Genbank accession numbers EU665309 – EU665314). Five single nucleotide polymorphisms (SNP) were seen between the sequence of the fabp6 transcripts cloned by us and the coding sequence of the *fabp6* gene (Fig. 1). Only one SNP in the coding sequence, located at nucleotide (nt) position +1633 in Fig. 1, changed the deduced amino acid sequence of Fabp6, a change of valine (GTC) to isoleucine (ATC). We attribute the five SNPs to differences between established strains in zebrafish. The coding sequence derived from the fabp6 gene at the Wellcome Trust Sanger Institute was derived from the Tuebingen strain of zebrafish, while the sequences for the fabp6 cDNA generated in this study were derived from the AB strain of zebrafish (see http://zfin.org for strain details).

The coding sequence of the *fabp6* gene contained an open reading frame of 393 bp (not including the stop codon), with 5' and 3' untranslated regions of 50 and 74 bp, respectively (Fig. 1). The open reading frame codes for a polypeptide of 131 amino acids with a molecular mass of 14 406 Da and an isoelectric point of 6.59. With the exception of some Fabp10s, which have an isoelectric point of 8.8-9.0, all other FABPs have isoelectric points of approximately 6 [18].

The zebrafish *fabp6* gene consists of four exons of 112, 174, 89 and 141 bp, coding for 22, 59, 30 and 20 amino acids, respectively. The intron/exon structure of the *fabp6* gene from zebrafish is consistent with that of all other *fabp* genes studied to date [1] with the exception of the muscle-type FABP (M-FABP) genes from desert locust [19], which lacks intron II, and the *fabp1b* gene from zebrafish, which contains an additional intron in the 5'-untranslated region [6]. Each of the intron/exon splice junctions in the zebrafish *fabp6* gene conform to the GT/AG rule proposed by Breathnach and Chambon [20].

Alignment of the zebrafish Fabp6 sequence with mammalian FABP6 sequences (human, rat, mouse and pig) and to paralogs of other zebrafish Fabps and human FABPs showed that zebrafish Fabp6 shared greatest sequence identity with human (55.3%), mouse (50%), rat (50%) and pig (49.2%) FABP6 sequences (Fig. 2). Sequence identity of the zebrafish Fabp6 with paralogous zebrafish Fabps and human FABPs varied from 42.1% to 21.8%. Phylogenetic analysis revealed the inclusion of zebrafish Fabp6 with human, rat, mouse and pig FABP6s in a distinct clade with a robust bootstrap value of 99/100 (Fig. 3). The phylogenetic tree indicates a closer evolutionary relationship between FABP6s and FABP1s, and a more distant relationship between FABP6s and FABP7s, a finding consistent with early phylogenetic studies of rat and human FABP6 and FABP1 [10, 12]. The sequence alignment (Fig. 2) and phylogenetic analysis (Fig. 3) strongly suggests that the ESTs and genomic sequence retrieved from DNA assembly Zv7, scaffold 296.3 (Wellcome Trust Sanger Institute zebrafish genome sequence) described above, code for Fabp6 in zebrafish.

114

Linkage group assignment of the zebrafish fabp6 gene by radiation hybrid mapping and its conserved gene synteny with mammalian FABP6/Fabp6 genes

To provide additional evidence that the gene located on the DNA assembly Zv.7, scaffold 296.3, indeed codes for zebrafish Fabp6, we determined the linkage group (chromosome) assignment of the zebrafish fabp6 gene and examined its conserved gene synteny with the human, rat and mouse FABP6/Fabp6 genes. Using the LN54 panel of radiation hybrids [21] and specific primers to exon 2 and introns 2, respectively (see Fig. 1 and experimental procedures), the zebrafish fabp6 gene was mapped to linkage group (chromosome) 21 at a distance of 26.79 cR from the marker fc08c06 with a LOD (logarithm of the odds [to the base 10]) of 10.8 (mapping data available at http://dir.nichd.nih.gov/lmg/devb.htm). This result is consistent with the chromosomal location of fabp6 on Zv6 at the Wellcome Trust Sanger Institute database, but not with the latest version, Zv7, which places the zebrafish fabp6 gene on chromosome 3. We have previously observed incompatibilities between radiation hybrid mapping data for other zebrafish fabp genes and their chromosomal assignment in the Wellcome Trust Sanger Institute genome database for the zebrafish. Later, versions of the zebrafish genome sequence have been corrected in agreement with the chromosomal assignment of fabp genes by radiation hybrid mapping.

The conserved gene synteny between the zebrafish *fabp6* gene on chromosome 21 and human *FABP6* gene on chromosome 5 is extensive (Table 1). Conserved gene synteny was also evident between the zebrafish *fabp6* gene and the *Fabp6* genes on rat chromosome 10 and mouse chromosome 11. Not all the genes that show conserved gene synteny between zebrafish chromosome 21

and human chromosome 5 are located on rat chromosome 10 and mouse chromosome 11. Other genes are located on rat chromosomes 2, 17, 18 and 20, and mouse chromosome, 13, 15 and 18, suggesting chromosomal rearrangements or translocations in these regions after divergence of the human and rodent lineages. Despite these chromosomal rearrangements, the conserved gene synteny shown in Table 1 strongly indicates that a common linkage group containing the *FABP6/Fabp6/fabp6* gene was inherited from a common ancestor of fishes and mammals. The conserved gene synteny (Table 1), sequence identity (Fig. 2) and phylogenetic analysis (Fig. 3) provide compelling evidence that the putative zebrafish *fabp6* gene described here and the mammalian *FABP6/Fabp6*

Distribution of fabp6 gene transcripts in zebrafish embryos and larvae

To determine the spatio-temporal distribution of *fabp6* transcripts during zebrafish embryonic and larval development, we performed whole-mount *in situ* hybridization to zebrafish embryos and larvae at various developmental stages (Fig. 4). *fabp6* transcripts were not detected in embryos at 48 h postfertilization (hpf), but a very strong hybridization signal was detected in the distal region of the zebrafish intestine at 72 hpf (Fig. 4A) indicating that initiation of *fapb6* gene transcription occurred between 48 and 72 hfp. The distribution of *fabp6* transcripts remained constant in the distal region of the intestine of zebrafish larvae from 3 to 7 days postfertilization (Fig. 4A-C). A transverse section of a 4-day-old larva showed the presence of *fabp6* transcripts located predominately in epithelial cells of the intestine (Fig. 4B).

To our knowledge, only two studies have investigated the tissue-specific distribution of FABP6/Fabp6 transcripts during embryogenesis [14,15]. In the mouse, Sacchettini et al. [14] showed by dot-blot hybridization that no Fabp6 transcripts were detected in any tissues during fetal life, or throughout the suckling period of 1-12 postnatal days. Mouse Fabp6 transcripts were first detected and restricted to the ileum at the beginning of the suckling/weaning transition at postnatal days 12-14. In contrast, Crossman et al. [15] did detect Fabp6 transcripts in mouse embryos. They used Northern blot analysis and quantified mRNA steady-state levels by scanning autoradiograms of RNA extracted from total intestine and sections along the entire length of the intestine (*i.e.*, from the gastroduodenal junction to the rectum). Fabp6 transcripts were first detected in RNA from total intestine at E18 which is the stage at which the 'proximal-to-distal wave of cytodifferentiation of the pseudo-stratified gut epithelium to a monolayer had reached the ileum' [14]. During postnatal development, Fabp6 transcripts were restricted to the distal third of the small intestine and cecum. No Fabp6 transcripts were detected in the duodenum, jejunum or 12 other extraintestinal tissues (the latter tissues were not specified). The transcriptional initiation of the zebrafish fabp6 gene in the distal region of the intestine at around 72 hpf, prior to hatching (Fig. 5), occurs at approximately the same developmental stage as the transcription initiation of the mouse Fabp6 gene in the ileum at E18 [14].

Tissue-specific distribution of fabp6 gene transcript in adult zebrafish

We explored the tissue-specific distribution of *fabp6* transcripts in adult zebrafish by RT-PCR amplification from total RNA extracted from various tissues and by *in situ* hybridization of a *fabp6*-specific antisense oligonucleotide probe to

sections of adult zebrafish. A *fabp6*-specific RT-PCR product of expected size was amplified from total RNA extracted from liver, heart, intestine, ovary and kidney (Fig. 5, top panel). No *fabp6*-specific RT-PCR product was amplified from total RNA extracted from skin, brain, gill, eye or muscle. As a positive control to determine the integrity of the RNA samples used in these assays, transcripts for the constitutively expressed elongation factor 1α (*ef1* α) gene were amplified by RT-PCR. A product of the expected size was generated from RNA extracted from all tissues assayed (Fig. 5, bottom panel).

In situ hybridization of an antisense *fabp6* oligonucleotide probe to adult zebrafish sections revealed an intense hybridization signal in the distal region of the intestine (Fig. 6, 1A) and in the adrenal homolog of the kidney (Fig. 6, 1D). Less-intense hybridization signals were observed in the liver (Fig. 6, 1B) and the ovary (Fig. 6, 1C). Despite the difference in sensitivity of the two methods employed, the tissue distribution of *fabp6* transcripts in adult zebrafish assayed by RT-PCR and by *in situ* hybridization was identical.

In adult mammals, the reported tissue distributions of *FABP6/Fabp6* gene transcripts and its protein have varied, probably due to the assay techniques used. For example, Fujita *et al.* [10] used Northern blot analysis to detect a single-sized *FABP6* transcript in RNA extracted from the terminal region of the human ileum, whereas RT-PCR generated an abundant *FABP6*-specific product from total RNA extracted from the ileum, and to a much lesser extent from RNA extracted from the human ovary and placenta. Unfortunately, the authors do not state whether other tissues were assayed by RT-PCR in which *FABP6* transcripts were not detected. Rat *Fabp6* transcripts were detected by Northern blot analysis in RNA extracted from the ileum and ovary, but not in RNA extracted from the stomach, jejunum,

colon. adrenal. heart. or liver [12]. Iseki al. [11] used brain. et immunocytochemistry to localize the rat Fabp6 protein and in situ hybridization to localize Fabp6 transcripts to the enterocytes of the ileum, luteal cells of the ovary and a subpopulation of steroid endocrine cells of the adrenal gland. Sato et al. [13] also detected rat FABP6 in the adrenal gland and ovary. In adult mouse, Fabp6 transcripts were only detected by blot hybridization in the intestine, and not in the liver, stomach, pancreas, kidney, spleen, testis, skeletal muscle, heart or lung [14].

With the exception of one report [12], these studies consistently show that the FABP6/Fabp6 gene transcripts are expressed at high levels in the ileum and to a lesser extent in the ovary and adrenal gland of adult mammals. In zebrafish, we showed by RT-PCR and by in situ hybridization that fabp6 transcripts were detected at high levels in the distal region of the intestine, the tissue homologous to the mammalian ileum. The presence of *fabp6* transcripts suggests that Fabp6 may well play a role in the uptake of lipids from the distal region of the zebrafish intestine, which is similar to the suggested role for FABP6 in the uptake of bile salts from the mammalian ileum [11, 12]. Zebrafish fabp6 transcripts were shown by RT-PCR assay (Fig. 5) and by in situ hybridization (Fig. 6) to be abundant in the ovary and kidney of adult zebrafish, similar to the distribution of mammalian FABP6/Fabp6 transcripts, which are also generally found in the ovary and adrenal gland. In fishes, the adrenal homolog is not as compact as the adrenal gland found in mammals. In fishes, adrenal tissue exists as aminergic chromaffin and interregnal cells mostly inside the head kidney with the two tissues being either mixed, adjacent, or completely separated [22]. The distribution of the hybridization signal for zebrafish fabp6 transcripts in the adrenal homolog of the kidney (Fig. 6, 1D) is consistent with the structure of the adrenal homolog in teleost fishes. With the

exception of the zebrafish liver and heart, the overall pattern of adult tissue distribution of zebrafish *fapb6* and mammalian *FABP6/Fabp6* gene transcripts, and the transcriptional initiation of these genes at similar embryonic stages of development, is surprisingly concordant, in contrast to some other members of the multigene family of iLBP genes (*e.g., fabp1a/b, fabp10, fabp11, rbp2*) [3,6,16,23,24]. As FABP6 has been implicated in human colorectal cancer [25] and type 2 diabetes [26], zebrafish may serve as a useful model experimental system to investigate the role of FABP6 in these disease states.

Acknowledgements

The authors wish to thank Santhosh Karanth (Department of Biology, Dalhousie University, Halifax, Canada) and Paul Wright (University of Virginia Health Sciences Center, Charlottesville, VA, USA) for technical assistance during the course of these studies. This work was supported by funds from the Natural Sciences and Engineering Research Council of Canada (to J.M.W.), the Canadian Institutes of Health Research (to E.D.W.), and the National Institute of Health, the European Commission as part of the ZF-Models integrated project in the 6th Framework Program (to B.T. and C.T.). F.A-C. was a recipient of a scholarship from FAPESP, "Fundação de Amparo à Pesquisa do Estado de São Paulo", Brazil.

References

[1] Bernlohr DA, Simpson MA, Hertzel AV & Banaszak LJ (1997) Intracellular lipidbinding proteins and their genes. Annu Rev Nutr 17, 277-303.

120

- [2] Schaap FG, Van der Vusse GJ & Glatz JFC (2002) Evolution of the family of intracellular lipid binding proteins in vertebrates. *Mol Cell Biochem* 239, 69-77.
- [3] Agulleiro MJ, André M, Morais S, Cedrà J & Babin PJ (2007) High transcript level of fatty acid-binding protein 11 but not of very low-density lipoprotein receptor is correlated to ovarian follicle atresia in a teleost fish (*Solea senegalensis*). *Biol Reprod* **77**, 504-516.
- [4] Hertzel AV & Bernlohr DA (2000) The mammalian fatty acid-binding protein multigene family: molecular and genetic insights into function. *Trends Endocrine Met* **11**, 175-180.
- [5] Zimmerman AW & Veerkamp JH (2002) New insights into the structure and function of fatty acid-binding proteins. *Cell Mol Life Sci* **11**, 1096-1116.
- [6] Sharma MK, Liu R-Z, Thisse C, Thisse B, Denovan-Wright EM & Wright JM (2006) Hierarchical subfunctionalization of *fabp1a*, *fabp1b*, and *fabp10* tissuespecific expression may account for retention of these duplicated genes in the zebrafish (*Danio rerio*) genome. *FEBS J* **273**, 3216-3229.
- [7] Ockner RK, Manning JA, Poppenhausen RB & Ho WK (1972) A binding protein for fatty acids in cytosol of intestinal mucosa, liver, myocardium, and other tissues. *Science* **177**, 56-58.
- [8] Haunerland NH & Spener F (2004) Fatty acid-binding proteins insights from genetic manipulations. *Prog Lipid Res* 43, 328-349.
- [9] Storch J & Thumser AEA (2000) The fatty acid transport function of fatty acidbinding proteins. *Biochim Biophys Acta* **1486**, 28-44.
- [10] Fujita M, Fujii H, Kanda T, Sato E, Hatakeyama K & Ono T (1995) Molecular cloning, expression and characterization of a human intestinal 15-kDa protein. *Eur J Biochem* 233, 406-413.
- [11] Iseki S, Amano O, Kanda T, Fujii H & Ono T (1993) Expression and localization of intestinal 15 kDA protein in the rat. *Mol Cell Biochem* **123**, 113-120.
- [12] Gong Y-Z, Everett ET, Schwartz DA, Norris JS & Wilson FA (1994) Molecular cloning, tissue distribution, and expression of a 14-kDa bile acid-binding protein from rat ileal cytosol. *Proc Natl Acad Sci* **91**, 4741-4745.
- [13] Sato E, Fujii H, Fujita M, Kanda T, Iseki S, Hatakeyama K, Tanaka T & Ono T (1995) Tissue-specific regulation of the expression of rat intestinal bile acidbinding protein. *FEBS Lett* **374**, 184-186.

- [14] Sacchettini JC, Hauft SM, Van Camp SL, Cistola DP & Gordon JI (1990) Developmental and structural studies of an intracellular lipid binding protein expressed in the ileal epithelium. *J Biol Chem* 265, 19199-19207.
- [15] Crossman MW, Hauft, SM & Gordon JI (1994) The mouse ileal lipid-binding protein gene: a model for studying axial patterning during gut morphogenesis. *J Cell Biol* **126**, 1547-1564.
- [16] Walz DA, Wider MD, Snow JW, Dass C & Desiderio DM (1988) The complete amino acid sequence of porcine gastrotropin, an ileal protein which stimulates gastric acid and pepsinogen secretion. *J Biol Chem* 28, 14189-14195.
- [17] Kanda T, Odani S, Tomoi M, Matsubara Y & Ono T (1991) Primary structure of the 15-kDA protein from rat intestinal epithelium. *Eur J Biochem* **197**, 759-768.
- [18] Denovan-Wright EM, Pierce M, Sharma MK & Wright JM (2000) cDNA sequence and tissue-specific expression of a basic liver-type fatty acid binding protein in adult zebrafish (*Danio rerio*). *Biochim Biophys Acta* 1492, 227-232.
- [19] Wu Q, Andolfatto P & Haunerland NH (2001) Cloning and sequence of the gene encoding the muscle fatty acid binding protein from desert locus, *Schistocerca gregaria*. *Insect Biochem Mol Biol* **31**, 553-563.
- [20] Breathnach R & Chambon P (1981) Organization and expression of eukaryotic split genes coding for proteins. Annu Rev Biochem 31, 349-383.
- [21] Hukriede NA, Joly L, Tsang M, Miles J, Tellis P, Epstein JA, Barbazuk WB, Li FN, Paw B, Postlethwait JH, Hudson TJ, Zon LI, McPerson JD, Chevrette M, Dawid IB, Johnson SL & Ekker M (1999) Radiation hybrid mapping of the zebrafish genome. *Proc Natl Acad Sci* **96**, 9745-9750.
- [22] Gallo VP & Civinini A (2003) Survey of the adrenal homolog in teleosts. Int Rev Cytology 230, 89-187.
- [23] Liu R-Z, Denovan-Wright EM & Wright JM (2003) Structure, linkage mapping and expression of the heart-type fatty acid-binding protein gene (*fabp3*) from zebrafish (*Danio rerio*). *Eur J Biochem* 270, 3223-3234.
- [24] Liu R-Z, Denovan-Wright EM, Degrave A, Thisse, C, Thisse B & Wright JM (2004) Spatio-temporal distribution of cellular retinol-binding protein gene transcripts (*CRBPI* and *CRBPII*) in the developing and adult zebrafish (*Danio rerio*). *Eur J Biochem* 271, 339-348.

- [25] Ohmachi T, Inoue H, Mimori K, Tanaka F, Sasaki A, Kanda T, Fujii H, Yanaga K & Mori M (2006) Fatty acid binding protein 6 is overexpressed in colorectal cancer. *Clin Cancer Res* **12**, 5090-5095.
- [26] Fisher E, Nitz I, Lindner I, Rubin D, Boeing H, Möhlig M, Hampe J, Schreiber S, Schrezenmeir J & Döring F (2007) Candidate gene association study of type 2 diabetes in a nested case-control study of the EPIC-Potsdam cohort role of fat assimilation. *Mol Nutr Food Res* **51**, 185-191.
- [27] Westerfield M (1995) The zebrafish Book: a Guide for the Laboratory Use of Zebrafish (Danio rerio), 3rd edn. University of Oregon Press, Eugene, USA.
- [28] Thompson JD, Gibson TJ, Plewniak F, Jeanmougin F & Higgins HG (1997) The CLUSTALW windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Res* 24, 4876-4882.
- [29] Hall T (1999) BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucleic Acids Symp Ser* **41**, 95-98.
- [30] Thisse C & Thisse B (2008) High-resolution *in situ* hybridization to wholemount zebrafish embryos. *Nat Protoc* **3**, 59-69.
- [31] Pattyn F, Robbrecht P, Speleman F, De Paepe A & Vandesompele (2006) RTPrimerDB: the real-time PCR primer and probe database, major update 2006, *Nucleic Acids Res* 34 (Database issue), D684-688.
- [32] Denovan-Wright EM, Newton RA, Armstrong JM, Babity JM & Robertson HA (1998) Acute administration of cocaine, but not amphetamine, increases the level of synaptotgmin IV mRNA in the dorsal striatum of rat. *Mol Brain Res* 55, 350-354.

Table, Figures and Legends

Chromosomal Position											
Genes	Zebrafish	Human		Rat				Mouse			
	LG	5	2	10	17 /20	18	11	13	15	18	
c6 c7 rpl37 fgf10 taf9 f2rl11 thbs4 bhmt glrx ell2 pcsk1 rnf14 cdc23 pou4f3 sept8 skp1a vdac1 cnot8 ddx46 pdim4 rapgef6 sara2 tcf7 zcchc10 spry4 zmat2 rbm22 larp1 sap300 rnf145 fabp6 sgcd mat2b drd1 fgfr4 rars ubtd2 cnot6 nola2	21 21 21 21 21 21 21 21 21 21	$\begin{array}{c} 5p13\\ 5p13\\ 5p13\\ 5p13.\\ 5p13.\\ 5p13.p12\\ 5q11.2-q13.1\\ 5q13\\ 5q13\\ 5q13\\ 5q13\\ 5q14\\ 5q15\\ 5q15\\ 5q15- 5q15- 5q15- 5q15- 5q15- 5q15- 5q15- 5q31- 15q31- 15q31- 15q31- 15q31- 15q31- 15q31- 15q33- 5q33- 35q33- 35q33- 35q33- 35q33- 35q33- 35q33- 35q35- 356- 356- 355- 356- 355- 356- 355- 356- 355$	2q16 2q16 2q15-q16 2q12 2q12 2q12 2q11 2q11 2q11-q12 - -	10q22 10q22 10q22 10q22 10q22 10q22 10q22 10q22 10q22 10q22 10q22 10q22 10q22 10q22 10q22 10q22 10q22 10q22 10q22 10q22	17p14 - - 20p12 17p14 -	18p11 18p12 18p11 - - 18p11 18q12.1 -	11 28.5cM 11 31.0cM 11 29.0cM 11 B1.3 11 B1.3 11 B1.3 11 B1.3 11 28.5cM 11 28.5cM 11 B2 11 B1.1 11 24.0cM 11 B1.2 11 B1.1 11 A4 11 A4 11 A4 11 B1.2 11 28.5cM	13 75.0cM 13 D1 13 46.99cM 13 D1 13 44.0cM 13 C1 13 44.0cM 13 B2	15 3.0cM 15 3.0cM 15 A1	18 17.0cM 18 17.0cM 18 24.0cM 18 18.0cM 18 B2 18 D2	

Table 1:	Conserved	gene synteny	between	zebrafish	i linkage	group	(chromosome) 21	and	human	chromos	some
5, rat chr	omosome 10) and mouse c	hromosor	ne 11.								

tccaaacatatcatgaccctgagcatgctgccaaatta -560 -480 gttaaaatgtactttaaggacaacagagtgaatgttttggagtggccatcacaaagccctgagctcaatcctatagaaa atttgtgggcagagttgaaaaagcttgtgcgagcaaaacagccaacaaatttgacttagttacaccaattccgtcagga -400 ggaatgagccaaaattcgttcaaactattgtgagaagcttgtggaaggatacccaaaacatttgaccaaagttatacag -320 -240 tttaaaatcaaagctaaaaaattcttatggtaatttttttgtattgtattataattgtattgcacgattataaagtgat ${\tt tttgcccatgtatggcatatcatcaaccatgtattttgcccatgtaaggcatatcatcaccacgtgagcagccgt}$ -160 gcggtcatgcactttatgacttactctattttgtggggccatcataaatgtttatggcttgattatgatgctggtgtac-80 $\verb|cactaaggccatgagaaagagtcagagttgaggaacagttgggta \underline{\texttt{TATAA}} a \verb|caccgcatacttcacccatcagtacctc||$ -1 AGCTCTCAACCCGCTCTTCTTCTCCGCTCAATCAACACCAAAAACC ATG GCT TTC AAC GGC AAG TGG GAA A +70 M A F N G K W E CC GAA TCT CAG GAG GGA TAT GAA CCA TTC TGC AAA CTG ATC G +112 TΕ E G Y E P F S 0 C K T. Т gtgaggct... INTRON I... cacctcag (+113 - +1576)GT ATC CCT GAT GAT GTC ATC GCA AAG GGC CGT GAC TTC AAG CTT GTG ACA GAG ATC GTC +1635G I P D D V I A K G R D F K L V T E I V/I CAG AAC GGA GAT GAC TTC ACA TGG ACC CAG TAC TAC CCC AAT AAC CAT GTT GTG ACC AAC +1694 O N G D D F Т W Т 0 Υ Y P Ν N Η V V Т N Α AAA TTC ATC GTA GGC AAA GAG AGC GAC ATG GAG ACT GTA GGA GGG AAG AAA TTT AAG +1751 Т Κ F Т V G ΚE S D М E V G G K K F Κ gtgtggca... INTRON II... ttcaccag (+1752 - +3729)С GGC ATA GTT TCC ATG GAA GGA GGC AAG CTG ACC ATA AGC TTC CCC AAA TAT CAA CAA ACA +3789 G I V S M E G G K L T I S F P K Y Q Q т ACT GAG ATC AGC GGT GGA AAG CTG GTG GAG +3819TEISGGKLV E (+3820 - +4140)gtgagata... INTRON III... ttttcag ACC TCC ACA GCC AGT GGC GCC CAG GGT ACC GCT GTT CTT GTG CGC ACA AGC AAG AAG GTT +4200 T S T A S G A Q G T A V L VR Т K S Κ т G TAA CCGCATTGTTGAGATGAATGACCCTTTTCTGCTCTGACACTCAATTTACTGAATAAAGTGTACGAGCTAAACTGA +4278 * TAAA +4282

Figure 1. The nucleotide sequence of the zebrafish *fabp6* gene. The coding sequence is shown in uppercase letters with the deduced amino acid sequence indicated below. The size of each intron is shown with each exon/intron splice junctions (gt-ag) highlighted in bold font and underlined and the 5' upstream sequence of the *fabp6* gene is presented by lowercase letters with a putative TATA box in uppercase letters, underlined and in bold font. Single nucleotide polymorphisms (SNPs) based on differences between cDNA sequences from the Tuebingen and AB strains of zebrafish are positioned above the genomic sequence and in bold font. A polyadenylation signal sequence, "AATAAA", is underlined and in bold font.

ZF	FABP6	MAFNGKWETESQEGYEPFCKLIGIPDDVIAKGRDFKLVTEIVQNGDDFTWTQYYPNNHVVTNKFIVGK
HU	FABP6	TF.MEKN.DE.MLSSE.A.NIVQ.D.QS.H.SGG.TMT
MO	FABP6	SY.FEKN.DE.M.RL.L.GERNIIVQ.D.QS.S.SGGNIMSTI
RA	FABP6	TY.FEKN.DE.M.RL.L.EERN.IIVQ.D.ENS.S.SGGNIMSTI
ΡI	FABP6	TY.IEKN.DE.M.RLAL.S.A.D.A.NL.IIS.VK.D.QNS.QGG.SIT.TI
ΖF	FABP1B	S.TYQLF.E.M.AV.LM.EK.I.S.SEENQ.KV.VT-TGSK.L.S.TI.Q
HU	FABP1	S.SYQLQNF.A.M.AL.EEL.QK.I.G.SKH.KF.IT-AGSK.IQ.E.TE
ΖF	FABP1A	TYQLH.NF.A.M.AV.VEVEK.I.SISH.D.KKV.VT-AGTK.ILYS.TE
HU	FABP3	.VDL.T.KLVDSKNFDDYM.SL.VGFATRQVASMT.PT.I.EKIL.LKTH-STFKNTEIS.KL.V
ΖF	FABP7A	.VDCAT.KLVDSQNFDEYM.SL.VGFATRQV.NVT.PTIV.SHEKVVIKTL-STFKNTEIS.KL.E
HU	FABP2	DST.KVDRS.N.DK.MEKM.VNIVKRKLAAHDN.KLT.T.E.NKVKES-SAFRNIEVV.EL.V
HU	FABP7	.VECAT.KLTNSQNFDEYM.AL.VGFATRQV.NVT.PTVI.S.EKVVIRTL-STFKNTEIS.QL.E
ΖF	FABP3	.ADI.T.NLKESKNFDEYM.GVGFATRQVANMT.PT.I.SKEVLKTV-STFKSTEIN.KL.E
HU	FABP4	.CDV.T.KLV.S.NFDDYM.EV.VGFATRKVAGMA.PNMI.SVVI.IKSE-STFKNTEIS.L.Q
ΖF	FABP2	TT.KVDRN.NK.MEQM.VNMVKRKLAAHDN.KITLE.TK.NVKEV-STFRTLEIN.TL.V
ΖF	FABP7B	.VDC.T.KLV.SDNFDEYM.SLGFATRQV.NVT.PTLV.SKE.EKVVIKTQ-STFKNTEIS.TL.E

ZF FABP6	ESDMETVGGKKFKGIVSMEG-GKLTISFPKYQQTTEISG-GKLVETSTASGAQGTAVLVRTSKKV-	100.0%
HU FABP6	NIQ.MTAT.QVVNNHSVDVIG.VTYE.VRLA	55.3%
MO FABP6	.CE.Q.MAT.KVVAENHS.VVDIIGDVTYE.VRLA	50.0%
RA FABP6	.CE.Q.MAT.KVVADNHS.VVDIIGDVTYE.VR.A	50.0%
PI FABP6	.C.IIAT.QVVVNS.NHH.AVDVVG.VTYE.VLA	49.2%
ZF FABP1B	.A.ILT.E.V.TT.NRNKVVLNRITSILVDTNTN.L.LG.LVYK.IR.A	42.1%
HU FABP1	.CELMT.E.V.TV.QLDNVTT.KNIKSVLNDIITN.M.LGDIVFK.IRI-	39.3%
ZF FABP1A	.CELFT.DRA.TV.Q.DNAFVKGIESVLDDTISN.LSFN.IVYK.I.RRIS	36.3%
HU FABP3	.F.ET.ADDR.V.STLDVHLQKWDGQETTLVR.LIDIL.L.HGT.VCTYE.EA	26.8%
ZF FABP7A	.F.ET.ADDRHV.STLDN.VQVQRWDGKETKFVRKDM.M.L.FE.VAYE.A-	25.5%
HU FABP2	TFNYNLAD.TELR.TW.LNIGK.KRTDNGNELNTVRIDEQ.YVYE.VEAK.IFD-	24.8%
HU FABP7	.F.ET.ADDRNC.SVLDDVHIQKWDGKETNFVRKDM.M.L.FGDVVAHYE.A-	24.0%
ZF FABP3	.F.ET.ADDR.V.SVITLDLHVQKWDGKETTLLR.V.D-NN.TL.L.LGDIVST.HYV.AE	23.9%
HU FABP4	.F.EV.ADDR.V.STITLDV.VHVQKWDGKSTTIKRKRED-DVECVMK.VTST.VYERA-	22.6%
ZF FABP2	TF.YSLAD.TELT.SWVIDT.KGT.TRKDNGKVLTTVRT.VNEQSYSYD.VEAK.IF.RA-	22.6%
ZF FABP7B	.FEET.ADDRHC.ST.LLKNQ.VHVQKWDGKETTFIRKDM.MKL.FGDVEALYE.A-	21.8%

Figure 2. Sequence alignment and amino acid sequence identity of zebrafish Fabp6 and FABP6 from different species, and paralogues of other zebrafish Fabps and human FABPs. The deduced amino acids sequence of the zebrafish Fabp6 (Ensembl Peptide ID: ENSDARP00000065447) was compared to sequences of FABP6s from human (HU FABP6; U19869), mouse (MO FABP6; CAI24826), rat (RA FABP6; NP_058794), pig (PI FABP6; P10289), zebrafish Fabp paralogues, FABP1A (ZF FABP1A; DQ062095), FABP1B (ZF FABP1B; DQ062096), FABP2 (ZF FABP2; AAH75970), FABP3 (ZF FABP3; NP_694493), FABP7A (ZF FABP7A; NP_571680), FABP7B (ZF FABP7B; AAQ92970), and human FABPs, FABP1 (Hu FABP1; M10617), FABP2 (HU FABP2; M18079), FABP3 (HU FABP3; X56549), FABP4 (HU FABP4; NP_00133) and FABP7 (HU FABP7; CAI15449). Dots indicate amino acids identity. Dashes have been introduced to maximize alignment. The percentage amino acid sequence similarities between the zebrafish Fabp6 and other FABPs are shown at the end of each sequence.



Figure 3. A neighbour-joining tree showing the phylogenetic relationship of zebrafish fabp6 with selected paralogous and orthologous FABP/Fabps from zebrafish and mammals. The bootstrap values (based 100 duplicates) are indicated at the nodes. The sequences used were: zebrafish FABP6 (Ensembl Peptide ID: ENSDARP00000065447); mammalian sequences for FABP6 from human (HU FABP6, U19869), mouse (MO FABP6, CAI24826), rat (RA FABP6, NP_058794) and pig (PI FABP6, P10289); zebrafish FABP1A (ZF Fabp1A, DQ062095), FABP1B (ZF Fabp1B, DQ062096), FABP2 (ZF Fabp2, AAH75970), FABP3 (ZF Fabp3, NP_694493), FABP7A (ZF Fabp7A, NP_571680) and FABP7B (ZF Fabp7B, AAQ92970); human FABP1 (Hu FABP1, M10617), FABP2 (HU FABP2, M18079), FABP3 (HU FABP3, X56549), FABP4 (HU FABP4, NP_00133) and FABP7 (HU FABP7, CAI15449). The distinct clade of FABP6/Fabp6s is boxed in grey. Scale bar = 0.2 substitutions per site.



Figure 4. Spatio-temporal distribution of *fabp6* transcripts during zebrafish embryonic and larval development was determined by whole-mount *in situ* hybridization. *fabp6* transcripts were first detected at 72 hours postfertilization (hpf) in the distal region of the intestine (Fig. 4A) and remained confined to this region of the intestine up to 7 day postfertilization (Fig. 4C), the last time point assayed. Fig. 4B shows the distribution of *fabp6* transcripts throughout the enterocytes of the intestine in a transverse section at 4 days postfertilization.



Figure 5. RT-PCR detection of *fabp6* transcripts in RNA extracted from tissues of adult zebrafish. RT-PCR generated a *fabp6* mRNA-specific product from RNA extracted from adult zebrafish liver (L), heart (H), intestine (I), ovary (O) and kidney (K). No *fabp6* mRNA-specific product was generated by RT-PCR of RNA extracted from adult zebrafish skin (S), brain (B), gills (G), eyes (E), muscle (M), or the negative control (-) lacking total RNA derived from a whole fish. As a positive control for the integrity of each RNA template, an *ef1á* mRNA-specific product was generated from all the adult zebrafish tissues analyzed.



Figure 6. Tissue-specific detection of *fabp6* gene transcripts by *in situ* hybridization of an antisense riboprobe to sections of adult zebrafish. Panel **1** shows the distribution of the *fapb6* transcript in the distal region of the intestine (A), liver (B), ovary (C) and the adrenal homolog in the fish kidney (D). Panel **2** shows the location of distal region of the intestine (A), liver (B), ovary (C) and the adrenal homolog in the fish kidney (D) and the adrenal homolog in the fish kidney (D) in an adjacent tissue section stained with cresyl violet.

CAPÍTULO 4

Análise comparativa de expressão de transcritos gênicos em tecido cerebral de *zebrafish* (*Danio rerio*) selvagem e transgênico para o gene do hormônio do crescimento (GH), utilizando uma abordagem de DDRT-PCR A comparative expression analysis of gene transcripts in brain tissue of wild-type and GH-transgenic zebrafish (*Danio rerio*) using a DDRT-PCR approach

Abstract

Previous data on homozygote individuals of the GH-transgenic zebrafish F0104 line indicated that, despite expressing a higher amount of exogenous growth hormone in relation to hemizygote animals, they presented a significant reduced growth rate (similar to non-transgenic individuals). In the present study, hemizygote (HE) and homozygote (HO) samples of this GHtransgenic fish line were compared to non-transgenic (NT) samples of the species through a DDRT-PCR (Differential Display Reverse Transcriptase -Polymerase Chain Reaction) approach, with the goal of identifying candidate differentially expressed transcripts in brain tissue that could be involved in growth process. Two isolated fragments evidenced a high identity level with the genes that code for the proteins p300 (E1A binding protein p300) and ADCY2 (adenylate cyclase 2). Densitometric analyses of the p300 amplification product pointed to a significant lower gene expression in the transgenic genotypes, specially in HE (17.41±5.32) than in HO (32.25±3.21) samples, when compared to NT samples (49.35±12.32). Inferences on the ADCY2 transcript suggested a higher expression level in NT animals (22.73±3.80) than in HE (14.05±4.95) and HO individuals (10.63±0.67). These results corroborate previous data that evidenced that p300 and adenylate cyclase are involved in muscle formation/growth and in GH release, respectively. The obtained data can give support to further analyses into the elucidation of the mechanisms beyond vertebrate growth and GH expression gene control.

Introduction

Zebrafish (*Danio rerio*) is a well-established model organism in scientific research (Kimmel 1989, Westerfield 1995), especially in behavioral, physiologic, and developmental analyses due to several characteristics of the species. The large, robust and transparent embryos and eggs, and the rapid early development are traits that facilitate experimental manipulation and observation. Moreover, the continuous reproduction, external development, relatively short generation time, large number of progeny, and small size adults, represent important features that allow for large-scale experiments, even in reduced spaces (Kimmel 1989, Westerfield 1995, Higashijima *et al.* 1997, Detrich *et al.* 1998, Dahm *et al.* 2007, Bass & Gerlai 2008, Fadool & Dowling 2008).

The facility to manipulate this species permitted gene transfer experiments, leading to the development of several transgenic lines, which contribute to new research on vertebrate gene expression (Sprague *et al.* 2003). Growth hormone (GH) transgenic zebrafish represents one of the most explored lines of the species with an exogenous gene. There is a great interest in GH gene overexpression in fish species, since generally it can be directly related to a high growth performance of the animals in a short period of time, an important trait in aquaculture (Fletcher & Davies 1991, Hew *et al.* 1995, Zbikowska 2003, Devlin *et al.* 2004, Devlin *et al.* 2006, Figueiredo *et al.* 2007a, Rosa *et al.* 2008).

Even though some studies have shown that only a 20% increase in growth rate can be achieved by an exogenous GH application in zebrafish (Simpson *et al.* 2000, Morales *et al.* 2001, Biga & Goetz 2006), contrasting with other growth hormone-transgenic fish, as coho salmon, that are extremely fast growing (Devlin *et al.* 1994), GH-transgenic *Danio rerio* can be used as an experimental model on basic research analyses to understand vertebrate growth mechanisms and GH expression gene control. Moreover, the growth hormone is also involved in other physiological processes, since it can regulate the metabolism of several molecules, as lipids, proteins, and maintenance of all major immune cell types maintenance (Jeay *et al.* 2002), and influence the stress response and behavior (Yoshizato *et al.* 1998).

A growth hormone-transgenic line of *Danio rerio*, denominated F0104 lineage, is composed by genetically modified animals with an exogenous GH gene from the marine fish *Odonthestes argentinensis* (Figueiredo *et al.* 2007a). A significant high growth performance was evidenced in hemizygote GH-transgenic animals of this line, although individuals of the homozygote genotype did not present a differential growth rate when compared to wild-type zebrafish. However, homozygote animals presented a higher GH expression than hemizygote and non-transgenic animals, leading to the assumption that a GH excess do not correspond to a higher growth performance in zebrafish (Figueiredo *et al.* 2007b).

Due to growth differences among non-transgenic, and homozygote and hemizygote zebrafish of the F0104 line, it is expected that genes involved in growth process are differentially expressed in these animals. Therefore, the present study intended to identify candidate DETs (Differentially Expressed Transcripts) in these three different samples using a DDRT-PCR (Differential Display Reverse Transcription - Polymerase Chain Reaction) approach. The results can give new insights into the elucidation of the mechanisms beyond vertebrate growth and of the GH expression gene control.

Material and Methods

Animal samples

Adult individuals of zebrafish (*Danio rerio*) were obtained from Fundação Universidade Federal do Rio Grande (Rio Grande, RS), using the transgenic line F0104 that has the carp (*Cyprinus carpio*) β -actin promoter driving the expression of two exogenous genes - the marine silverside fish *Odonthestes argentinensis* growth hormone (msGH) gene and the jellyfish (*Aequorea victoria*) GFP (Green Fluorescent Protein) as a reporter gene to monitor for transformed zygotes (Figueiredo *et al.* 2007a). Different genotypes were obtained through crossing GH-transgenic hemizygote individuals leading to a progeny with a proportion of 25% homozygote (HO), 50% hemizygote (HE), and 25% non-transgenic (NT) animals. Genotypes were determined through UV fluorescence analysis, due to the GFP expression, with an epifluorescence microscope (excitation = 485 nm; emission = 520 nm). HO individuals were identified by presenting a higher fluorescence level.
RNA isolation

Brain tissue was collected from two animals of each genotype (HO, HE, and NT) and immediately stored at -80°C until RNA extraction. Approximately 100mg of the brain samples were mechanically homogenized with 1mL of TRizol Reagent (Invitrogen) and total RNA extraction followed the manufacturer's protocol. RNA samples were eluted in RNase-free water and quantified (NanoDrop 1000 Spectrophotometer) by measuring the optical density (OD) at 260nm. RNA purity was ensured by obtaining a 260/280nm OD ratio ≥1.80.

RT-PCR and cDNA amplification

Each RNA sample (5µg) was reverse transcribed (RT-PCR) with the commercial kit SuperScript III RT (Invitrogen) using the oligonucleotide AP (5'-GGCCACGCGTCGACTAGTAC(T)₁₇-3'), according to the manufacturer's instructions. A DDRT-PCR approach was used in order to amplify the cDNA samples. Oligonucleotides with 15-24 bp were designed based on VNTR core sequences (Table 1) and used as single primers for cDNA amplification. Each cDNA amplification reaction consisted of 1µL of cDNA, 0.2mM of primer, 1x 25mM MgCl₂ PCR buffer, 0.2mM of dNTPs, and 0.2 unit of Platinum *Taq* DNA polymerase (Invitrogen), in a final volume of 25µL. Reactions were carried out with an initial denaturation step at 95°C for 2 minutes, followed by 40 cycles at 94°C for 50 seconds, 55°C for 2 minutes, and 72°C for 50 seconds, with an additional extension step at 72°C for 5 minutes. DDRT-PCR products (10µL)

were fractionated on 6% polyacrilamide gel, stained with 0.17% silver nitrate (AgNO₃) (Sambrook & Russel 2001), and visualized under white light. The molecular weight of the amplified fragments were assigned through comparison with 1Kb DNA ladder (Invitrogen).

Densitometric analysis

The amplified fragments corresponding to putative differentially expressed transcripts were quantified by densitometry as Integrated Optical Density (IOD) using Imaging Master VDS Software version 3.2 (GE Healthcare Life Sciences). Mean and standard deviations (SD) were determined using Macrosoft Excel Software.

DNA isolation and reamplification

Selected DNA fragments that seemed to correspond to genes with differential expression among the analyzed zebrafish samples were reamplified in order to obtain a higher DNA content to be used on cloning and nucleotide sequencing. The fragments were cut from the polyacrilamide matrix and used directly (without any DNA purification) on reamplification reactions, as described in and Dakis & Kouretas (2002). The amplification reactions followed the conditions described above. PCR products were analyzed through 6% polyacrilamide gel electrophoresis.

Cloning, sequencing, and sequence analysis

PCR products were cloned into pGEM-T (Promega) vector and used to transform competent cells of the *E. coli* strain DH5 α (Invitrogen), following the manufacturers' instructions. Clones were submitted to automated sequencing on an ABI 377 Automated DNA Sequencer (Applied Biosystems) with a DYEnamic ET Terminator Cycle Sequencing kit (GE Healthcare Life Sciences), following the manufacturer's instructions, and using primers complementary to vector arms. Nucleic acid sequence database searches were performed using BLAST/N (Altschul et al. 1990) at the National Center for Biotechnology Information (NCBI) website (http://www.ncbi.nlm.nih.gov/blast). Sequence alignments were obtained by Clustal-W function (Thompson et al. 1994) and the consensus sequences were manually determined.

Results and Discussion

Muscle development and growth are dynamic processes controlled by a number of factors including growth hormone, insulin-like growth factors (IGFs), anabolic steroids, thyroid hormone, cytokines and cyclins. The GH, that plays a major role in somatic growth, is mainly produced at the anterior pituitary gland, under the control of hypothalamic factors as GHRH (GH release hormone) and somatostatin (GHIH - growth hormone inhibiting hormone) and is further released in circulation blood, acting in specific organs through its association with different receptors present at the surface of target cells (Florini *et al.* 1996, Le Roith *et al.* 2001). Nowadays, several analyses in disorders that result from under or overproduction of GH, and in transgenic animals that overexpress this hormone have been conducted in order to understand these hormone's interactions with other proteins.

Regardless the mechanisms responsible for the actions of GH, the resulting effect of its overexpression is generally an increase in muscle mass, as observed for some fish species (Nam et al. 2001, Devlin et al. 2006). However, a higher GH transgene expression level is not related to an enhancement of growth rate in other fish, as observed in the tilapia Oreochromis niloticus and O. hornorum (Rahman et al. 1998, Martínez et al. 1999). Homozygotes of the GH-transgenic Danio rerio F0104 line also demonstrated a reduced growth, similar to non-transgenic animals, even though they presented a higher GH expression (Figueiredo et al. 2007b). The different growth performance between hemizygote (HE) and homozygote (HO) GH-transgenic zebrafish of this line lead to analyses on genes that could be involved in growth process, including IGF-I (insulin-like growth factor I), GHG (GH receptor), and SOCS (suppressor of cytokine signaling). The results permitted to identify distinct expression levels of these genes among the different GH-transgenic genotypes and the non-transgenic (NT) animals (Figueiredo et al. 2007b, Studzinski 2008).

In order to implement data on differentially expressed genes in NT, HE, and HO zebrafish, total RNA of brain tissue of these three different samples were obtained and used on a DDRT-PCR approach. cDNA amplification was achieved using oligonucleotides that were designed based on VNTR (Variable Number of Tandem Repeats) core sequences that correspond to minisatellite short and highly conserved regions (Jeffreys *et al.* 1985). The use of longer

140

primers (14-24pb) on differential display strategies can lead to detection of DNA polymorphism through RAPD-like results. However, since these oligonucleotides are longer than RAPD primers, this methodology can also be effectively carried out at relatively high stringencies, thus yielding more reproducible results (Wasko & Alves-Costa*). Two of the primers (INS and HBV3) used to amplify the cDNA templates lead to the identification of presumptive differentially expressed transcripts among the analyzed samples.

DDRT-PCR using primer INS

cDNA amplifications using primer INS lead to the identification of an intense stained fragment of approximately 1.000 bp, present in NT, and in GH-transgenic HE and HO individuals of zebrafish (Figure 1a), that seemed to correspond to a transcript with differential gene expression among the three samples - the polyacrilamide electrophoresis results permitted the visualization of a fragment with distinctive stain intensities among the different samples (Figure 1a). Densitometric analyses of this amplified product, through IOD values, suggested a significant lower gene expression level in the transgenic genotypes, specially in HE (17.41±5.32) than in HO (32.25±3.21) samples, when compared to NT samples (49.35±12.32) (Figure 2a). The suggested expression levels of the analyzed transcript in HE and HO samples represent approximately 35% and 65% of the IOD value obtained for the non-transgenic samples, respectively.

Cloning and sequencing of the transcript obtained with the primer INS permitted the characterization of a 941 bp DNA fragment (Figure 3). Database searches for nucleotide similarity indicated a 95% identity level with a partial region of the gene that codes for the p300 protein of *Danio rerio* (accession

141

number XM_001332682, NM_001510744) and also a high identity with the p300 gene of several other vertebrates, as *Ctenopharyngodon idella* (81%), *Gallus gallus* (81%), and *Mus musculus* (78%). The p300, also known as E1A binding protein p300, is involved in several processes, as cellular differentiation and proliferation, cycle cell regulation, growth, apoptosis, and histone acetylation (Huret 2000, Vleugel *et al.* 2006). It has been described as a transcriptional coactivator that interacts with other proteins, leading to the activation of the transcriptional process of several genes (Huret 2000, Tu & Luo 2007), including those related to cancer (Vachtenheim *et al.* 2006, Vleugel *et al.* 2006) and neural pathologies development (Renoult *et al.* 2007, Francis *et al.* 2007), and to immune responses (Zhang *et al.* 2005).

Recently, it was proposed that the p300 is also involved in muscle tissue formation and growth. The p300 protein can regulate the transcription of the myogenic regulatory factor MyoD (Ji *et al.* 2003) and of the gene *Cugbp1* (Huichalaf *et al.* 2007). The CUGBP1 protein (CUG triplet repeat, RNA binding protein 1) is essential for the muscle differentiation process and any disturbance in the *Cugbp* gene expression leads to a delay in the myogenesis process or to a muscle dystrophy (Bhagwati *et al.* 1996).

Moreover, the interaction between p300 and GH was already evidenced for some genes, as the *RhoA* gene (*Ras* homolog gene family member A) and the early response *c-fos* gene, whose proteins regulate the actin cytoskeleton in the formation of fibers and the cell proliferation and differentiation, respectively (Ling & Lobie 2004, Cui *et al.* 2005). It seems that the growth hormone acts in the regulation of these genes, since GH can stimulate the p300 protein to occupy the promoter of these genes leading to their activation (Cui *et al.* 2005).

The present results indicate that the p300 gene, isolated from the brain of zebrafish, may have different expression levels in non-transgenic and GHtransgenic animals - HE individuals seem to present a lower expression, while an intermediary and a higher expression levels could be associated to HO and NT animals, respectively. These data reinforce the p300 role in muscle growth and its interaction with the growth hormone.

DDRT-PCR using primer HBV3

The use of the primer HBV3 to amplify the cDNA samples resulted on several faint and diffuse bands (Figure 1b). However, a fragment of approximately 900 bp was clearly visible in the NT, and in the GH-transgenic HE and HO samples of *Danio rerio* (Figure 1b). Although the difference in the stain intensity of this fragment among the three analyzed samples was not so evident, it was possible to notice a darker staining in the NT samples, which led us to infer that this fragment could also correspond to a differentially expressed transcript. The densitometric analyses of this transcript suggested a higher expression level in NT animals (22.73±3.80) than in HE (14.05±4.95) and HO samples (10.63±0.67) (Figure 2b). In the HE and HO samples, the expression level of the analyzed transcript represented approximately 62% and 47% of the IOD value obtained for the NT animals, respectively. Therefore, the IOD value decrease observed in HE and HO individuals can be directly related to the increased exogenous GH level in these two samples of zebrafish.

Nucleotide sequencing analysis of the transcript obtained with the primer HBV3 evidenced a DNA fragment with 831 bp (Figure 4). BLAST

143

database searches for nucleotide similarity indicated a high identity level (94%) with the 3' end of the gene that codes for the protein adenylate cyclase 2 (ADCY2) found in brain tissue of *Danio rerio* (accession numbers CR759914 and GI6270118). The Adenylate cyclase (AC) catalyses the conversion of ATP into cAMP that then acts to regulate a wide variety of cellular processes. The AC activity can be regulated by hormones, neurotransmitters interactions with their receptors, and calmodulin (Aikawa *et al.* 2000, Niewiadomski *et al.* 2002, Zawilska *et al.* 2003).

Recently, it was evidenced that the neuropeptide PACAP (pituitary adenylate cyclase-activating polypeptide) can also modulate the AC activity, leading not only to the formation of cAMP, but also playing a role in mediating growth hormone release in the pituitary gland of vertebrates (Matsuda et al. 2008, Mitchell et al. 2008), including fish species, as grass carp (Wong et al. 2005), common carp (Xiao et al. 2002), sockey salmon (Parker et al. 1997), and European eel (Montero et al. 1998). Wong et al. (2005) verified that increased concentrations of PACAP and/or higher concentrations of cAMP were effective in inducing GH release. Therefore, the performed densitometric data suggest that the lower adenylate cyclase expression level in HO zebrafish could result in a decrease in GH release thus leading to the reduced growth performance of these animals. Since Figueiredo et al. (2007b) evidenced that these HO animals, despite presenting a higher level of exogenous GH, were expressing approximately only one third of the endogenous GH observed in non-transgenic animals, the p300 should be involved in the endogenous growth hormone release control.

Final considerations

The growth hormone has physiological actions in several organs, since growth hormone receptors (GHR) have been identified in different tissues (Hughes & Friesen 1985, Kelly *et al.* 1991). As the GH is a multiple-function hormone, its excess can cause several physiological and behavioral collateral effects in the organism, as decrease in fertility (Cecim *et al.* 1995a, b), increase in oxygen consumption (Cook *et al.* 2000, Mackenzie *et al.* 2003), premature aging (Wolf *et al.* 1993, Bartke *et al.* 1998), decrease in learning and memory (Meliska *et al.* 1997), and oxidative stress (Rollo *et al.* 1996, Brown-Borg *et al.* 2001, Rosa *et al.* 2008). Therefore, the presence of a higher level of exogenous GH in transgenic fish could also leads to metabolic alterations that could influence not only the animal growth but also several other traits.

Previous analyses evidenced that HE GH-transgenic zebrafish reached a final average mass higher than HO and NT animals and that, surprisingly, HO fish presented a growth performance similar to NT animals (Figueiredo *et al.* 2007b). Taking into account the amount of endogenous and exogenous growth hormone present in HO and HE animals, both transgenic samples produce higher levels of biological active GH than NT individuals. Therefore, the hormone excess could be inducing a negative feedback system over the endogenous GH expression (Figueiredo *et al.* 2007b). These data could be related to regulation systems of intracellular GH that involves IGF-I (insulinlike growth factor I), GHR (GH receptor), and SOCS proteins (suppressor of cytokine signaling) (Figueiredo *et al.* 2007b, Studzinski 2008). The present data also indicate that the p300 and ADCY2 are also involved in a regulation system for GH in zebrafish in the case of high circulating hormone levels. Therefore, it seem that the reduced growth performance observed not only in homozygote GH-transgenic *Danio rerio* but also in other vertebrates with an exogenous GH gene is a result of the biological action of several proteins and their interaction. Further research is needed to characterize the significance of these proteins in controlling vertebrate muscle growth.

Acknowledgements

This work was supported by grants from FAPESP (Fundação da Amparo à Pesquisa do Estado de São Paulo), CNPq (Conselho Nacional de Desenvolvimento Científico e Tecnológico), and CAPES (Coordenação de Aperfeiçoamento de Pessoal de Nível Superior).

References

- Aikawa N, Karasawa A, Ohmori K (2000). Effect of zaldaride maleate, an antidiarrheal compound, on intracellular cyclic nucleotide-mediated intestinal ion secretion in rats. *European Journal of Pharmacology* **394**: 233-236.
- Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ (1990) Basic local alignment search tool. *Journal of Molecular Biology* **215**: 403-410.
- Bartke A, Brown-Borg HM, Bode AM, Carlson J, Hunter WS, Bronson RT (1998) Does growth hormone prevent or accelerate aging? *Experimental Gerontology* 33: 675-687.
- Bass SLS & Gerlai R (2008) Zebrafish (*Danio rerio*) responds differentially to stimulus fish: The effects of sympatric and allopatric predators and harmless fish. *Behavioural Brain Research* **186**: 107-117.
- Biga PR & Goetz FW (2006) Zebrafish and giant danio as models for muscle growth: determinate vs. indeterminate growth as deermined by morphometric analysis. *American Journal of Physiology* **291**: 1327-1337.
- Bhagwati S, Ghatpande A, Leung B (1996) Identification of two nuclear proteins which bind to RNA CUG repeats: significance for myotonic dystrophy. *Biochemistry and Biophysics Research Community* **228**: 55-62.
- Brown-Borg H, Johnson WT, Rakoczy S, Romanick M (2001) Mitocondrial oxidant generation and oxidative damage in ames dwarf and GH transgenic mice. *Journal of American Aging Association* **24**: 85-100.
- Cecim M, Kerr J, Bartke A (1995a) Effects of bovine growth hormone (bGH) transgene expression or bGH treatment on reproductive functions in female mice. *Biology of Reproduction* **52:** 1144-1148.
- Cecim M, Fadden C, Kerr J, Steger RW, Bartke A (1995b) Infertility in transgenic mice overexpressing the bovine growth hormone gene, Disruption of the neuroendocrine control of prolactin secretion during pregnancy. *Biology of Reproduction* **52**: 1187-1192.

- Cook JT, Mcniven MA, Sutterlin AM (2000) Metabolic rate of pre-smolt growth-enhanced transgenic Atlantic salmon (*Salmo salar*). *Aquaculture* **188**: 33-45.
- Cui TX, Piwien-Pilipuk G, Huo JS, Kaplani J, Kwok R, Schwartz J (2008). Endogenous CCAAT/Enhancer binding protein β and p300 are both regulated by growth hormone to mediated transcriptional activation. *Molecular Endocrinology* **19**(8): 2175-2186.
- Dahm R, Schonthaler HB, Soehn AS, Marle J, Vrensen GFJM (2007) Development and adult of the eye lens in the zebrafish. *Experimental Eye Research* **85**: 74-89.
- Dakis D & Kouretas D (2002) Differential display of m-RNAs in fish gonads by modified DD-PCR. *In Vivo* **16**: 103-105.
- Detrich HW, Westerfield M, Zon LI (1998) *The zebrafish: Biology.* Academic Press, New York.
- Devlin RH, Yesaki TY, Biagi CA, Donaldson EM, Swanson P, Chan WK (1994) Extraordinary salmon growth. *Nature* **371**: 209-210.
- Devlin RH, Biagi CA, Yesaki TY (2004) Growth, viability and genetics characteristics of GH transgenic coho salmon strains. *Aquaculture* **236**: 607-632.
- Devlin RH, Sundstro LF, Muir WM (2006) Interface of biotechnology and ecology for environmental risk assessments of transgenic fish. *Trends in Biotechnology* **24**: 89-97.
- Fadool JM & Dowling JE (2008) Zebafish: a model system for the study of eye genetics. *Progress in Retinal and Eye Research* **27**: 89-110.
- Figueiredo MA, Lanes CFC, Almeida DV, Marins LF (2007a) Improving to production of transgenic fish germlines: In vivo evaluation of mosaicism in zebrafish (*Danio rerio*) using a green fluorescent protein (GFP) and growth hormone cDNA transgene co-injection strategy. *Genetics and Molecular Biology* **30**: 31-36.

- Figueiredo MA, Lanes CFC, Almeida DV, Proietti MC, Marins LF (2007b) The effect of GH overexpression on GHR and IGF-I gene regulation in different genotypes of GH-transgenic zebrafish. *Comparative Biochemistry Physiology*, *Part D* **2**: 228-233.
- Fletcher GL & Davies PL (1991) Transgenic fish for aquaculture. *Genetic Engineering* **13**: 331-371.
- Florini JR, Ewton DZ, Coolican SA (1996) Growth hormone and the insulinlike growth factor system in myogenesis. *Endocrine Reviews* **17**: 481-517.
- Francis YI, Diss JKJ, Kariti M, Stephanou A, Latchman DS (2007) p300 activation by presenilin 1 but not by its M146L mutant. *Neuroscience Letters* **413**: 137-140.
- Harris AS & Wrigth JM (1995) Nucleotide sequence and genomic organization of cichlid fish minisatellites. *Genome* **38**: 177-184.
- Hew CL, Fletcher GL, Davies PL (1995) Transgenic salmon: tailoring the genome for food production. *Journal of Fish Biology* **47**:1-19.
- Higashijima S, Okamoto H, Ueno N, Hotta Y, Eguchi G (1997) High-frequency generation of transgenic zebrafish which reliably express GFP in whole muscles or the whole body by using promoters of zebrafish origin. *Developmental Biology* **192**: 289-299.
- Hughes JP & Friesen (1985) The nature and regulation of the receptors for pituitary growth hormone. *Annual Reviews of Physiology* **47**: 469-482.
- Huichalaf CH, Sakai K, Wang G-L, Timchenko NA, Timchenko L (2007) Regulation of the promoter of CUG triplet repeat binding protein, Cugbp 1, during myogenesis. *Gene* **396**:391-402.
- Huret JL (2000) EP300 (E1A binding protein p300). *The Atlas of Genetics and Cytogenetics in Oncology and Haematology*. URL http://AtlasGeneticsOncology.org.
- Jeay S, Sonenshein GE, Postel-Vinay MC, Kelly PA, Baixeras E (2002) Growth hormone can act as a cytokine controlling survival and proliferation of immune cells: new insights into signalling pathways. *Molecular and Cellular Endocrinology* **188**: 1-7.

- Jeffreys AJ, Wilson V, Thein SL (1985) Hypervariable minisatellite regions in human DNA. *Nature* **314**: 67-73.
- Ji AM, Dao D, Chen JX, MacLellan WR (2003) EID-2, a novel member of the EID family of p300-binding proteins inhibits transactivation by MyoD. *Gene* **318**: 35-43.
- Kelly PA, Djiane J, Postel-Vinary MC, Edery M (1991) The prolactin/growth hormone receptor family. *Endocrine Reviews* **12**: 235-251.
- Kimmel CB (1989) Genetics and early development of zebrafish. *Trends in Genetics* **5**: 283-288.
- Leroith D, Bondy C, Yakar S, Liu JL, Butler A (2001) The somatomedin hypothesis: 2001. *Endocrine Reviews* **22**: 53-74.
- Ling L & Lobie PE (2004) RhoA/ROCK activation by growth hormone abrogates p300/histone deacetylase 6 repression of State5-mediated transcripton. *Journal of Biology Chemestry* **279**: 32737-32750.
- Martínez R, Arenal A, Estrada MP, Herrera F, Huerta V, Vásquez J, Sánchez T, Fuente J (1999) Mendelian transmission, transgene dosage and growth phenotype in transgenic tilapia (*Oreochromis hornorum*) showing ectopic expression of homologous hormone. *Aquaculture* **173**: 271-283.
- Matsuda K, Nejigaki Y, Satoh M, Shimaura C, Tanaka M, Kawamoto K, Uchiyama M, Kawauchi H, Shioda S, Takahashi A (2008) Effect of pituitary adenylate cyclase-activating polypeptide (PACAP) on prolactin and somatolactin release from the goldfish pituitary in vitro. *Regulatory Peptides* **145**: 72-79.
- Mckenzie DJ, Martinez R, Morales A, Acosta J, Morales R, Taylor EW, Steffensen JF, Estrada MP (2003) Effects of growth hormone transgenesis on metabolic rate, exercise perfomance and hypoxia tolerance in tilapia hybrids. *Journal of Fish Biology* **63**: 398-409.
- Meliska CJ, Burke PA, Bartke A, Jensen RA (1997) Inhibitory avoidance and appetitive learning in aged normal mice, Comparison with transgenic mice having elevated plasma growth hormone levels. *Neurobiology of Learning and Memory* **68**: 1-12.

- Mitchell G, Sawisky GR, Grey CL, Wong CJ, Uretsky AD, Chang JP (2008) Differential involvement of nitric oxide signaling in dopamine and PACAP stimulation of growth hormone release in goldfish. *General and Comparative Endocrinology* **155**: 318-327.
- Moller N & Norrelund H (2003) The role of growth hormone in the regulation of protein metabolism with particular reference to conditions of fasting. *Hormone Research* **59**: 62-68.
- Montero M, Yon L, Rousseau K, Arimura A, Fournier A, Dufour S, Vaudry H (1998) Distribution, characterization, and growth hormone-releasing activity of pituitary adenylate cyclase activing polypeptide in the European eel (*Anguilla anguilla*). *Endocrinology* **139**: 4300-4310.
- Morales R, Herrera M, Arenal A, Cruz A, Hernandez O, Pimentel R, Guillen I, Martinez R, Estrada M (2001) Tilapia chromosomal growth hormone gene expression accelerates growth in transgenic zebrafish (*Danio rerio*). *Electronic Journal of Biotechnology* **4**: 51-58.
- Murray MJ, Haldeman BA, Grant FJ, O'Hara N (1988) Probing the human genome with minisatellite-like sequences from the human coagulation factor-VII gene. *Nucleic Acids Research* **16**: 4166.
- Nakamura Y, Leppert M, O'Connell P, Wolff R, Holm T, Culver M, Martin C, Fujimoto E, Hoff M, Kumlin E, White R (1987) Variable number of tandem repeat (VNTR) markers for human-gene mapping. *Science* 235: 1616-1622.
- Nam YK, Noh JK, Cho YS, Cho HJ, Cho KN, Kim CG, Kim DS (2001) Dramatically accelerate growth and extraordinary gigantism of transgenic mud loach *Misgurnus mizolepis. Transgenic Research* **10**: 353-362.
- Niewiadomski P, Nowak JZ, Sedkowska P, Zawilska JB (2002) Rapid desensitization of receptors for pituitary adenylate cyclase-activating polypetide (PACAP) in chick cerebral cortex. *Polish Journal of Phamacology* 54: 717-721.

- Parker DB, Power ME, Swanson P, Rivier J, Sherwood NM (1997) Exon skipping in the gene encoding pituitary adenylate cyclase-activating polypeptide in salmon alters the expression of two hormones that stimulate growth hormone release. *Endocrinology* **138**: 414-423.
- Rahman MA, Mak R, Ayad H, Smith A, Maclean N (1998) Expression of a novel growth hormone gene results in growth enhancement in transgenic tilapia (*Oreochromis niloticus*). *Transgenic Research* **7**: 357-369.
- Renoult L, Prévost M, Brodeur M, Lionnet C, Joober R, Malla A, Debruille JB (2007). P300 asymmetry and positive symptom severity: A study in the early stage of a first episode of psychosis. *Schizophrenia Research* **93**: 366-373.
- Rollo CD, Carlson J, Sawada M (1996) Accelerated aging of giant transgenic mice is associated with elevated free radical processes. *Canadian Journal of Zoology* **74**: 606-620.
- Rosa CE, Figueiredo MA, Lanes CFC, Almeida DV, Monserrat JM, Marins, LF (2008) Metabolic rate and reactive oxygen species productions in different genotypes of GH-transgenic zebrafish. *Comparative Biochemistry and Physiology, Part B* **149**: 209-214.
- Sambrook J & Russel DW (2001) *Molecular cloning. A Laboratory Manual*, 3rd Edition. Cold Spring Harbor Laboratory Press, New York.
- Simpson P, Peterson B, Hughes N, and Schelling GT (2000) Recombinant bovine somatotropin enhances growth rates in two species of ornamental fish; giant danios (*Danio aequipinnatus*) and zebrafish (*Danio rerio*). *Journal of Animal Sciences* **78**: 139.
- Sprague J, Clements D, Conlin T, Edwards P, Frazer K, Schaper K, Segerdell E, Song P, Sprunger B, Westerfield M (2003) The zebrafish information network (ZFIN): the zebrafish model organism database. *Nucleic Acids Research* **31**: 241-243.
- Studzinski ALM (2008) Regulação negativa do eixo somatotrófico em fígado de peixes transgênicos homozigóticos superexpressando o gene do hormônio do crescimento (GH). Master Thesis. Fundação Universidade Federal do Rio Grande, Rio Grande, RS.

- Thompson JD, Higgins DG, Gibson TJ (1994) Clustal W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Research* **22**: 4673-4680.
- Tu AW & Luo K (2007) Acetylation of Smad2 by the co-activator p300 regulates activin and TGFβ response. *Journal of Biological Chemistry* **282**(29): 21187-21196.
- Vachtenheim J, Sestáková B., Tuhácková Z (2006) Inhibition of MITF transcriptional activity independent of targeting p300/CBP coactivators. *Pigment Cell Research* **20**: 41-51.
- Vleugel MM, Shvarts D, Wall E, Diest PJ (2006) p300 and p53 levels determine activation of HIF-1 downstream targets in invasive breast cancer. *Human Pathology* **37**: 1085-1092.
- Xiao D, Chu MMS, Lee EKY, Lin H-R, Wong AOL (2002) Regulation of growth hormone release in common carp pituitary cells by pituitary adenylate cyclase-activating polypeptide: signal transduction involves cAMP and calcium-dependent mechanisms. *Neuro Endocrinology* **76**: 325-338.
- Yoshizato H, Fujikawa T, Soya H, Tanaka M, Nakashima K (1998) The growth hormone (GH) gene is expressed in the lateral hypothalamus: enhancement by GH-releasing hormone and repression by restraint stress. *Endocrinology* **139**: 2545-2551.
- Westerfield M (1995) *The Zebrafish Book. A guide for the laboratory use of zebrafish (Brachydanio rerio)*. University of Oregon Press. Eugene, Oregon.
- Wolf E, Kahnt E, Ehrlein J, Hermans W, Brem G, Wankle R (1993) Effects of long-term elevated serum levels of growth hormone on life expectancy of mice: lessons from transgenic animal models. *Mechanism of Ageing and Development* 68: 71-87.
- Wong AO, Li W, Leung CY, Huo L, Zhou H (2005) Pituitary adenylate cyclaseactivating polypeptide (PACAP) as a growth hormone (GH)-releasing factor in grass carp. I. Functional coupling of cyclic adenosine 3',5'monophosphate and Ca2+/calmodulin-dependent signaling pathways in PACAP-induced GH secretion and GH gene expression in grass carp pituitary cells. *Endocrinology* **146**: 5407-5424.

- Zawilska JB, Niewiadomski P, Nowak JZ (2003) Pituitary adenylate cyclaseactivating polypetide (PACAP)-Evoked increase in cyclic AMP production in chick cerebral cortex: lack of a role of the protein kinase C pathway. *Polish Journal of Phamacology* **55**: 245-250.
- Zbikowska HM (2003) Fish can be first Advances in fish transgenesis for commercial applications. *Transgenic Research* **12**: 379-389.
- Zhao L-J, Subramanian T, Chinnadurai G (2006) Changes in C-terminal binding protein 2 (CtBP2) corepressor complex induced by E1A and modulation of E1A transcriptional activity by CtBP2. *Journal of Biological Chemistry* **281**: 36613-36623.

Table, Figures and Legends

Table 1:	Oligonucleotides that were designed based on VNTR core sequences
and used	as single primers for the amplification of the cDNA samples of Danio
rerio.	

Primer	Sequence (5' - 3')	Reference
INS	ACAGGGGTGTGGGG	Nakamura <i>et al.</i> (1987)
YNZ22	CTCTGGGTGTCGTGC	Nakamura <i>et al.</i> (1987)
HBV5	GGTGTAGAGAGGGGT	Nakamura <i>et al.</i> (1987)
HBV3	GGTGAAGCACAGGTG	Nakamura <i>et al.</i> (1987)
Fvllex8	ATGCACACACAGG	Murray <i>et al.</i> (1988)
EMBL	AGAGCTTCAGGCTGGGCAGCTAAG	Harris & Wright (1995)



Figure 1: DDRT-PCR patterns of non-transgenic (NT) and of GH-transgenic hemizygote (HE) and homozygote (HO) zebrafish of the F0104 line, visualized trough 6% polyacrilamide gel electrophoresis. (a) brain cDNA amplification using primer INS; (b) brain cDNA amplification using primer HBV3. L, 1 Kb DNA molecular marker. Arrows indicate putative differentially expressed transcripts among the three analyzed samples.



Figure 2: Densitometric analyses of the fragments that correspond to putative differentially expressed transcripts identified through DDRT-PCR in non-transgenic (NT) and in GH-transgenic hemizygote (HE) and homozygote (HO) zebrafish of the F0104 line. (a) estimated expression level of the fragment amplified with the primer INS that corresponds to the gene that codes for protein p300 (E1A binding protein p300); (b) estimated expression level of the fragment amplified with the primer HBV3 that corresponds to the gene that codes for protein ADCY2 (adenylate cyclase 2).

Figure 3: Consensus nucleotide sequence of a partial region of the gene that codes for protein p300, isolated from brain tissue of *Danio rerio* through a DDRT-PCR approach using primer INS. The underlined regions are homologous to the primer sequence.

5' <u>ggtgaagcacangtg</u> caaacagtattatgtgggcaatcctaaaagaga48
tgtcaaaaacaaagaggtcggtccaggtggctagcaccagaaaacaataa98
acacanatgatctaacatggcagagtaaggcaaggagaaatgctttgtaa148
aggctgatttatactcctgcgtcaaacactggcgtatgctacggcgctga198
$\verb cgcatagcccttcgccctggccgtcgccgtcactgacgtgcaccactcta248 $
aaaattggaactacacatgggaacgacgcgtatcgcaaagtctgtgattg298
gtcggcttggtagcgttgacgagtcggggcaggaccgagacccgtgcgaa348
tggtgcaaacccaatgtagcgattgtttacaagtgtggagtcccgtgaag398
$\verb gagcttcagatggaaagttttgttttgtgtttacctcatagttaaagttg448 $
$\verb+ctgcatgtccgccggttgctgcctcaaaatgagcgagtttcagccacttg498$
tacattccggaagtgttcaggaaaagcaaaaagcagcgaagaaactcaa548
$\verb cacaaaggaacatttacacctcactgccaactagcatttcggaagtgtta598 $
atccagaccaacagagacagcgcgcagaagtataaatgcacagccacgtg648
$\verb cgttgcatgcaccgtgggttacgccggtcacttgacgcagaagtattaat698 $
$\verb caggcttaatgtgtccaggtaacaacaagactcagtaattgcatgtatgt$
atgtgtctgtgagtgtgtgtgtggggcgttgcatgaatagtccatgtaa798
tgagtgcttaaccagctt <u>cacctgtgcttcacc</u> 3'831

Figure 4: Consensus nucleotide sequence of a partial region of the gene that codes for protein ADCY2, isolated from brain tissue of *Danio rerio* through a DDRT-PCR approach using primer HBV3. The underlined regions are homologous to the primer sequence.

Considerações Finais e Conclusões

O objetivo principal do presente trabalho, especificado inicialmente, de isolar e caracterizar transcritos diferencialmente expressos em duas espécies de peixes - *Leporinus macrocephalus* e *Danio rerio* - foi alcançado com sucesso, dada a obtenção de resultados referentes às diferentes sub-propostas de pesquisa.

Assim, as análises desenvolvidas no presente estudo evidenciaram:

- Duas isoformas de α-actina esquelética na espécie Leporinus macrocephalus, geneticamente e provavelmente funcionalmente distintas, cujas características podem estar relacionadas à demanda energética dos tecidos musculares esqueléticos branco e vermelho.
- Um gene com expressão diferencial em tecido cerebral de machos e fêmeas de *Leporinus macrocephalus*, relacionado à proteína APBA2 (*amyloid beta (A4) precursor protein-binding, family A, member 2*), cuja expressão está associada à produção da proteína β-amilóide a qual, por sua vez, relaciona-se à doença de Alzheimer.
- A estrutura do gene *fabp6* (*fatty acid-binding protein 6*) de *Danio rerio*, similar a grande parte dos demais membros da família multigênica *FABP* de outros vertebrados, e sua distribuição tecidual em embriões, larvas e adultos, também similar à evidenciada em outros organismos, especialmente mamíferos.

Dois genes candidatos diferencialmente expressos, relacionados à proteína p300 (*E1A binding protein p300*) e à proteína ADCY2 (*adenylate cyclase 2*) envolvidas na formação/crescimento muscular e na liberação de GH, respectivamente, entre exemplares de *Danio rerio* não-transgênicos e transgênicos homozigotos e hemizigotos para gene do hormônio de crescimento.

Dada a importância da obtenção e ampliação de dados genéticos em peixes, que constituem um grupo extremamente diversificado e basal entre os vertebrados, os resultados obtidos contribuem para a melhor compreensão da evolução, função e organização molecular de diversos genes e podem servir de base a futuros estudos. Além disso, os dados obtidos confirmam a utilidade de Danio rerio como espécie modelo em estudos genéticos e indicam que Leporinus macrocephalus representa um potencial modelo experimental em análises direcionadas а genes envolvidos no desenvolvimento de doenças humanas, como alterações neurodegenerativas, distrofias musculares e câncer.

SÚMULA CURRICULAR

(Outras atividades vinculadas e desenvolvidas no Doutorado)

1. Trabalhos apresentados em congressos:

- <u>ALVES, F.A.;</u> FORESTI, F., OLIVEIRA, C., MARTINS, C. 2005. Uso de seqüências de DNAr 5S como marcador molecular em tilápia do Nilo (*Oreochromis niloticus*). In: "XVI Encontro Brasileiro de Ictiologia", João Pessoa, Paraíba, Brasil.
- <u>ALVES, F.A.</u>; MARTINS, C., WASKO, A.P. 2005. Caracterização molecular de genes de actina isolados de músculos branco e vermelho de *Leporinus macrocephalus* (Pisces, Anostomidae). In: *"51°. Congresso Brasileiro de Genética",* Águas de Lindóia/ SP, Brasil. Genetics and Molecular Biology (Suppl.) 26, GA123, p.123.
- MATOS, F.D.C.; <u>ALVES, F.A.;</u> MARTINS, C.; ALVES, A.L.; FORESTI,
 F.; OLIVEIRA, C.; WASKO, A.P. 2005. Caracterização do DNAr 5S
 em espécies da família Sciaenidae (Pisces, Perciformes). In: "51°. *Congresso Brasileiro de Genética*", Águas de Lindóia/ SP, Brasil.
 Genetics and Molecular Biology (Suppl.) 26, GA199, p.199.
- WONG, M.S.L.; <u>ALVES, F.A.</u>; WASKO, A.P. 2005. Identificação de um marcador RAPD associado a fêmeas de *Brycon cephalus* (Pisces, Bryconidae). In: "51° Congresso Brasileiro de Genética", Águas de Lindóia/ SP, Brasil. Genetics and Molecular Biology (Suppl.) 26, GA229, p.229.
- <u>ALVES-COSTA, F.A.;</u> MARTINS, C.; WASKO, A.P. 2006. Identificação de transcritos sexo-específicos em *Leporinus macrocephalus* (Pisces, Anostomidae) através da metodologia de DDRT-PCR. In:
 "XXVI Congresso Brasileiro de Zoologia", Universidade Estadual de Londrina UEL, Londrina/ PR, Brasil. Nº. 858_1791.

- WONG, M.S.L.; <u>ALVES-COSTA, F.A.</u>; WASKO, A.P. 2006. Prospecção de fragmentos sexo-específicos em *Brycon cephalus* (Pisces, Bryconidae) através da metodologia de RAPD. In: "XXVI Congresso Brasileiro de Zoologia", Universidade Estadual de Londrina UEL, Londrina/ PR, Brasil. Nº. 2123_1791.
- <u>ALVES-COSTA, F.A.;</u> MARTINS, C.; WASKO, A.P. 2006. Identificação e caracterização de transcritos com potencial expressão diferencial entre os sexos de *Leporinus macrocephalus* (Pisces, Anostomidae). **In**: "52° Congresso Brasileiro de Genética" and "12°. Congresso de la Associación Latinoamericana de Genética", Bourbon Cataratas Resort & Convention Center, Foz do Iguaçu/PR, Brasil. p. 187.
- WONG, M.S.L.; <u>ALVES-COSTA, F.A.</u>; WASKO, A.P. 2006. Identificação e caracterização de marcadores RAPD associados ao sexo em *Brycon cephalus* (Pisces, Bryconidae). In: "52° Congresso Brasileiro de Genética" and "12° Congresso de la Associación Latinoamericana de Genética", Bourbon Cataratas Resort & Convention Center, Foz do Iguaçu/PR, Brasil. p. 235.
- GOMES, M.L.; <u>ALVES-COSTA, F.A.</u>; NISHIDA, S.M.; WASKO, A.P.
 2006. Análise de locos microssatélites em codorna japonesa *Coturnix japonica* (Galliformes, Phasianidae) para identificação de polimorfismos, visando a elaboração de testes de paternidade. In: *"52° Congresso Brasileiro de Genética" and "12°. Congresso de la Associación Latinoamericana de Genética"*, Bourbon Cataratas Resort & Convention Center, Foz do Iguaçu/PR, Brasil. p. 264.

<u>ALVES-COSTA, F.A.;</u> MARTINS, C.; WASKO, A.P. 2006. Prospecção de DETS (Differentially Expressed Transcripts) em machos e fêmeas de *Leporinus macrocephalus* (Anostomidae). **In**: *"XI Brazilian Symposium on Fish Cytogenetics and Genetics and I International Congress of Fish Genetics"*, Universidade Federal de São Carlos, São Carlos/SP, Brasil. p. 277. WONG, M.S.L.; <u>ALVES-COSTA, F.A.;</u> WASKO, A.P. 2006. Identificação e caracterização de fragmentos de DNA sexoespecíficos em *Brycon cephalus* (Characidae) através da metodologia de RAPD. In: "XI Brazilian Symposium on Fish Cytogenetics and Genetics and I International Congress of Fish Genetics", Universidade Federal de São Carlos, São Carlos/SP, Brasil. p. 210.

- <u>ALVES-COSTA, F.A.;</u> MARTINS, C.; WASKO, A.P. 2006. Uso de seqüências de DNAr 5S como marcadores moleculares em espécies marinhas da família Sciaenidae (Pisces, Perciformes). **In**: "*VI Jornadas de Ciencias del Mar and XIV Coloquio de Oceanografía",* Puerto Madryn, Chubut, Argentina, p. 158.
- TONIATO, J.; MARTINS, C.; <u>ALVES-COSTA, F.A.</u>; ROCHA, K.K.H.R. 2007. Diferenciação molecular da tilápia do Nilo com base em marcadores moleculares de baixo custo. **In**: "XXII Reunião Anual da FESBE", Águas de Lindóia/SP, Brasil. Métodos, 49.011 -Equipamentos e Ensino.
- <u>ALVES-COSTA, F.A.;</u> FIGUEIREDO, M.A.; LANES, C.F.C.; ALMEIDA, D.V.;
 MARINS, L.F.; WASKO, A.P. 2007. Isolamento e caracterização de DETs ("Differentially Expressed Transcripts") em tecido cerebral de uma linhagem de de peixe transgênico (*Danio rerio*) superexpressando o gene do hormônio do crescimento (GH). In: "53°. Congresso Brasileiro de Genética", Hotel Monte Real Resort, Águas de Lindóia/SP, Brasil. p. 168.
- GOMES, M.L.; <u>ALVES-COSTA, F.A.;</u> NISHIDA, S.M.; WASKO, A.P.
 2007. Elaboração de testes de paternidade em codorna japonesa *Coturnix japonica* (Galliformes, Phasianidae). In: "53°. Congresso Brasileiro de Genética", Hotel Monte Real Resort, Águas de Lindóia/SP, Brasil. p.257.

- TONIATO, J.; MARTINS, C.; <u>ALVES-COSTA, F.A.</u> 2007. Aplicação de marcadores RFLP-PCR na identificação de espécies de tilápias (Perciformes: Cichlidae). In: "53°. Congresso Brasileiro de Genética", Hotel Monte Real Resort, Águas de Lindóia/SP, Brasil. p.91
- GOMES, M.L.; <u>ALVES-COSTA, F.A.</u>; NISHIDA, S.M.; WASKO, A.P.
 2007 Testes de paternidade para a codorna japonesa *Coturnix japonica* (Galliformes, Phasianidae): ferramenta para evidenciar a competição pós-cópula. In: *"XXV Encontro Anual de Etologia",* UNESP/São José do Rio Preto, São José do Rio Preto, SP, Brasil. Oral presentation.

2. Participações em congressos:

- XVI Encontro Brasileiro de Ictiologia, realizado em João Pessoa. Paraíba, no período de 24 a 28 de janeiro de 2005.
- 51°. *Congresso Brasileiro de Genética* realizado em Águas de Lindóia, SP, no período de 7 a 10 de setembro de 2005.
- XXVI Congresso Brasileiro de Zoologia, realizado na Universidade Estadual de Londrina e Centro Universitário Filadélfia, Londrina, PR, de 12 a 17 de fevereiro de 2006.
- 52º. Congresso Brasileiro de Genética e 12º. Congresso de la Associación Latinoamericana de Genética, realizado no Bourbon Cataratas Resort & Convention Center, Foz do Iguaçu, PR, de 3 a 6 de setembro de 2006.

XI Brazilian Symposium on Fish Cytogenetics and Genetics and I International Congress of Fish Genetics, realizado na Universidade Federal de São Carlos, São Carlos, SP, de 10 a 13 de outubro de 2006. VI Jornadas de Ciencias del Mar y XIV Coloquio de Oceanografía, realizada na cidade de Puerto Madryn, Chubut, Argentina, de 4 a 8 de dezembro de 2006.

3. Cursos assistidos:

- *Evolução de proteínas através de embaralhamento de DNA*, realizado durante o 51°. *Congresso Brasileiro de Genética,* em Águas de Lindóia, SP, no período de 7 a 10 de setembro de 2005, totalizando 3 horas.
- *Expressão Gênica: novas formas de análise e abordagens*", realizado durante o 52°. Congresso Brasileiro de Genética, Bourbon Cataratas Resort & Convention Center, Foz do Iguaçu, PR, no período de 3 a 6 de setembro de 2006, totalizando 3 horas.
- *Melhoramento Genético em Peixes*", realizado durante o XI Brazilian Symposium on Fish Cytogenetics and Genetics and I International Congress of Fish Genetics, na Universidade Federal de São Carlos, São Carlos, SP, no período de 10 a 13 de outubro de 2006, totalizando 6 horas.

4. Cursos ministrados:

Curso de extensão Universitária "Introdução à Biologia Molecular", realizado no Laboratório de Genética de Peixes, junto ao Departamento de Ciências Biológicas, UNESP, Bauru, SP, durante o período de 16 a 19 de agosto de 2005, com duração de 24 horas.

- Curso "*Marcadores Citogenéticos e Moleculares: conceitos e aplicações*", ministrado durante o VI Workshop de Genética, no Instituto de Biociências, UNESP, Botucatu, SP, no período de 20 a 21 de maio de 2006, apresentando carga horária equivalente a 8 horas.
- Curso "Marcadores Moleculares e Sua Aplicação em Teste de Paternidade: uma abordagem sobre conservação genética", ministrado durante o VII Workshop de Genética, no Instituto de Biociências, UNESP, Botucatu, SP, no período de 20 a 21 de maio de 2007, apresentando carga horária equivalente a 8 horas.

5. Monitoria em cursos de extensão:

- Monitoria na disciplina "Métodos de Estudo em Biologia Molecular", sob a responsabilidade do Prof. Cesar Martins, no curso de pósgraduação em Ciências Biológicas (Genética), ministrada de 10 a 19 de outubro de 2005.
- Colaboração como monitora durante o curso de extensão universitária (I Curso de Inverno) "Manipulação de Ácidos Nucleicos: PCR, RT-PCR e PCR em tempo real", realizado no Instituto de Biociências, UNESP, Botucatu, SP, no período de 10 a 14 de julho de 2006
- Colaboração como monitora da oficina "Experimentando Genética", caracterizada como parte do Prpgrama de Extensão "Difundindo e Popularizando a Ciência", realizada no Instituto de Biociências da UNESP, Botucatu, SP, no período de 15 a 20 de janeiro de 2007, totalizando 48 horas de atividade.
- Colaboração como monitora durante curso de extensão universitária (II Curso de Inverno - "Manipulação de Ácidos Nucleicos: PCR, RT-PCR e PCR em tempo real", realizado no Instituto de Biociências, UNESP, Botucatu, SP, no período de 16 a 20 de julho de 2007.

- "Expressão de Proteínas Recombinantes", realizada durante curso de extensão universitária "Introdução à Biologia Molecular", Instituto de Biociências de Botucatu, Universidade Estadual Paulista - UNESP (Botucatu, SP), no dia 13 de julho de 2007.
- "Brazilian Conservation Genetics Issues", realizada na Acadia University, Wolfville, Nova Scotia, Canadá, no dia 10 de outubro de 2007.

7. Publicações:

- <u>Alves-Costa FA</u>, Wasko AP, Oliveira C, Foresti F, Martins C (2006) Genomic organization and evolution of the 5S ribosomal DNA in Tilapiini fishes. *Genetica* 127:243-252.
- <u>Alves-Costa FA,</u> Martins C (2006) DNAr 5S um novo marcador molecular para análise genética de tilápias. *Revista Biotecnologia, KL3 Publicações Ltda* 35: 22-27.
- Wasko AP, Bento AP, Ferreira DC, Souza DB, <u>Alves-Costa FA</u>, Rossetto FO, Gatinho IA, Maia IG, Mazzuchelli J, Paiva LRS, Ribolla PEM, Almeida RA, Barbetta SR, Salles VP, Teixeira WG, Martins C (2007) Aliando conceitos e criatividade: proposta de dramatização na área de genética e biologia celular para alunos do ensino médio. *Genética na Escola 4*: 34-38.
- <u>Alves-Costa FA</u>, Matos FDC, Foresti F, Oliveira C, Martins C, Wasko AP (2008) 5S rDNA characterization in twelve Sciaenidae fish species (Teleostei: Perciformes): depicting gene diversity and molecular markers. *Genetics and Molecular Biology* 31, 1 (suppl): 303-307.

<u>Alves-Costa FA</u>, Denovan-Wright EM, Thisse C, Thisse B, Wright JM (2008) Spatial-temporal distribution of fatty acid-binding protein 6 (*fabp6*) gene transcripts in the developing and adult zebrafish (*Danio rerio*). *FEBS Journal* 275: 3325-3334.

8. Estágios realizados:

- Estágio supervisionado pelo Prof. Dr Luis Fernando Marins, no Laboratório de Biologia Molecular do Departamento de Ciências Fisiológicas, FURG (Fundação Universidade do Rio Grande), realizado de 14 a 26 de outubro de 2006, com participação no desenvolvimento da tecnologia de DDRT-PCR (*Differential Display Transcriptase Reverse – Polymerase Chain Reaction*) dentro do projeto "Peixes Geneticamente Modificados para o Gene do Hormônio de Crescimento (GH)", sob a coordenação do mesmo professor.
- Estágio supervisionado pelo Prof. Dr. Jonathan M. Wright, Department of Biology, Dalhousie University, Halifax, Nova Scotia, Canada, no período de 30 de julho a 07 de dezembro de 2007, desenvolvendo o trabalho de pesquisa "Caracterização da organização e da expressão gênica da proteína FABP6 ("Fatty Acid-binding Protein 6") na espécie de peixe *Danio rerio* (zebrafish)".

9. Participação em projetos de pesquisa:

- "Isolamento e caracterização de transcritos sexo-específicos em peixes utilizando a técnica de DDRT-PCR ("Differential Display Reverse Transcriptase"): modelo inicial para implementação de linha de pesquisa em expressão gênica diferencial". Projeto de Jovem Pesquisador Supervisor(a): Dra. Adriane Pinto Wasko) – FAPESP nº. 03/06761-1. Laboratório de Biologia Molecular e do Desenvolvimento, Departamento de Morfologia, Instituto de Biociências, UNESP (Botucatu, SP). Período de Dezembro de 2003 a Novembro de 2006.
- "Identificação de marcadores sexo-específicos em Leporinus macrocephalus (Pisces, Anostomidae) utilizando a técnica de RAPD". Projeto de Iniciação Científica (Aluna: Marina Seik Lien Wong) – FAPESP nº. 04/10328-4,. Departamento de Morfologia, Instituto de Biociências, UNESP (Botucatu, SP). Dezembro de 2004 à Novembro de 2005.
- "Identificação de marcadores sexo-específicos em *Brycon cephalus* (Pisces, Bryconinae) utilizando a técnica de RAPD". Projeto de Iniciação Científica (Aluna: Fernanda Del Campos de Matos) – FAPESP nº. 04/10911-1, Departamento de Morfologia, Instituto de Biociências, UNESP (Botucatu, SP). Período de Dezembro de 2004 a Novembro de 2006.
- "Comparação e aperfeiçoamento de técnicas para isolamento de DNA de aves através de amostragem não-invasiva". Projeto de Iniciação Científica (Aluno: Tiago José Benedito Eugênio). Laboratório de Biologia Molecular e do Desenvolvimento, Departamento de Morfologia, Instituto de Biociências, UNESP (Botucatu, SP). Período de Abril de 2005 a Março de 2006.

- "Identificação de marcadores RAPD sexo-específicos e DNAs satélites em Brycon cephalus (Pisces, Bryconinae): contribuições a caracterização genética da espécie" Projeto de Iniciação Científica (Aluna: Marina Sek Lien Wong) –FAPESP nº. 04/10328-4, Departamento de Genética, Instituto de Biociências, UNESP (Botucatu, SP). Período de Dezembro de 2006 a Novembro de 2007.
- "Isolamento e caracterização de transcritos sexo-específicos em Leporinus macrocephalus (Pisces, Characiformes), utilizando a técnica de DDRT-PCR (Differential Display Reverse Transcriptase)" – Projeto de Doutorado (Aluna de pós-graduação: Fernanda Antunes Alves da Costa) FAPESP nº. 04/10024-5,. Laboratório de Biologia Molecular Animal, Departamento de Genética, Instituto de Biociências, UNESP (Botucatu, SP). Período de Fevereiro de 2005 a Agosto de 2008.
- "Determinação das relações entre o comportamento de acasalamento e o sucesso reprodutivo na codorna japonesa (*Coturnix japonica*): uma abordagem multidisciplinar através de análises etológicas e genéticas". Projeto de Mestrado (Aluna de pós-graduação Magali Lira Gomes) FAPESP nº.05/57996-4, Laboratório de Biologia Molecular Animal, Departamento de Genética, Instituto de Biociências, UNESP (Botucatu, SP). Período de Abril de 2005 a Agosto de 2008.
10. Disciplinas cursadas:

Disciplinas	Créditos	Conceito
Genética Molecular em Animais de Produção (mamíferos)	03	A
Fundamentos em Genética	06	A
Tópicos especiais em genética: expressão, purificação e caracterização de proteínas recombinantes	03	A
Desenvolvimento embrionário comparado: modelos de estudo	03	A
Sistemática Molecular	03	A
Análisos cromossâmicas molocularos: tácnicas do	03	٨
hibridação <i>in situ</i>	05	~
Organização e evolução dos genomas	04	А
Difundindo Genética: popularização da ciência pelos genes	04	A
Mecanismos de Regulação e Métodos de Análise da	03	А
Conservation Genetics Professores Responsáveis: Adriane Pinto Wasko	03	A
(Departamento de Genética, Instituto de Biociências,		
UNESP, Botucatu, SP) e Stephen Mockford (Acadia University, Wolfville, Nova Scotia, Canada)		

Livros Grátis

(<u>http://www.livrosgratis.com.br</u>)

Milhares de Livros para Download:

Baixar livros de Administração Baixar livros de Agronomia Baixar livros de Arquitetura Baixar livros de Artes Baixar livros de Astronomia Baixar livros de Biologia Geral Baixar livros de Ciência da Computação Baixar livros de Ciência da Informação Baixar livros de Ciência Política Baixar livros de Ciências da Saúde Baixar livros de Comunicação Baixar livros do Conselho Nacional de Educação - CNE Baixar livros de Defesa civil Baixar livros de Direito Baixar livros de Direitos humanos Baixar livros de Economia Baixar livros de Economia Doméstica Baixar livros de Educação Baixar livros de Educação - Trânsito Baixar livros de Educação Física Baixar livros de Engenharia Aeroespacial Baixar livros de Farmácia Baixar livros de Filosofia Baixar livros de Física Baixar livros de Geociências Baixar livros de Geografia Baixar livros de História Baixar livros de Línguas

Baixar livros de Literatura Baixar livros de Literatura de Cordel Baixar livros de Literatura Infantil Baixar livros de Matemática Baixar livros de Medicina Baixar livros de Medicina Veterinária Baixar livros de Meio Ambiente Baixar livros de Meteorologia Baixar Monografias e TCC Baixar livros Multidisciplinar Baixar livros de Música Baixar livros de Psicologia Baixar livros de Química Baixar livros de Saúde Coletiva Baixar livros de Servico Social Baixar livros de Sociologia Baixar livros de Teologia Baixar livros de Trabalho Baixar livros de Turismo