

---

Ariane Lima Xavier

Influência de fatores ambientais na distribuição  
espacial de uma comunidade de anuros em  
área de altitude no domínio das caatingas

**Orientador:** Dr. Marcelo Felgueiras Napoli

Dissertação apresentada ao Programa  
de Pós-Graduação em Ecologia e  
Biomonitoramento da Universidade  
Federal da Bahia como parte dos  
requisitos necessários à obtenção do  
título de mestre em Ecologia e  
Biomonitoramento.

Salvador  
2008

---

# **Livros Grátis**

<http://www.livrosgratis.com.br>

Milhares de livros grátis para download.

---

## *Ficha Catalográfica*

---

### **Fotografias**

Banco de imagens do Laboratório de Taxonomia e História Natural de Anfíbios da Universidade Federal da Bahia: fotografias de Rafael O. Abreu (capa, fig. 7a-7e, 7g-7j, 7m-7p) e de Deise Cruz (Fig. 7l).  
Prof<sup>a</sup> Dra. Flora Juncá (Fig. 7f)

Capa: *Bokermannohyla oxente*

Biblioteca Central Reitor Macêdo Costa - UFBA

\_\_\_\_ Xavier, Ariane Lima

“Influência de fatores ambientais na distribuição espacial de uma comunidade de anuros em área de altitude no domínio das caatingas” / Ariane Lima Xavier – 2008.

Inclui Bibliografia

Orientador: Marcelo Felgueiras Napoli

Dissertação (mestrado) – Universidade Federal da Bahia, Instituto de Biologia, 2008.

1. Estrutura de comunidade de anuros 2. Variáveis ambientais 3. Floresta Estacional  
4. Campo Rupestre. I. Napoli, Marcelo Felgueiras. II. Universidade Federal da Bahia. Instituto de Biologia. III. Título.

### **Comissão Julgadora**

---

Prof<sup>a</sup>. Dr<sup>a</sup>. Gilda Vasconcellos de Andrade

---

Prof<sup>a</sup>. Dr<sup>a</sup>. Flora Acuña Juncá

---

Prof. Dr. Marcelo Felgueiras Napoli (Orientador)

*Para aqueles que  
me fizeram acreditar  
em anjos da guarda.*

---

## *Agradecimentos*

---

Agradeço aos meus adoráveis pais, José e Alice, pelo amor e confiança que sempre dedicaram a mim. Aos meus amados, Alanza e Rodrigo, obrigada por acreditarem em mim. À Aline, companheira e amiga, pela boa vontade em me escutar desde o tão comentado delineamento amostral à longa elaboração final. Espero um dia poder lhe retribuir todo o apoio.

Ao meu orientador, Prof. Dr. Marcelo Felgueiras Napoli, por apresentar-me o mundo dos sapos e da pesquisa, pela cessão de conhecimentos, com muita paciência, amizade e “bom humor” característico... Com admiração, obrigada!

Ao Prof. Dr. Miguel da Costa Accioly pelo convite para participar da elaboração do Plano de Manejo do Parque Estadual das Sete Passagens (PESP), o que tornou possível a construção deste trabalho. Agradeço por sua atenção, sempre pronto ajudar em todos os momentos em que precisei e não foram poucos.

À Dona Iracy e Seu Assis, presentes no início da caminhada, agradeço pela confiança, abrigo e amizade. Por vocês guardo profundo respeito, carinho, admiração e saudades.

Aos amigos Rodrigo e Elaine, corações enormes que me abrigaram durante esses anos. Meus amigos, companheiros, “pais”... preocupados e presentes nos momentos em que sabiam, ou que nem imaginavam, o quanto eu estava precisando. A vocês não só agradeço como dedico essa dissertação, “co-orientadores”...rsrsrs.

A todos os colegas do mestrado, em especial aos amigos Pablo, sempre presente quando eu precisei, Tasso e Roberta por vocês tenho grande carinho e admiração! Ao HOTSCOPS, pelos momentos de alegria e amizade que construímos: Igor, Perimar,

Márcio, Elaine, Rodrigo, Pablo, Paty, Eder, Tasso, Roberta, Magno e Fernandinha. Comprovamos empiricamente que a construção do conhecimento vai beeeem além dos muros da faculdade, perto de Itapuã...rs.

Aos amigos do Amphibia: Deise Cruz, Laís Encarnação, Milena Camardelli, Patricia Fonseca, Rafael Abreu e Thais Dória. Obrigada por me receberem em suas vidas. Guardo amor e respeito por todos vocês. Para Rafa agradeço em dobro, por todo carinho e dedicação que demonstrou na reta final. E àqueles que me acompanharam e ajudaram muito em campo: equipe AMPHIBIA, Natália, Wilton, Tiago Jordão, Tiago Filadelfo, Camila, Tasso, Heverton e Marina. Vocês tornaram o trabalho possível.

À Prof. Dr<sup>a</sup>. Flora Acuña Juncá, Rafael Abreu e Deise Cruz por cederem as fotografias utilizadas neste trabalho.

Ao Sr. José Manoel Pereira (Zélis), gestor do PESP, pelo apoio durante a realização dos trabalhos e aos guardas florestais: Adão, Evanaldo, Erivaldo, Geovan, Junior, Marivaldo e Vilobaldo, e aos guias Gê e Nilson obrigada pela boa vontade e amizade que vocês nos dedicaram em todos os momentos.

À Fundação de Amparo à Pesquisa do Estado da Bahia – FAPESB por ter me concedido bolsa durante a realização do mestrado.

Ao Instituto Brasileiro do Meio Ambiente e dos Recursos Naturais Renováveis – IBAMA por ter cedido a licença de coleta.


À Secretaria Estadual do Meio Ambiente e dos Recursos Hídricos – SEMARH por fornecer os “shapes” utilizados neste trabalho.

# *Sumário*

---

<b>Introdução geral</b>	<b>1</b>
<b>Manuscrito.</b> Revista <i>Biodiversity and Conservation</i>	
Abstract	<b>7</b>
<b>1. Introduction</b>	<b>8</b>
<b>2. Material and methods</b>	
2.1. Study area	<b>10</b>
2.2. Sampling methods	<b>11</b>
2.3. Data analyses	<b>14</b>
<b>3. Results</b>	
3.1. Composition and Structure of the Anuran Community	<b>14</b>
3.2. Environmental Variables	<b>19</b>
3.3. Anuran Community vs. Environmental Variables	<b>20</b>
<b>4. Discussion</b>	
4.1. Habitat Structure and Composition of the Anuran Community	<b>21</b>
4.2. Anuran Conservation	<b>24</b>
<b>5. Acknowledgments</b>	<b>31</b>
<b>7. References</b>	<b>33</b>
<b>Conclusão Geral</b>	<b>45</b>
<b>Referências Bibliográficas</b>	<b>47</b>
<b>Anexo</b>	<b>53</b>

# I INTRODUÇÃO GERAL

munidades são moldadas por interações bióticas (Pianka 1973) e podem responder de maneira independente aos mesmos fatores ambientais (Homes et al. 1986). A identificação das funções que tais fatores desempenham na determinação dos padrões de distribuição e abundância das espécies é o primeiro passo para a compreensão dos processos de inter-relação das espécies com o ambiente (Underwood et al. 2000).<sup>[AMP1]</sup>

Os padrões de organização das comunidades refletem predisposição social (Ricklefs 2001), requerimentos específicos de habitats (Pianka 1967; Willson 1974), distribuição de recursos (Duellman 1990; Mesquita et al. 2006) e magnitudes das interações sociais (Sredl 1992; Eterovick 2003), sendo que muitas vezes estas ações agem em conjunto, o que dificulta o estabelecimento de relações de causalidade (Begon 2006). Entretanto, a existência ou a aparente falta de padrões na estrutura das comunidades pode refletir condições históricas que atuaram no passado e que não agem mais nos dias atuais (Inger 1969; Connell 1980).

Nenhum sistema ecológico pode ser estudado isoladamente do ambiente no qual está inserido (Begon et al. 2006). A heterogeneidade estrutural do ambiente é um dos fatores que tem sido reconhecido por apresentar associação com a riqueza de espécies em diferentes grupos zoológicos (MacArthur & MacArthur 1961; Parris 2004). É esperado que em áreas com maior disponibilidade de microambientes e nichos



ecológicos ocorra maior riqueza de espécies (Putman 1996), já que estes ambientes permitiriam a coexistência de um número maior de espécies, quando comparados a ambientes mais homogêneos (Cardoso et al. 1989; Gascon 1991). No entanto, a depender deve-se ponderar que o efeito da heterogeneidade ambiental pode variar consideravelmente dependendo do que é reconhecido como hábitat para o táxon estudado (Tews 2004).

A interação fisiológica dos anfíbios com seu ambiente é um sistema complexo e dinâmico de processos relacionados (Duellman & Trueb 1986), difíceis de serem isolados. Essa estreita relação faz os anfíbios serem considerados bioindicadores de variação nas condições ambientais, uma vez que suas características ecofisiológicas (p. ex. pele permeável e ciclo de vida bifásico) possibilitam marcada suscetibilidade a alterações de parâmetros físico-químicos da água, bem como da estrutura da vegetação que os circundam (Vitt et al. 1990; Wake 1991).

Abordagens descritivas ainda representam grande parte do esforço em trabalhos de pesquisa com anuros (p.ex. Duellman 1988; Feio & Ferreira 2005; Uetanabaro 2007; Vieira et al. 2007) e as relações investigadas entre a estrutura da comunidade e as variáveis do ambiente muitas vezes limitam-se a associações com temperatura (p.ex. Bertoluci 1998), pluviosidade (p.ex. Rossa-Feres et al. 1999; Bernarde 2007; Bernarde & Kokubum 1999) ou a ambas (p.ex. Pombal Jr. 1997; Conte & Machado 2005; Zina et al. 2007). Poucos são os estudos que buscaram relacionar os padrões de distribuição das espécies e os fatores ambientais que os influenciam. Associações positivas da complexidade estrutural da vegetação com a riqueza, diversidade ou composição de espécies foram encontradas por Parris & McCarthy (1999), Vallan (2002), Krishnamurthy (2003), Parris (2004) e Bastazini et al. (2007), embora em Santos et al.

(2007) o hidroperíodo tenha sido considerado dentre os descritores de complexidade estrutural o que melhor representou a riqueza registrada nos ambientes estudados.

O domínio morfoclimático das caatingas (ver Ab'Sáber 1977 para definição dos domínios morfoclimáticos) é um dos domínios brasileiros menos investigados para os anuros (Rodrigues 2003). Estudos ecológicos que versem especificamente sobre estrutura de comunidades de anfíbios no domínio das caatingas são escassos, pontuais e, por vezes, limitam-se a monografias, dissertações e teses não publicadas (p. ex. Cascon 1987; Lima 1999). Isso é consistente para o Parque Estadual das Sete Passagens (PESP), localizado na Serra de Jacobina, designação regional da Serra do Espinhaço, representando a vertente norte da Ecorregião “Complexo da Chapada Diamantina” (ver Velloso et al. 2002 para caracterização das Ecorregiões do bioma Caatinga). A Serra de Jacobina foi definida pelo Ministério do Meio Ambiente (Rodrigues et al. 2004) como área prioritária para conservação por apresentar “informação insuficiente”. O conhecimento proporcionado pela elaboração do Plano de Manejo do PESP (MF Napoli comunicação pessoal) ao qual esteve associada a execução desta dissertação, subsidiou alteração na categorização destas prioridades. A Serra de Jacobina (incluindo aí o PESP) foi definida pelo MMA (2007) como área com relevância extremamente alta para conservação devido à diversidade de fisionomias e registros de espécies novas, endêmicas, restritas e ameaçadas. O PESP está inserido em uma das áreas consideradas como zona de exceção do domínio das caatingas (ver Ab'Sáber 1977 para definição de zona de exceção), caracterizado pela presença de maciços montanhosos que podem ultrapassar 2000m de altitude e condições de temperatura e umidade que proporcionam a ocorrência de ecossistemas singulares, como os campos rupestres, as matas estacionais decíduais e as semi-decíduais montanas. Ambientes de altas elevações (>700m),

montanhosos e acidentados, como os encontrados no PESP, propiciam barreiras à dispersão de diversos grupos filogenéticos (Naniwadekar & Vasudevan 2007), apresentando localmente condições climáticas e geográficas que, refletidas na vegetação, justificam a presença de espécies singulares. O presente trabalho é o primeiro a investigar as relações entre a composição da comunidade de anuros e a estrutura do hábitat para a Ecorregião “Complexo da Chapada Diamantina”.

Diante da insuficiência de dados para o domínio das caatingas e do crescente consenso sobre o declínio nas populações de anfíbios no mundo (Crump et al. 1992; Marsh 2001; Funk & Mills 2003; Beebee & Griffiths 2005), há uma clara necessidade de estudos que além de descreverem a diversidade de espécies de ambientes desconhecidos para a ciência, procurem estabelecer as relações que estruturam estas comunidades bióticas. Os resultados obtidos nestes estudos podem subsidiar ações de manejo a serem adotadas por gestores ambientais e dar suporte a estratégias regionais de conservação em áreas prioritárias do bioma Caatinga, especialmente em regiões onde a disponibilidade de dados sobre diversidade e distribuição é incipiente.

A presente pesquisa tem como objetivos (1) determinar a composição e a distribuição espacial de uma comunidade de anuros em área montanhosa coberta por campos rupestres e florestas decíduas e semi-decíduas no domínio das caatingas e (2) buscar por variáveis ambientais que sejam capazes de explicar o padrão de distribuição espacial desta comunidade.

## II MANUSCRITO PARA APRECIACÃO

Este capítulo apresenta o manuscrito intitulado **“Influência de fatores ambientais na distribuição espacial de uma comunidade de anuros em área de altitude no domínio das caatingas”**, que se destina à submissão para publicação no periódico científico BIODIVERSITY AND CONSERVATION. Os resultados aqui discorridos, assim como a discussão e conclusões derivadas, decorrem do desenvolvimento da presente dissertação. Os critérios de redação e formatação seguem as normas deste periódico, as quais se encontram disponíveis na íntegra no ANEXO desta dissertação.

Running head: **Environmental Variables vs. Anurans Composition**

**Influence of environmental variables to the spatial distribution of an anuran community in a exception area within the Caatinga Domain**

**Ariane Lima Xavier • Marcelo F. Napoli**

Ariane Lima Xavier

Programa de Pós-Graduação em Ecologia e Biomonitoramento, Instituto de Biologia, Universidade Federal da Bahia, Rua Barão de Geremoabo 147, Campus Universitário de Ondina, 40170-290 Salvador, Bahia, Brazil

Phone: +55 71 3283-6562, Fax: +55 71 3283-6511, e-mail: [arianelx@yahoo.com.br](mailto:arianelx@yahoo.com.br)

Marcelo F. Napoli

Museu de Zoologia, Departamento de Zoologia, Instituto de Biologia, Universidade Federal da Bahia, Rua Barão de Geremoabo 147, Campus Universitário de Ondina, 40170-290 Salvador, Bahia, Brazil

Phone: +55 71 3283-6562, Fax: +55 71 3283-6511, e-mail: [napoli@ufba.br](mailto:napoli@ufba.br)

**Abstract** The present study searched for associations between an anuran community composition and environment variables to test which variable would better explain spatial patterns of anurans within a mountain environment covered by semideciduous forests and rocky mountain fields. The study area was a State Park in the Municipality of Miguel Calmon, Bahia State, Brazil, an area comprised in the Caatinga Morphoclimatic Domain. We sampled 20 plots by active method during two rainy seasons, and measured 13 environment variables. The environment matrix was reduced to four synthetic axes by the principal component analysis (PCA), and anuran community (abundance matrix) was reduced to a single synthetic axis by the nonmetric multidimensional scaling method (NMS). We tested the absence of association between environment axes (PCA) and community composition (NMS axis) by multiple regression analysis. The PC1 (40%) was the only axis significantly associated to anuran composition, in which variables with major contributions were directly related to the vegetation structure. The anuran community was ordered along an environment gradient, from open areas featured by rocky mountain fields (“Campo Rupestre”) to primary semideciduous mountain forest patches. Anuran composition in Sete Passagens State Park is very distinct from those in Caatinga, which denotes the importance in considering both phytogeographies to create protection areas in the Brazilian semi-arid.

**Keywords** Anuran community structure · Brazilian semi-arid · Campo Rupestre · Chapada Diamantina · Habitat gradient · Landscape context · Semideciduous forest

## **Abbreviations**

PESP Sete Passagens State Park

SU Sample Unit

NMS Nonmetric Multidimensional Scaling

PCA Principals Components Analysis

## **Introduction**

The identification of factors that regulate spatial distribution patterns and abundance of species is an important step to realize the processes that determine the communities' biologic diversity (Magurran 1988; Norton 1991; Underwood et al. 2000). Such patterns come from different processes, reflecting specific requirements of habitats (Pianka 1973; Sredl 1992), resources distribution (Duellmann 1990; Mesquita et al. 2006), and extension of the social interactions (Connell 1980; Eterovick 2003).

Regardless of the importance on conduct researches to find the structural processes that shape the biotic communities, descriptive approaches still represent a huge part of the effort in research works on anurans in Brazil (e.g. Feio and Ferreira 2005; Santos et al. 2005; Uetanabaro 2007). Several studies related community structure to environmental variables and indicated associations with temperature (e.g. Bertoluci 1998), rainfall (e.g. Bernarde and Kokubum 1999; Rossa-Feres et al. 1999; Bernarde 2007), or both (e.g. Pombal Jr. 1997; Conte and Machado 2005; Zina et al. 2007).

However, these studies did not provide general explanations for the organization patterns found at anuran communities. In Brazil, only few studies on these communities searched for relationships between patterns of species distribution and environmental variables (e.g. Giaretta 1999; Bastazini et al. 2007; Afonso and Eterovick 2007). Such studies were carried out mainly to the Atlantic Forest Biome and studies focusing on anuran communities in the Brazilian semi-arid are still rare (e.g. Arzabe et al. 1998).

The Caatinga Morphoclimatic Domain (see Ab'Sáber 1977 for definition of Morphoclimatic Domain), comprised within the Brazilian semi-arid, is highlighted as the most unknown area regarding its taxa composition, turning more difficult the development of ecological studies. The exception zones distributed within this domain (see Ab'Sáber 2003 for definition of exception zones) are known as areas of extremely high relevance for conservation due to their singular physiographic features, climate, and association with endemic, threatened, and/or new species. Up till now, there are no studies conducted in a landscape of the ecoregion Chapada Diamantina Complex (see Velloso et al. 2002 for definition of the ecoregions from Brazilian Caatinga), an exception area, focusing on the landscape context of spatial distribution of anuran communities.

Ecological investigations on spatial distribution of anurans, and the understanding of how environmental variables influence the anuran community allows to know of main ecological demands for the species survival. In this scope, the results obtained here can help finding management actions, and to support regional strategies of conservation in priority areas of the Caatinga biome, especially in regions where there is a poor availability of data about diversity and distribution.



The aims of the present research are: (1) to analyze the composition and spatial distribution of an anuran community in an exception area known as “Campo Rupestre” (a rocky mountain area, characterized by stony fields and Deciduous and Semideciduous Forests) in the Caatinga Morphoclimatic Domain, and (2) to search for environmental variables able to explain the spatial distribution pattern of this community.

## **Material and methods**

### **Study area**

The present study was carried out at the Sete Passagens State Park (Parque Estadual das Sete Passagens – PESP; 11°26'S, 40°33'W), in the Municipality of Miguel Calmon, State of Bahia, Brazil (Fig. 1). This park is placed in the ecoregion Chapada Diamantina Complex (Velloso et al. 2002), comprised in the Caatinga Morphoclimatic Domain (Ab'Sáber 1977), the latter with vegetation morphologically and physiologically adapted to semiarid conditions. At the PESP, elevations ranged from approximately 800 m to 1300 m, with an estimated area of 2821 ha, being part of the residual group of mountains named Serra de Jacobina, a regional designation of the northern portion of the Serra do Espinhaço mountain range (see Davis et al. 1997 for characterization of the Serra do Espinhaço mountain range). The topography is represented by a sequence of edges and aligned bars, with deep kerfs, steep cliffs, short flat peaks, and canyons. This

topography is classified as extremely unstable with medium slopes ranging from 30° to 45°, excepting for the tops. Almost all rivers surrounding the Park have their headsprings inside its boundaries. In accordance to Köppen classification, the weather is of the BSw type, semi-arid, rainy in summer and dry in winter. The rainy period is from October to April, and the rainiest months are November to January. Due to the complex geomorphology and its elevated altitudes, PESP temperature reaches less than 14° C in winter, establishing a difference with its surroundings, where the minimum temperature recorded is 20° C. Such characteristics are reflected in the plant cover, composed by seasonal Semideciduous Forests and mountains immersed in a matrix of Campo Rupestre (the same of Campos Rupestres Montane Savanna of WWF 2008; see also Rizzini 1979 for the characterization of the Brazilian Campo Rupestre), which contrast with lowland areas (up to ca. 500 m de altitude) surrounding the Park that present shallow soils covered by a typical Caatinga landscape (see Ab'Sáber 2003 for the characterization of the Caatinga landscape).

### Sampling methods

The sample design followed Bastazini et. al. (2007). We distributed 20 sample units (SUs) (plots: P1–P20), measuring 60 x 25 m each, within the overall landscape of the PESP, composed by areas of Campo Rupestre and patches of seasonal Semideciduous Forests (Fig. 1). The sample units were placed along existing trails, as the extremely hilly topography led to access difficulties, mainly at night, preventing the equivalence between the amount of plots distributed in forest patches and areas of Campo Rupestre. Yet, as plots were placed with a minimum arbitrary distance of 300 m to avoid violation

of the premise of plots independence, the available sampling area for each main phytophysiology was reduced. To control for sampling biases in searching for relationships among SUs, we categorized each plot by its main microhabitat features only after statistical analyses.

We sampled SUs in two periods of consecutive rainy seasons, as follows: 10 SUs from 18–22 November 2006 (P1–P10), and 10 SUs from 16–20 January 2007 (P11–P20). Each sampling period lasted for five days. To control for temporal biases in the characterization of the main landscape units (forested areas vs. rocky open areas), in each sampling occasion we distributed 40% of the SUs in open areas and 60% in forested areas. Therefore, environmental temporal variation will affect the sampling of the two main physiognomies similarly, and habitat or microhabitat effects will not be confused with time variation. We looked for anurans in each SU during 40 min a day, after the 1800 h, totalling 6 h and 40 min of active search per day (total effort of 66 h and 40 min considering all SUs and both sampling periods). We covered the SUs with homogeneous effort with a team of three pairs of researches, which were not changed during the study. These pairs of observers and the sequence of visitations to SUs were changed randomly during the sampling period.

We measured 13 environmental variables in each SU. (1) Air moisture and (2) air temperature, both measured every night in each SU before the 40 minutes of active search, using a thermo-hygrometer Minipa MT 242. (3) Estimated percentage of water cover in the SU. (4) Type of water body in the SU: we calculated the average velocity achieved by a matchstick (0.082 g and 3.5 cm long) through 1 m of water course for three consecutive times, and then the arithmetic mean; the same matchstick and chronometer were used for this measurement in all SUs with water bodies. (5)

Maximum depth and (6) width of the water body. (7) Number of terrestrial and (8) epiphytic bromeliads. The next five environmental variables were taken from 10 points chosen at random in the SU, following a point map used for all SUs (Fig. 2); points were not changed during the study. A circle of 3 m diameter was drawn around each point, and the following variables were measured. (9) Soil moisture: 10 soil samples were taken from the SU, one per point; samples were mixed and sealed in a plastic bag. Moisture was measured by the “Moisture at 65° C” analysis (Embrapa 1997). (10) Percentage of leaf litter and (11) percentage of understory, shrub, and herb layers (covering of forest layers between canopy and litter layers); ordinal variables ranging from 0 to 4: 0 (0%), 1 (1–25%), 2 (25–50%), 3 (50–75%), and 4 (75–100%). (12) Canopy height, ordinal variable, ranging from 0 to 4: 1 (1–5 m), 2 (5–10 m), 3 (10–15 m), and 4 (>15 m). (13) Density of trees (point-quarter method; Krebs 1999).

In order to search for differences in anuran composition and species richness between the Sete Passagens State Park — PESP (an exception area) and the nearby Caatinga landscapes (xeric plant cover), we surveyed for anurans within adjacent and contiguous areas in the Municipality of Miguel Calmon, Bahia State, in 09–11 October 2006 and 17–19 February 2007. In addition, we obtained data on a collection of anurans from another contiguous area to PESP, with the same plant cover type (Municipality of Jacobina, Itaitu, 11°19'S, 40°28'W; ca. 570 m altitude, 18–20 April 2003, by M. F. Napoli), deposited in the Amphibians' Collection of the Museu de Zoologia da Universidade Federal da Bahia (UFBA).

The specimens captured were deposited in the Amphibians' Collection of the Museu de Zoologia da Universidade Federal da Bahia (UFBA). Names of the taxa followed

Faivovich et al. (2005), Frost et al. (2006), Grant et al. (2006), Chaparro et al. (2007), and Hedges et al. (2008).

### Data analyses

For the purpose of analysis, we considered only SUs with the presence of anurans ( $n = 16$ ; P7–P9 and P19 were discarded). We produced four data matrices: (A) a matrix of 16 SUs (plots) vs. 15 anuran species (attributes), representing the dependent variable of the study, namely community composition. We used the “concrete community” concept of McCune and Grace (2002), which is characterized as the collection of organisms found at a specific place and time. In the matrix, each cell corresponded to the number of specimens of the species in the SU. (B) A matrix with relative abundances, calculated from the matrix A by dividing each cell by the line sum to reduce differences in weight of sample units in analyses. (C) A second matrix of relative abundances, calculated from the matrix A by dividing each cell by the column sum to reduce differences in weight of species in the analysis. (D) A matrix with 16 SUs vs. 13 environmental variables, representing the independent variable in the analysis. We considered the relative number of specimens collected per species as an adequate comparative index of plot composition, since the sample effort was the same in the sample units.

We used the Nonmetric Multidimensional Scaling method (NMS), an ordination technique, to seek and display the strongest anuran community structure using only the anuran species data set (matrix B) (McCune and Grace 2002). We selected only one dimension (ordination axis) to the NMS solution, and used the Bray-Curtis distance measure. To avoid the local minima problem, we made 50 starting configurations, using

as stability criteria the instability value of 0.0005, 20 iterations to evaluate the stability of the solution, and 500 as the maximum number of iterations. The Monte Carlo test (a randomization test) was used to evaluate whether the NMS extracted a stronger axis than expected by chance (500 iterations for the randomized results). The proportion of variance represented by the NMS axis, based on the correlation between distance in the ordination space (Euclidian distance) and distance in the original space (Sorensen distance), was obtained by a standardized Mantel test ( $r$ ). We removed *Ischnocnema aff. ramagii* from the analysis because this species turned the test unfeasible (its inclusion led to a high final stress in the NMS solution). This taxon was represented in all 16 SUs considered for the analysis, showing a high abundance in forested areas, as well as in Campo Rupestre. As the final stress of the NMS ordination depends on the quantity of objects used (plots) and variables (species), very abundant species regularly distributed within almost all replicas can influence the quality of ordination and, therefore, the final stress. We followed McCune and Grace (2002) in removing this taxon from the analysis, as the removal of species occurring in more than half SUs leads to a lower final stress. The quality of the indirect ordination of the anuran community obtained from the NMS solution was represented in a composite graph (Fig. 3), using projections of the relative abundances of each species vs. the NMS axis (plots) (matrix C).

The null hypothesis of absence of association between the community composition (NMS axis) and the spatial distribution of SUs was tested by the Spatial Autocorrelation Analysis (Sokal and Oden 1978). A spatial correlogram was constructed using Moran's  $I$  coefficients in four distance classes, whose upper bound limits (in km) were: 1.3, 2.1, 3.1, and 6.4. These limits were chosen irregularly, but in a way that each distance class contained approximately the same number of distance pairs of SUs. For each distance

class we computed a correlation coefficient, and the statistical significance of the overall correlogram was tested by the Bonferroni criterion, in which a correlogram is considered significant if at least one of its Moran's I is significant at  $P \leq 0.05/k$ , where "k" is the number of distance classes. Significant and positive Moran's I coefficients indicate that plots a given distance apart are similar for the variable under analysis (NMS axis: community composition), while significant and negative Moran's I values indicate dissimilar plots for that variable at this distance (Diniz-Filho et al. 2000). The result of the spatial autocorrelation analysis was graphically represented by a spatial correlogram, where the values of the Moran's I coefficient were plotted vs. a horizontal axis representing the geographical distance classes among SUs.

We used the Principals Components Analysis (PCA) to reduce the environmental data set (matrix D) to a smaller number of orthogonal synthetic variables that represent most of the original information. The principal components were obtained from a correlation matrix, and those with eigenvalues  $\geq 1$  were rotated to a new Varimax solution (McCune and Grace 2002).

The null hypothesis of absence of association between the environmental data set (principal components) and the anuran community composition (NMS axis) was tested by multiple regression analysis. The significance value adopted was of  $P \leq 0.05$ . In cases with multiple comparisons, working with the same set of data, the  $P$  critic value was corrected by the Bonferroni method (Bland 2004).

## Results

### Composition and Structure of the Anuran Community

We sampled in the PESP 567 anurans belonging to 5 families, 9 genera, and 15 species. The family with the highest amount of specimens was Hylidae, followed by Brachycephalidae. The five most abundant species were *Ischnocnema* aff. *ramagii* (44.8%), *Ischnocnema* sp. (16.6%), *Bokermannohyla oxente* (14.5%), *Scinax* sp. (6.3%), and *Proceratophrys* aff. *cristiceps* (4.4%), corresponding together to 86.6% of the total sampled individuals. Out of nine SUs located at Campo Rupestre, only five presented anurans, which were distributed within six species (40% of the species in PESP). Anurans occurred in all forested SUs. *Scinax pachycrus* was the only species not found in forested areas, and *Bokermannohyla* cf. *circumdata*, *Proceratophrys* aff. *cristiceps*, *Rhinella crucifer*, *Scinax eurydice*, *S.* aff. *catharinae*, and *Trachycephalus* aff. *mesophaeus* were restricted to forested areas. *Bokermannohyla oxente*, *Scinax* sp., and *Ischnocnema* sp. were the most widespread species in PESP, the former two mainly associated to forested environments, while the latter to Campo Rupestre.

Comparisons of species richness and composition of anurans between the PESP and adjacent areas covered by Caatinga vegetation evidenced very distinctive anuran communities (Table 1). The only species in common were *S. pachycrus*, *Hypsiboas crepitans*, *H. faber*, and *Phyllomedusa bahiana*, all usually related to forest borders and/or open habitats with lentic water bodies.



The NMS axis was able to express structure in the anuran community of the PESP (Fig. 3). Although stress associated with NMS axis was high (24.5), it is not implied that the obtained stress leads to a low level of interpretation. According to McCune and Grace (2002), an increase in sample units leads to a rise in stress (>15%), which does not interfere with the results interpretation. The variance represented by the NMS axis explained 68% of the variance in the original multidimensional space (Mantel test:  $r = 0.81$ ,  $P \leq 0.001$ ). The Monte Carlo test resulted in a probability of 0.02, indicating that the NMS extracted a stronger axis than expected by chance. The NMS axis ordered the anuran community through an environmental gradient that reflects six main landscape categories. Category 1 (P20): Campo Rupestre, mostly (80%) an open area, marked by sparse treelets and shrubs with thick bark, and also with a patch of Deciduous Forest (20%), with shallow ground and a narrow and deep water well. Category 2 (P10, P16, P17, and P18): similar to the former category, but without water bodies on the ground. *Scinax pachycrus* was exclusive to the former two categories, in which *Ischnocnema* sp. reached its highest abundance values. Category 3 (P6 and P13): hybrid areas, characterized by mixing features of open areas (herbaceous vegetation and bare rocks) and riparian forests, not dense, with understory layer and water bodies on the ground. *Hypsiboas crepitans*, *H. faber*, *Scinax* sp., *Phyllomedusa bahiana*, and *Proceratophrys* aff. *cristiceps* were associated to this landscape category, often captured inside forested areas, but also in forest edges and open areas. The forest cover of categories 3 and 4 (see below) suffered different degrees of deforestation by human activities in the last 20 years, and is at present in an initial stage of natural forest recovery. Category 4 (P3 and P4): forest areas without water bodies on the ground, with many epiphytic bromeliads, herbaceous vegetation, and bryophytes. *Strabomantis aramunha* was restricted to plots

of this landscape category, but also captured from open areas external to the sampled plots in cavities under rocks on the side of a stream. Categories 5 (P1, P5, P14, P15, and P11) and 6 (P2 and P12): include the largest forest patches in the PESP covered by dense vegetation. These areas are somewhat inaccessible for human activities due to their long trails in hard topographies, which are represented by a sequence of edges and aligned bars, with deep kerfs, steep cliffs, short flat peaks, and canyons. Category 5 differs from category 6 by the presence of water bodies on the ground, absent in the latter. *Bokermannohyla* cf. *circumdata*, *Rhinella crucifer*, and *Scinax* aff. *catharinae* were restricted to these latter two categories.

### Environmental Variables

The PCA applied to 13 environmental variables generated four principal components with eigenvalues larger than 1.0 (Table 2). They represented 84.7% of the entire variance after the varimax solution. The first principal component accounted for 40.0% of the total variance, and presented high loadings ( $\geq 0.7$ ) on variables related to the vegetation structure; in decreasing order: (1) percentage of leaf litter, (2) density of trees, and (3) canopy height; (4) soil moisture (loadings = 0.68). In the second axis (19.5%), the variables that mostly contributed were associated to water bodies on the ground, as follows: (1) maximum width of the water body, (2) maximum depth of the water body, (3) estimated percentage of water cover in the SU, and (4) type of body of water in the SU. In the third axis (14.8%), out of three most important variables, two were related to atmospheric features: (1) air temperature, (2) air moisture, and (3)

number of epiphytic bromeliads. The fourth axis (9.6%) was strongly and positively related to the number of terrestrial bromeliads.

The projection of PC1 on PC2 discriminated three groups of SUs (Fig. 4): (1) SUs with water bodies on the ground (P1, P5, P6, P11, P13–P15, P20); (2) SUs without water bodies on the ground in open areas (P10, P16–P18); and (3) SUs without water bodies on the ground in forested or transitional areas (P2–P4, P12). Within areas without water bodies on the ground, the ordination of SUs followed a plant cover gradient throughout PC1, with moderately forested SUs (P3 and P4) placed between non forested (P10, P16–P18) and forested plots (P2 and P12). A similar plant cover gradient was observed for SUs with water bodies on the ground: (1) open areas: 80% of the SU covered by a matrix of Campo Rupestre (P20); (2) transitional areas: riparian forest (55% of the SU) along with a matrix of Campo Rupestre (45% of the SU), the latter covered by herbaceous vegetation (P13) or by rocks and bromeliads (P6), being that P13 with greater forest layering than P6; (3) forested areas (P1, P5, P11, P14–A15): with the highest density of trees, diameter at breast height reaching 40 cm, and canopy reaching 15 m high

#### Anuran Community vs. Environmental Variables

We tested the null hypothesis of association absence between the environmental data set (independent variables: principal components with eigenvalues  $\geq 1.0$ ) and anuran community composition (dependent variable: NMS axis) using a multiple regression analysis, which was strongly significant ( $P < 0.0001$ ). However, out of four possible

partial regressions, only PC1 vs. the NMS axis was significant ( $P < 0.0004$ , Bonferroni corrected). The other three partial plots showed no regression pattern. The partial regression plot of PC1 on the NMS axis (Fig. 5) accounted for 82% of the variation and reflected an environmental gradient, that is, from open areas of Campo Rupestre to forest environments. Considering only the forested areas, there is a environmental gradient from hybrid areas (P6, P13) to full forested areas (P1–P5, P11–P12, P14–P15), determined by gradual increase of leaf litter, density of trees, and canopy height, and gradual decrease of understory, shrub, and herb layers.

The distribution pattern of the anuran community (NMS axis) was not related to the spatial distribution of SUs, as none of the four Moran's I coefficients calculated for the dataset (community composition; 1 variable vs. 4 distance classes) were significant at the 5% level or less, as well as the overall correlogram ( $P > 0.15$ ), and therefore not rejecting the null hypothesis of absence of spatial autocorrelation among the SUs.

## **Discussion**

### **Habitat Structure and Composition of the Anuran Community**

The variables that markedly related to the spatial distribution of anuran species in the Sete Passagens State Park were primary associated to the vegetation structure, describing an environmental gradient from open areas of Campo Rupestre to forested areas. Forest patches presented greater species richness than those with direct sunlight,

probably due to the structural complexity granted by forests, which provide more predictable (Colwell 1974), complex (Putman 1994), and heterogeneous habitats than shrubby environments (see Tews 2004 for a brief review on the relation between species diversity and habitat heterogeneity).

Similar results were found by other studies that searched for environmental variables to explain richness, diversity or species composition in anuran communities (e.g. Parris and McCarthy 1999; Atauri and Lucio 2001; Vallan 2002; Bastazini et al. 2007). On the other hand, Vasconcelos and Rossa-Feres (2005), studying an anuran community from an anthropized area originally covered by seasonal forests in the State of São Paulo, Brazil, did not find association of anuran species richness with habitat heterogeneity, but found with air temperature and air moisture; Santos et al. (2007), analysing a similar area of Vasconcelos and Rossa-Feres (2005), stated that the hydroperiod was the environmental variable that better explained the anuran species richness. Nevertheless, these studies were done in highly anthropized habitats, i.e. small secondary forest patches surrounded by pastures. Moreover, the environmental variables used to describe the habitat heterogeneity were associated only to water bodies on the ground, as inner vegetation, marginal vegetation, and vegetation surrounding water bodies, clearly denoting the lack of complex forest stratification (e.g. litter, herb, shrub, understory, and canopy layers).

In the Campo Rupestre almost all anurans were captured in terrestrial bromeliads. Bromeliads are considered in the literature important ecological units, as they provide moisture, shelter, and breeding sites, mainly in environments with high air temperatures, low soil and air moistures, and absence of water bodies on the ground (Dejean and Olmsted 1997; Richardson 1999; Shineider and Teixeira 2001; Teixeira et al. 2002).

However, terrestrial bromeliads were not able to explain the anurans spatial distribution in the studied area. This result could be due to the homogeneous distribution of these plants along the SUs, without leading to a gradient or a clear cut distribution pattern among SUs, as that encountered by Bastazini et. al. (2007). These authors studied an anuran community in a Brazilian Restinga (see Suguio and Tessler 1984 for a brief characterization of the Restinga environment), and found a clear cut pattern in the spatial distribution of terrestrial bromeliads among SUs (few terrestrial bromeliads per SU in forested areas, and more than a hundred in open areas of sand dunes habitats). For the Restinga landscape, terrestrial bromeliads were among the environmental variables that most explained the changes in anuran composition.

Most investigations on anuran communities are usually limited to water bodies on the ground, such as breeding ponds and streams (e.g. Arzabe et al. 1998; Arzabe 1999; Vasconcelos and Rossa-Feres 2005; Zina et al. 2007), mainly because water is conventionally considered the most important factor in shaping the spatial distribution of the anuran fauna. Nevertheless, this relation has been scarcely tested (e.g. Zimmerman and Bierregaard 1986; Eterovick 2003; Afonso and Eterovick 2007; Bastazini et al. 2007). In the present study, variables associated to water reservoirs on the ground (represented by PC2) were not associated to the spatial distribution of the anuran fauna. Despite of the anurans' dependence on water bodies that function as breeding sites for many species, environments with more diversified vegetal stratification lead to low direct sunstroke, high availability of microhabitats, and wet soils with high accumulation of organic materials, which together provide environmental conditions of moisture and temperature that are sufficient to support a large variety of adult forms of anurans. The anuran spatial distribution therefore were

not regulated only by the presence of water bodies, though many species need to reach them over a period of their life cycles to deposit their egg clutches, but mainly by the degree of vegetal layering of that habitat. This statement is supported by the results of Bastazini et al. (2007), in which change in plant communities was the most important environmental factor acting on the structure of an anuran community of a Restinga habitat.

### Anuran Conservation

The anuran species richness of the Sete Passagens State Park (PESP) (15 species) is similar to other mountain regions in the Caatinga Domain (12–18 species; Azarbe 1999; Borges-Nojosa and Cascon 2005; Vieira 2007). On the other hand, its richness is noticeably smaller than the overall Chapada Diamantina Complex (44 species; Juncá 2005), once the latter includes a larger amount of distinct landscapes (e.g. Caatinga, Campo Rupestre, Cerrado, and Deciduous and Semideciduous Forests).

The Chapada Diamantina region is one of the five biogeographic sub-regions of endemism proposed by Galindo-Leal and Câmara (2005) for the Tropical Atlantic Forest (based on birds, primates, and butterflies). The PESP is placed in the Bahia's north-western limit of the original Tropical Atlantic Forest covering, but its anurans composition is related to a mixture of distinct phytophysionomies. *Scinax pachycrus* and *Hypsiboas crepitans* are common species to xeric environments of the Caatinga Domain, while *Bokermannohyla* cf. *circumdata*, *Hypsiboas faber*, *Ischnocnema* aff. *ramagii*, *Phyllomedusa bahiana*, *Scinax* aff. *catharinae*, *Scinax eurydice*, and *Rhinella*

*crucifer* are related to the Atlantic Forest Domain (Fig. 5). One particular record deserves to be highlighted. The record of *Bokermannohyla* cf. *circumdata* represents the northernmost limit of the *B. circumdata* species group, which is distributed mainly over mountain stream habitats in the Atlantic Tropical Forests, occurring over 350 m (Napoli and Caramaschi 2004). Species of the group are also observed inhabiting gallery forests within high elevation fields in the Caatinga, Cerrado, and Tropical Atlantic Forest domains (e.g. *B. diamantina*, *B. circumdata*, *B. feioi*, *B. ibitipoca*, *B. nanuzae*, *B. ravida*, and *B. sazimai*) (Napoli and Juncá 2006). Up till now, *B. diamantina* was the only species of the group known for the Brazilian semi-arid (type-locality Serra do Barbado, Chapada Diamantina, Bahia State). The occurrence of *B. cf. circumdata* at the Serra de Jacobina mountain encloses an important biogeographical and conservational value, since the presence of this taxon points to a relict of the original Atlantic Forest range, and also set the northern limits of the *B. circumdata* species group. Napoli and Juncá (2006) had already stated that the occurrence of species of the *B. circumdata* group in Cerrado and Caatinga domains is due to the presence of gallery forests, which provide, in some degree, environmental conditions similar to the Atlantic Tropical Forests from coastal Brazil, and therefore these species should not be considered as characteristic taxa of Cerrado and Caatinga biomes.

The anuran species composition of the PESP, when compared to the anuran fauna of the surrounding open area covered by xeric vegetation (Caatinga plant cover) (Table 3), places the former almost as a forest island in the Caatinga landscape (see Vanzolini 1981 for an interesting discussion of the current mesic enclaves found in northeastern Brazil). Almost all anuran species from the surrounding open area are inhabitants of permanent and temporary ponds, sometimes invading forest edges, while anurans from



the PESP are usually related to forests. Five species were common to both areas, all of them related to open or transitional areas with or without lentic environments (*B. oxente*, *Scinax pachycrus*, *Hypsiboas crepitans*, *H. faber*, and *Phyllomedusa bahiana*). *Corythomantis grenningi*, a hylid tree frog widespread in the Caatinga Domain, was only found in a patch of forested area out of the PESP limits, bordering its base, and contiguous to the surrounding xeric Caatinga plant cover. These dry forest patches are physiognomically different from the mountain forests within the PESP, but is not under protection, as usually occurs for non mountainous landscapes in the Caatinga Domain. The PESP should not be understood as an isolated unit, but instead as ecologically and historically related to the adjacent Caatinga plant cover, lowland areas, and dry forests, which points out to the importance to include these different, but contiguous, phytophysionomies within new environmental protection areas in the Brazilian semi-arid.

**Acknowledgments** We are grateful to Miguel C. Accioly for the invitation to join the team that formulated the management plan of the Sete Passagens State Park (PESP); to Deise Cruz, Milena Camardelli, Patricia M. Fonseca, Thais A. F. Dória, Camila V. Bastazini, Herverton Cardona, Marina Bonfim, Natália Menezes, Wilton Fahning, Tasso Meneses, Thiago Filadelfo, and Tiago Jordão for the excellent field assistance; to Deise Cruz, Flora A. Juncá, and Rafael O. Abreu for photographs; to José M. Pereira (Zélis) for logistical assistance at the PESP. We acknowledge the Fundação de Amparo à Pesquisa do Estado da Bahia (FAPESB) for fellowships to Ariane L. Xavier, the Secretaria do Meio Ambiente (SEMA) for provide spatial data of the PESP; and the

Instituto Brasileiro do Meio Ambiente e dos Recursos Naturais Renováveis (IBAMA) (process numbers 02006.001462/2006-13 and 02006.002336/03-26), the SEMA, and local administration of the PESP for permitted collection of specimens.

## References

- Ab'Sáber AN (1977) Os domínios morfoclimáticos na América do Sul. *Geomorfologia* 52:1–22
- Ab'Sáber AN (2003) Os domínios de natureza no Brasil - potencialidades paisagísticas. Ateliê Editorial, São Paulo
- Afonso LG, Eterovick PC (2007) Spatial and temporal distribution of breeding anurans in streams in southeastern Brazil. *Journal of Natural History* 41:949–963
- Arzabe C (1999) Reproductive activity patterns in two different altitudinal sites within the Brazilian Caatinga. *Revista Brasileira de Zoologia* 16:851–864
- Arzabe C, Carvalho C X, Costa MAG (1998) Anuran Assemblages in Crasto Forest Ponds (Sergipe State, Brazil): Comparative Structure and Calling Activity Patterns. *Herpetological Journal* 8:111–113
- Atauari AA, Lucio JV (2001) The role of landscape structure in species richness distribution of birds, amphibians, reptiles and lepidopterans in Mediterranean landscapes. *Landscape Ecology* 16:147–159
- Bastazini CV, Munduruca JFV, Rocha PLB, Napoli MF (2007) Which environmental variables better explain changes in anuran community composition? A case study in the restinga of Mata de São João, Bahia, Brazil. *Herpetologica* 63:459–471

- Bernarde PS, Kokubum MNC (1999) Anurofauna do município de Guararapes, estado de São Paulo, Brasil (Amphibia, Anura). *Acta biológica leopoldensia* 21:89–97
- Bernarde PS (2007) Ambientes e temporada de vocalização da anurofauna no Município de Espigão do Oeste, Rondônia, Sudoeste da Amazônia - Brasil (Amphibia: Anura). *Biotaneotropica* 7:87–92
- Bertoluci JA (1998) Annual patterns of breeding activity in Atlantic Rainforest anurans. *Journal of Herpetology* 32:607–611
- Bland M (2004) Multiple significance tests and the Bonferroni correction. <http://www-users.york.ac.uk/~mb55/intro/bonf.htm>. Cited 04 Nov 2008
- Borges-Nojosa DM, Cascon P (2005) Herpetofauna da área Reserva da Serra das Almas, Ceará. In: Araújo FS, Rodal MJN, Barbosa MRV (orgs) *Análise das variações da biodiversidade do bioma caatinga: suporte a estratégias regionais de conservação*. Ministério do Meio Ambiente, Brasília
- Chaparro JC, Pramuk JB, Gluesenkamp AG (2007) A new species of aboreal *Rhinella* (Anura:Bufonidae) from cloud forest Southeastern Peru. *Herpetologica* 63:203–212
- Colwell RK (1974) Predictability, constancy and contingency of periodic phenomena. *Ecology* 55:1148–1153
- Connell JH (1980) Diversity and the coevolution of competitors, or the ghost of competition past. *Oikos* 35:131–138
- Conte EC, Machado RA (2005) Riqueza de espécies e distribuição espacial e temporal em comunidade de anuros (Amphibia, Anura) em uma localidade de Tijucas do sul, Paraná, Brasil. *Revista Brasileira de Zoologia* 22:940–948

- Davis SD, Heywood VH, McBride OH, Villa-Lobos J, Hamilton AC (1997) Centres of plant diversity: a guide and strategy for their conservation. WWF/IUCN, Cambridge
- Diniz-Filho JAF, Coelho ASG, Sant'ana CER (2000) Statistical inference of correlated evolution among macroecological traits using phylogenetic eigenvector regression. *Ecology Austral* 10: 27–36
- Duellman WE (1990) Herpetofaunas in Neotropical rainforests: comparative composition, history and resource use. In: Gentry AH (ed) *Four Neotropical Rainforests*. Yale University Press, New Haven
- Dejean A, Olmsted I (1997) Ecological studies on *Aechmea bracteata* (Swartz) (Bromeliaceae). *Journal of Natural History* 31:1313–1334
- EMBRAPA – Empresa Brasileira de Pesquisa Agropecuária (1997) *Manual de Métodos de análises de solo*. Embrapa, Rio de Janeiro
- Eterovick PC (2003) Distribution of anuran species among montane streams in south-eastern Brazil. *Journal of Tropical Ecology* 19:219–228
- Faivovich J, Haddad CFB, Garcia PCA, Frost DR, Campbell JA, Wheeler WC (2005) Systematic review of the frog family Hylidae, with special reference to Hylinae: phylogenetic analysis and taxonomic revision. *Bulletin of the American Museum of Natural History* 294:1–240
- Feio RN, Ferreira PL (2005) Anfíbios de dois fragmentos de Mata Atlântica no município de Rio Novo, Minas Gerais. *Revista Brasileira de Zoociências* 7:121–128
- Frost DR, Grant T, Faivovich J, Bain RH, Haas A, Haddad CFB, De Sá RO, Channing A, Wilkinson M, Donnellan SC, Raxworthy CJ, Campbell JA, Blotto BL, Moler

- P, Drewes RC, Nussbaum RA, Lynch JD, Green DM, WC Wheeler (2006) The amphibian tree of life. *Bulletin of the American Museum of Natural History* 297:1–370
- Galindo-Leal C, Câmara IG (2005) *Mata Atlântica: biodiversidade, ameaças e perspectivas*. Fundação SOS Mata Atlântica, São Paulo and Conservação Internacional, Belo Horizonte
- Giaretta AA, Facure KG, Sawaya RJ, Meyer JHM, Chemin N (1999) Diversity and abundance of litter frogs in a montane forest of Southeastern Brazil: seasonal and altitudinal changes. *Biotropica* 31:669–674
- Grant T, Frost DR, Caldwell JP, Gagliardo R, Haddad CFB, Kok PJR, Means DB, Noonan BP, Schargel WE, Wheeler WC (2006) Phylogenetic systematics of dart-poison frogs and their relatives (Amphibia: Athesphatanura: Dendrobatidae). *Bulletin of the American Museum of Natural History* 299:1–262
- Hedges SB, Duellman WE, Heinicke MP (2008) New World direct-developing frogs (Anura: Terrarana): molecular phylogeny, classification, biogeography, and conservation. *Zootaxa* 1737:1–182
- Juncá FA (2005) Anfíbios e Répteis. In: Juncá FA, Funch L, Rocha W (orgs) *Biodiversidade e Conservação da Chapada Diamantina*. Ministério do Meio Ambiente, Brasília
- Krebs CJ (1999) *Ecological Methodology*. Benjamin/Cummings, Menlo Park, San Francisco, California
- Magurran AE (1988) *Ecological Diversity and its Measurement*. Cambridge, University Press, London
- McCune B, Grace JB (2002) *Analysis of Ecological Communities*. Gleneden Beach, Oregon

- Mesquita DO, Costa GC, Colli GR (2006) Ecology of an Amazonian savanna lizard assemblage in Monte Alegre, Pará State, Brazil. *South American Journal of Herpetology* 1:61–71
- Napoli MF, Caramaschi U (2004) Two new species of the *Hyla circumdata* group from Serra do Mar and Serra da Mantiqueira, Southeastern Brazil, with description of the advertisement call of *H. ibitipoca* (Anura, Hylidae). *Copeia* 3:534–545
- Napoli MF, Juncá FA (2006) A new species of the *Bokermannohyla circumdata* group (Amphibia: Anura: Hylidae) from Chapada Diamantina, State of Bahia, Brazil. *Zootaxa* 1244:57–68
- Parris KM, McCarthy M (1999) What influences the structure of frog assemblages at forest streams? *Australian Journal of Ecology* 24:495–502
- Pianka ER (1973) The structure of lizard communities. *Annual Review of Ecology and Systematics* 4:53–74
- Pombal Jr JP (1997) Distribuição espacial e temporal de anuros (Amphibia) em uma poça permanente na Serra de Paranapiacaba, Sudeste do Brasil. *Revista Brasileira de Zoologia* 57:583–594
- Putman RJ (1996) *Community Ecology*. Chapman & Hall, London
- Richardson BA (1999) The bromeliad microcosm and the assessment of faunal diversity in a Neotropical Forest. *Biotropica* 31:321–336
- Rizzini CT (1979) *Tratado de fitogeografia do Brasil. Aspectos sociológicos e florísticos*. HUCITEC-EDUSP, São Paulo
- Rossa-Feres DC, Menin M, Izzo TJ (1999) Ocorrência sazonal e comportamento territorial em *Leptodactylus fuscus* (Anura, Leptodactylidae). *Ilheringia. Série Zoológica* 87:93–100

- Santos TG, Rossa-Feres DC, Casatti L (2007) Diversidade e distribuição espaço-temporal de anuros em região com pronunciada estação seca no sudeste do Brasil. *Iheringia, Série Zoológica* 97:37–49
- Schneider JAP, Teixeira RL (2001) Relacionamento entre anfíbios anuros e bromélias da restinga de Regência, Linhares, Espírito Santo, Brasil. *Iheringia, Série Zoológica* 91:41–48
- Sokal RR, Oden NL (1978) Spatial autocorrelation in biology. I Methodology. *Biological Journal Linnean Society* 10:199–228
- Sredl MJ, Collins JP (1992) The interaction of predation, competition, and habitat complexity in structuring an amphibian community. *Copeia* 18:607–614
- Suguio K, Tessler MG (1984) Planícies de cordões litorâneos quaternários do Brasil: Origem e nomenclatura. In: Lacerda LD, Araujo DSD, Cerqueira R, Turcq B (Eds) *Restingas Origem, Estrutura e Processos*. CEUFF, Niterói
- Tews J, Brose U, Grimm V, Tielbörger K, Tielbörger K, Wichmann MC, Schwagerl M, Jeltsch F. (2004) Animal species diversity driven by habitat heterogeneity/diversity: the importance of keystone structures. *Journal of Biogeography* 31:79–92
- Teixeira RL, Schneider JAP, Almeida GI (2002) The occurrence of amphibians in bromeliads from a southeastern Brazilian Restinga habitat, with special reference to *Aparasphenodon bruno*i (Anura, Hylidae). *Brazilian Journal of Biology* 62:263–268
- Underwood AJ, Chapman MG, Connell SD (2000) Observations in ecology: you can't make progress on processes without understanding the patterns. *Journal of Experimental Marine Biology and Ecology* 250:97–115

- Uetanabaro M, Souza FL, Filho PL, Beda AF, Brandão RA (2007) Anfíbios e répteis do Parque Nacional da Serra da Bodoquena, Mato Grosso do Sul, Brasil. *Biota Neotropica* 7:279–289
- Vallan D (2002) Effects of anthropogenic environmental changes on amphibian diversity in the rain forests of eastern Madagascar. *Journal of Tropical Ecology* 18:725–742
- Vanzolini PH (1981) A quasi-historical approach to the natural history of the differentiation of reptiles in tropical geographic isolates. *Papéis Avulsos da Zoologia* 34: 189-204
- Vasconcelos TS, Rossa-Feres DC (2005) Diversidade, distribuição espacial e temporal de anfíbios anuros (Amphibia, Anura) na região noroeste do estado de São Paulo, Brasil. *Biota Neotropica* 5:1–14
- Velloso AL, Sampaio EVB, Pareyn FGC (2002) Ecorregiões propostas para o Bioma Caatinga. Associação Plantas do Nordeste, Instituto de Conservação Ambiental The Nature Conservancy do Brasil, Recife
- Vieira WLS, Arzabe C, Santana GG (2007) Composição e distribuição espaço-temporal de anuros no Cariri Paraibano, Nordeste do Brasil. *Oecologia Brasileira* 11:383–396
- Zimmerman BL, Bierregaard RO (1986) Relevance of the equilibrium theory of island biogeography and species-area relations to conservation with a case from Amazônia. *Journal of Biogeography* 13:133–143
- Zina J, Enns J, Pinheiro SCP, Haddad CB, Toledo LF (2007) Taxocenose de anuros de uma mata semidecídua do interior do Estado de São Paulo e comparações com outras taxocenoses do Estado, Sudeste do Brasil. *Biota Neotropica* 7:1–9



## Figure Legends

**Fig. 1** Distribution map of the 20 sample units within the limits of the Sete Passagens State Park (PESP), Municipality of Miguel Calmon, State of Bahia, Brazil. **(a)** South America; in gray, Brazil; in black, Bahia State; white circle, geographical position of the PESP headquarters. **(b)** Topographic map of the State of Bahia, highlighting PESP's position in the Ecoregion of the Chapada Diamantina Complex. **(c)** PESP's polygon, highlighting its vegetation, hydrography, sample units (circles), and State Park headquarters (square). The Capão Forest is a forest patch inside an open area due to natural causes or to human activities, a name commonly used for this type of physiognomy by native inhabitants of the studied area

**Fig. 2** Distribution map of 10 environmental sampling points chosen at random within a hypothetical sampling unit. Five environmental variables were taken following this map: soil moisture, percentage of leaf litter, canopy height, density of trees, and percentage of understory, shrub, and herb layers

**Fig. 3** Composite graph of the anuran community structure from the Sete Passagens State Park based on weighted averaging from the NMS ordination. X-axis, sample units; Y-axis, relative abundances (in percentage) of an anuran species per sample unit (matrix C). Arabic numerals represent environmental categories: 1, Campo Rupestre with water bodies on the ground; 2, Campo Rupestre without water bodies on the ground; 3, hybrid habitat (mix of forest patch with streams and open areas of Campo Rupestre); 4, forest patches without water bodies on the ground, but of elevated air moisture and high density of epiphytes and herbaceous vegetation; 5,

dense forests with water bodies on the ground; 6, dense forests without water bodies on the ground. See text for a more detailed description of each category

**Fig. 4 (a)** Projection of individual scores resulting from the principal component analysis for 13 environmental variables of the combined 16 sample units (SUs) used in the account in the space of the first with the second axis. **(b)** Partial regression plot of the first principal component axis (PC1; environmental variables) on the nonmetric multidimensional scaling axis (NMS; anuran species composition). Regression was significant ( $P < 0.0004$ , Bonferroni corrected). Symbols represent types of environment: open and closed symbols, SUs without and with water bodies on the ground, respectively; asterisk, hybrid habitat (see legend of Fig. 3); circle, Campo Rupestre; square, dense forest; triangle, forest patch with many epiphytic bromeliads, herbaceous vegetation, and bryophytes

**Fig. 5** Anurans from the Sete Passagens State Park. **(a)** *Bokermannohyla* cf. *circumdata*; **(b)** *B. oxente*; **(c)** *Hypsiboas crepitans*; **(d)** *H. faber*; **(e)** *Scinax eurydice*; **(f)** *S. aff. catharinae*; **(g)** *Scinax* sp.; **(h)** *S. pachycrus*; **(i)** *Phyllomedusa bahiana*; **(j)** *Trachycephalus aff. mesophaeus*; **(l)** *Strabomantis aramunha*; **(m)** *Ischnocnema aff. ramagii*; **(n)** *Ischnocnema* sp.; **(o)** *Proceratophrys* sp.; **(p)** *Rhinella crucifer*

**Table 1** Abundances of anuran species in the Sete Passagens State Park (PESP), and within its surrounding area (xeric plant cover). Prevailing physiognomies of the PESP are ordered following PC1 ordination scores of 16 sample units used in the account, and species based on weighted averaging from NMS ordination (except for *Ischnocnema aff. ramagii*, which was not included in the analysis). Abundances from outside limits of the PESP are not given due to distinct sample efforts; species unique to this are alphabetically ordered. A hybrid habitat is a

mix of forest patch with streams and open areas of Campo Rupestre. CAA: Caatinga plant cover; DAM, human-made barrier within a stream, the latter characterized by slow backwaters that almost seem like a lentic system; LEN, lentic environment; LFP, lowland forest patches outside the PESP limits; LOT, lotic environment, characterized by springs and streams of noticeably flowing waters; WWA, areas without water bodies on the ground. Acronyms in parentheses indicate anuran families: BR, Brachycephalidae; BU, Bufonidae; CY, Cycloramphidae; HY, Hylidae; LE, Leptodactylidae; LP, Leiuperidae; ST, Strabomantidae

Species (Family)	Surrounding		Sete Passagens State Park						
	area		Campo Rupestre		Hybrid habitat		Semideciduous Forest patches		
	CAA	LFP	LEN	WWA	LOT	DAM	WWA	LOT	WWA
Sample units	—	—	P20	P10, P16–18	P6	P13	P3–4	P1, P5, P11, P14–15	P2, P12
<i>Ischnocnema</i> aff. <i>ramagii</i> (BR)			33	69	6	8	65	87	19
<i>Corythomantis grenningi</i> (HY)		•							
<i>Dendropsophus branneri</i> (HY)	•								
<i>Dendropsophus minutus</i> (HY)	•								
<i>Dendropsophus oliveirai</i> (HY)	•								
<i>Leptodactylus fuscus</i> (LE)	•								
<i>Leptodactylus ocellatus</i> (LE)	•								
<i>Leptodactylus vastus</i> (LE)	•								
<i>Odontophrynus</i> sp. (CY)	•								
<i>Phyllomedusa nordestina</i> (HY)	•								
<i>Physalaemus</i> aff. <i>kroyeri</i> (LP)	•								
<i>Procerathophrys</i> sp. (CY)	•								
<i>Pseudopaludicola</i> sp. (LP)	•								
<i>Rhinella granulosa</i> (BU)	•								

<i>Rhinella jimi</i> (BU)	•									
<i>Scinax pachycrus</i> (HY)	•			2						
<i>Ischnocnema</i> sp. (BR)			12	52	20	1	9	12		
<i>Phyllomedusa bahiana</i> (HY)	•		2			2		1		
<i>Scinax</i> sp. (HY)			5	10	5	15		6		
<i>Strabomantis aramunha</i> (BR)							2			
<i>Hypsiboas crepitans</i> (HY)	•					1				
<i>Hypsiboas faber</i> (HY)	•		1	1				1		
<i>Proceratophrys</i> aff. <i>cristiceps</i> (CY)					7	1	1	15	1	
<i>Trachycephalus</i> aff. <i>mesophaeus</i> (HY)							1	3		
<i>Bokermannohyla oxente</i> (HY)	•		1	5	6	7	5	49	10	
<i>Scinax eurydice</i> (HY)							1		3	
<i>Scinax</i> aff. <i>catharinae</i> (HY)								2		
<i>Bokermannohyla</i> cf. <i>circumdata</i> (HY)								1		
<i>Rhinella crucifer</i> (BU)								1	1	
<b>Total number of species</b>		17	2	6	6	5	7	7	11	5

**Table 2** Loadings from Principal Components Analysis (after Varimax rotation) for 13 environmental variables of 16 combined sample units from the Sete Passagens State Park, Municipality of Miguel Calmon, State of Bahia, Brazil. Projection of component scores are in Fig. 4a

Environmental variables	PC1	PC2	PC3	PC4
Air temperature	0.30	0.23	−0.83	−0.06
Air moisture	0.31	−0.06	0.81	0.29
Number of epiphytic bromeliads	0.25	0.41	0.70	−0.14
Number of terrestrial bromeliads	−0.05	−0.14	0.15	0.89
Canopy height	0.80	0.22	0.14	−0.08
Density of trees	0.82	0.09	−0.28	0.01
Maximum width of the water body	0.20	0.88	0.03	−0.07
Maximum depth of the water body	−0.10	0.86	−0.25	0.16
Soil moisture	0.68	0.57	0.37	0.03
Type of water body in the SU	0.28	0.77	0.30	−0.28
Percentage of understory, shrub, and herb layers	0.65	0.21	0.01	0.67
Estimated percentage of water cover in the SU	0.25	0.84	−0.13	−0.01
Percentage of leaf litter	0.93	0.08	0.12	0.06
Eigenvalues	5.21	2.54	1.93	1.25
% Total variance	40.0	19.51	14.88	9.61






FIG. 1




FIG. 2






FIG. 3




FIG. 4



FIG. 5

### III CONCLUSÃO GERAL

- Os anuros presentes no PESP refletem biogeografia complexa, apresentando espécies associadas a diferentes fitofisionomias, como caatinga, campo rupestre e floresta semi-decidual.
- A ocorrência de *Bokermannohyla* aff. *circumdata* na Serra de Jacobina agrega importante valor biogeográfico e conservacional, já que a presença deste táxon representa um relictos da distribuição original da espécie no domínio da Floresta Atlântica e o primeiro registro da espécie para o semi-árido.
- A comunidade de anfíbios anuros da área estudada revelou uma ordenação forte, apresentando uma associação de 82% com as variáveis relacionadas à estrutura da vegetação, sendo estas, em ordem crescente de contribuição: porcentagem da réplica coberta por serapilheira, altura do dossel e densidade de lenhosas. Evidenciou-se um gradiente relacionado primariamente à estrutura da vegetação: ambientes abertos – de transição – florestados. Em nosso estudo, áreas florestadas com vegetação densa (fechada), além de possuírem maior riqueza, abrigaram espécies cujas áreas de ocorrência estão relacionadas a formações do domínio Tropical Atlântico, enquanto que ambientes arbustivos como os de campo rupestre, estão geralmente relacionados a espécies generalistas e de ampla distribuição. Neste contexto, é necessário considerar durante a elaboração e execução das ações de manejo e uso do PESP, a relevância das áreas florestadas para a manutenção da sua anurofauna.

- O padrão de distribuição das espécies observado não reflete associação entre a posição geográfica das réplicas e a ordenação indireta da comunidade de anuros do PESP (eixo NMDS).
- A singularidade das espécies encontradas no PESP (matas estacionais) é percebida quando comparadas com aquelas registradas na área de entorno com formação vegetal de caatinga, havendo clara distinção entre as respectivas riqueza e composição de espécies de anuros. De interesse particular à conservação das espécies de anfíbios da Serra de Jacobina, destacamos que a manutenção das zonas protegidas pela poligonal do PESP protege apenas parte das espécies de anuros da região. As espécies que habitam áreas menos elevadas e circunjacentes ao PESP não estão sob o mesmo grau de proteção. Neste escopo, a área de entorno deve ter tratamento condizente com sua importância faunística, implicando em restrições severas a procedimentos que sejam poluidores ou que descaracterizem sua cobertura vegetal original.
- Existe uma tendência no Brasil em concentrar Unidades de Conservação em áreas montanhosas, o que é coincidente para UCs federais, estaduais e municipais de proteção integral no Estado da Bahia (MMA 2006). As regiões de baixadas, onde ocorrem extensas áreas de brejos e uma fauna com diferenças marcantes em relação às áreas de altitude, devem ser também priorizados em planejamentos para a estruturação de novas Unidades de Conservação (UCs).

## IV REFERÊNCIAS BIBLIOGRÁFICAS

- Ab'Sáber AN (1977) Os domínios morfoclimáticos na América do Sul. *Geomorfologia* 52:1–22
- Bastazini CV, Munduruca JFV, Rocha PLB, Napoli MF (2007) Which environmental variables better explain changes in anuran community composition? A case study in the restinga of Mata de São João, Bahia, Brazil. *Herpetologica* 63:459–471
- Beebee TJC, Griffiths RA (2005) The amphibian decline crisis. A watershed for conservation biology? *Biological Conservation* 25:271–285
- Begon M, Harper JL, Townsend CR (2006) *Ecology: Individuals, Populations, and Communities*. Blackwell Scientific Publications, Oxford
- Bernarde PS, Kokubum, MNC (1999) Anurofauna do município de Guararapes, estado de São Paulo, Brasil (Amphibia, Anura). *Acta biológica leopoldensia* 21:89–97
- Bernarde PS (2007) Ambientes e temporada de vocalização da anurofauna no Município de Espigão do Oeste, Rondônia, Sudoeste da Amazônia - Brasil (Amphibia: Anura). *Biotaneotropica* 7:87–92.  
<http://www.biotaneotropica.org.br/v7n2/pt/abstract?article+bn01507022007>,  
acessado em 15 de maio de 2008
- Bertoluci JA (1998) Annual patterns of breeding activity in Atlantic Rainforest anurans. *Journal of Herpetology* 32:607–611
- Connell JH (1980) Diversity and the coevolution of competitors, or the ghost of competition past. *Oikos* 35:131–138

- Conte EC, Machado RA (2005) Riqueza de espécies e distribuição espacial e temporal em comunidade de anuros (Amphibia, Anura) em uma localidade de Tijucas do sul, Paraná, Brasil. *Revista Brasileira de Zoologia* 22:940–948
- Cardoso AJ, Andrade GV, Haddad CFB (1989) Distribuição espacial em comunidades de anfíbios (Anura) no sudeste do Brasil. *Revista Brasileira de Biologia* 49:241–249
- Cascon P (1987) Observações sobre diversidade, ecologia e reprodução na anurofauna de uma área de caatinga. Dissertação de Mestrado, Universidade Federal da Paraíba, João Pessoa
- Crump ML, Hensley FR, Clark KL (1992) Apparent decline of the golden toad: underground or extinct? *Copeia* 1992:413–420
- Duellman WE (1988) Patterns of species diversity in anuran amphibians in the American tropics. *Annals of the Missouri Botanical Garden* 75:79–104
- Duellman WE (1999) Distribution Patterns of Amphibians in South America. In: Duellman WE (ed) *Patterns of Distribution of Amphibians*. The Johns Hopkins University Press, Baltimore
- Duellman WE, Trueb L (1994) *Biology of Amphibians*. The Johns Hopkins University Press, Baltimore, Maryland
- Eterovick PC (2003) Distribution of anuran species among montane streams in south-eastern Brazil. *Journal of Tropical Ecology* 19:219–228
- Feio RN, Ferreira PL (2005) Anfíbios de dois fragmentos de Mata Atlântica no município de Rio Novo, Minas Gerais. *Revista Brasileira de Zoociências* 7:121–128

- Funk WC, Mills LS (2003) Potential causes of population declines in Forest fragments in an Amazonian frog. *Biological Conservation* 111:205–214
- Gascon C (1991) Population and community-level analysis of species occurrences of central Amazonian rain forest tadpoles. *Ecology* 72:1731–1746
- Homes RT, Sherry TW, Sturges FW (1986) Bird community dynamics in a temperate deciduous forest: long-term trends at Hubbard Brook. *Ecological Monographs* 56:201–220
- Inger RF (1969) Organization of Communities of Frogs Along Small Rain Forest Streams in Sarawak. *The Journal of Animal Ecology* 38:123–148
- Krishnamurthy SV (2003) Amphibian assemblages in undisturbed and disturbed areas of Kudremukh National Park, Central Western Ghats, India. *Environmental Conservation* 30:274–282
- Lewinsonhn TM, Prado PI (2002) Biodiversidade brasileira: síntese do estado atual do conhecimento. Editora Contexto, São Paulo
- Lima DC (1999) Aspectos Sistemáticos, Zoogeográficos e Ecológicos da Herpetofauna da Serra de Maranguape, Estado do Ceará. Monografia de Graduação, Universidade Federal do Ceará, Fortaleza
- MacArthur RH, MacArthur JW (1961) On bird species diversity. *Ecology* 42:594–598
- Marsh DM (2001) Fluctuations in amphibian populations: a meta-analysis. *Biology Conservation* 101:327–335
- Mesquita DO, Costa GC, Colli GR (2006) Ecology of an Amazonian savanna lizard assemblage in Monte Alegre, Pará State, Brazil. *South American Journal of Herpetology* 1:61–71



- MMA – Ministério do Meio Ambiente (2007) Áreas Prioritárias para Conservação, Uso Sustentável e Repartição de Benefícios da Biodiversidade Brasileira: Atualização - Portaria MMA nº9, de 23 de janeiro de 2007. Ministério do Meio Ambiente, Secretaria de Biodiversidade e Florestas, Série Biodiversidade 31, Brasília
- Naniwadekar R, Vasudevan K (2007) Patterns in diversity of anurans along an elevational gradient in the Western Ghats, South India. *Journal of Biogeography* 34:842–853
- Parris KM, McCarthy M (1999) What influences the structure of frog assemblages at forest streams? *Australian Journal of Ecology* 24:495–502
- Parris KM (2004) Environmental and spatial variables influence the composition of frog assemblages in sub-tropical eastern Australia. *Ecography* 27:392–400
- Pianka ER (1973) The structure of lizard communities. *Annual Review of Ecology and Systematics* 4:53–74
- Pombal Jr JP (1997) Distribuição espacial e temporal de anuros (Amphibia) em uma poça permanente na Serra de Paranapiacaba, Sudeste do Brasil. *Revista Brasileira de Zoologia* 57:583–594
- Putman RJ (1996) *Community Ecology*. Chapman & Hall, London
- Ricklefs RE (2001) *The economy of nature*. Freeman and Company, New York
- Rodrigues MT (2003) Herpetofauna da Caatinga. In: Leal IR, Tabarelli M, Silva JMC (eds) *Ecologia e Conservação da Caatinga*. Editora Universitária da Universidade Federal de Pernambuco, Pernambuco
- Rodrigues MT, Carvalho CM, Borges-Nojosa DM, Freire EMX, Curcio FF, Oliveira FF, Silva HR, Dixo MBO (2004) Anfíbios e répteis: áreas e ações prioritárias para

- conservação da caatinga. Pp. 182–188. In Silva JMC, Tabarelli M., Fonseca MT, Lins LV (Orgs) Biodiversidade da Caatinga: áreas e ações prioritárias para conservação. Ministério do Meio Ambiente, Brasília
- Rossa-Feres DC, Jim J (1994) Distribuição sazonal em comunidades de anfíbios anuros na região de Botucatu, São Paulo. *Revista Brasileira de Biologia* 54:323–334
- Rossa-Feres DC, Menin M, Izzo TJ (1999) Ocorrência sazonal e comportamento territorial em *Leptodactylus fuscus* (Anura, Leptodactylidae). *Ilheringia, Série Zoológica* 87:93–100
- Santos TG, Rossa-Feres DC, Casatti L (2007) Diversidade e distribuição espaço-temporal de anuros em região com pronunciada estação seca no sudeste do Brasil. *Ilheringia, Série Zoológica* 97:37–49
- Sredl MJ, Collins JP (1992) The interaction of predation, competition, and habitat complexity in structuring an amphibian community. *Copeia* 18:607–614
- Tews J, Brose U, Grimm V, Tielbörger K, Tielbörger K, Wichmann MC, Schwagerl M, Jeltsch F. (2004) Animal species diversity driven by habitat heterogeneity/diversity: the importance of keystone structures. *Journal of Biogeography* 31:79–92
- Underwood AJ, Chapman MG, Connell SD (2000) Observations in ecology: you can't make progress on processes without understanding the patterns. *Journal of Experimental Marine Biology and Ecology* 250:97–115
- Uetanabaro M, Souza FL, Filho PL, Beda AF, Brandão RA (2007) Anfíbios e répteis do Parque Nacional da Serra da Bodoquena, Mato Grosso do Sul, Brasil. *Biota Neotropica* 7:279–289

- Vallan D (2002) Effects of anthropogenic environmental changes on amphibian diversity in the rain forests of eastern Madagascar. *Journal of Tropical Ecology* 18:725–742
- Velloso AL, Sampaio EVB, Pareyn FGC (2002) Ecorregiões propostas para o Bioma Caatinga. Associação Plantas do Nordeste, Instituto de Conservação Ambiental The Nature Conservancy do Brasil, Recife
- Vieira WLS, Arzabe C, Santana GG (2007) Composição e distribuição espaço-temporal de anuros no Cariri Paraibano, Nordeste do Brasil. *Oecologia Brasileira* 11:383–396
- Vitt LJ, Caldwell JP, Wilbur HM, Smith DC (1990) Amphibians as harbingers of decay. *BioScience* 40:418–418
- Wake DB (1991) Declining amphibian populations. *Science* 253:860–802
- Willson MF (1974) Avian community organization and habitat structure. *Ecology* 55:1017–1029
- Zina J, Ennser J, Pinheiro SCP, Haddad CB, Toledo LF (2007) Taxocenose de anuros de uma mata semidecídua do interior do Estado de São Paulo e comparações com outras taxocenoses do Estado, Sudeste do Brasil. *Biota Neotropica* 7:1–9

## V ANEXO

Normas para submissão de artigos a serem publicados no periódico Biodiversity and Conservation, publicado pela Springer, cujo conceito Qualis/CAPES em 2008 é A nas áreas de Ecologia e Conservação (disponível em: <http://www.springer.com/life+sci/journal/10531?detailsPage=contentItemPage&CIPageCounter=139710>; capturado em 26 de maio de 2008).

## Online Manuscript Submission

Springer now offers authors, editors and reviewers of *Biodiversity & Conservation* the option of using our fully web-enabled online manuscript submission and review system. To keep the review time as short as possible (no postal delays!), we encourage authors to submit manuscripts online to the journal's editorial office. Our online manuscript submission and review system offers authors the option to track the progress of the review process of manuscripts in real time. Manuscripts should be submitted to: <http://bioc.edmgr.com>

The online manuscript submission and review system for *Biodiversity & Conservation* offers easy and straightforward log-in and submission procedures. This system supports a wide range of submission file formats: for manuscripts - Word, WordPerfect, RTF, TXT and LaTeX; for figures - TIFF, GIF, JPEG, EPS, PPT, and Postscript.

**NOTE:** By using the online manuscript submission and review system, it is NOT necessary to submit the manuscript also in printout + disk. In case you encounter any difficulties while submitting your manuscript on line, please get in touch with the responsible Editorial Assistant by clicking on "CONTACT US" from the tool bar. The journal also publishes Editorials, Comments and Research notes. These types of articles should be submitted to the Journals Editorial Office in the usual way, but authors should clearly indicate that they are Editorials, Comments or Research notes.

## Electronic figures

Electronic versions of your figures must be supplied. For vector graphics, EPS is the preferred format. For bitmapped graphics, TIFF is the preferred format. The following resolutions are optimal: line figures - 600 - 1200 dpi; photographs - 300 dpi; screen dumps - leave as is. Colour figures can be submitted in the RGB colour system. Font-related problems can be avoided by using standard fonts such as Times Roman, Courier and Helvetica.

## Colour figures

Springer offers two options for reproducing colour illustrations in your article. Please let us know what you prefer: 1) Free online colour. The colour figure will only appear in colour on [www.springer.com](http://www.springer.com) and not in the printed version of the journal. 2) Online and printed colour. The colour figures will appear in colour on our website and in the printed version of the journal. The charges are EUR 950/USD 1150 per article.

## **Language**

We appreciate any efforts that you make to ensure that the language is corrected before submission. This will greatly improve the legibility of your paper if English is not your first language.

## **Manuscript Presentation**

The journal's language is English. British English or American English spelling and terminology may be used, but either one should be followed consistently throughout the article. Leave adequate margins on all sides to allow reviewers' remarks. Please double-space all material, including notes and references. Quotations of more than 40 words should be set off clearly, either by indenting the left-hand margin or by using a smaller typeface. Use double quotation marks for direct quotations and single quotation marks

for quotations within quotations and for words or phrases used in a special sense.

Number the pages consecutively with the first page containing:

running head (shortened title)

title

author(s)

affiliation(s)

full address for correspondence, including telephone and fax number and e-mail address

## **Abstract**

Please provide a short abstract of 100 to 250 words. The abstract should not contain any undefined abbreviations or unspecified references.

## **Key words**

Please provide 5 to 10 key words or short phrases in alphabetical order.

## **Abbreviations**

Abbreviations and their explanations should be collected in a list.

## **Symbols and units**

Please use the recommended SI units.

## **Nomenclature**

The correct names of organisms conforming with the international rules of nomenclature must be used. Descriptions of new taxa should not be submitted unless a specimen has been deposited in a recognized collection and it is designated as a type strain in the paper. Biodiversity and Conservation uses the same conventions for the genetics nomenclature of bacteria, viruses, transposable elements, plasmids and restriction enzymes as the American Society for Microbiology journals.

## **Figures**

All photographs, graphs and diagrams should be referred to as a 'Figure' and they should be numbered consecutively (1, 2, etc.). Multi-part figures ought to be labelled with lower case letters (a, b, etc.). Please insert keys and scale bars directly in the figures. Relatively small text and great variation in text sizes within figures should be avoided as figures are often reduced in size. Figures may be sized to fit approximately within the column(s) of the journal. Provide a detailed legend (without abbreviations) to

each figure, refer to the figure in the text and note its approximate location in the margin. Please place the legends in the manuscript after the references.

## **Tables**

Each table should be numbered consecutively (1, 2, etc.). In tables, footnotes are preferable to long explanatory material in either the heading or body of the table. Such explanatory footnotes, identified by superscript letters, should be placed immediately below the table. Please provide a caption (without abbreviations) to each table, refer to the table in the text and note its approximate location in the margin. Finally, please place the tables after the figure legends in the manuscript.

## **Section headings**

First-, second-, third-, and fourth-order headings should be clearly distinguishable but not numbered.

## **Appendices**

Supplementary material should be collected in an Appendix and placed before the Notes and Reference sections.



## Notes

Please use endnotes rather than footnotes. Notes should be indicated by consecutive superscript numbers in the text and listed at the end of the article before the References. A source reference note should be indicated by means of an asterisk after the title. This note should be placed at the bottom of the first page.

## Cross-referencing

In the text, a reference identified by means of an author's name should be followed by the date of the reference in parentheses and page number(s) where appropriate. When there are more than two authors, only the first author's name should be mentioned, followed by 'et al.'. In the event that an author cited has had two or more works published during the same year, the reference, both in the text and in the reference list, should be identified by a lower case letter like 'a' and 'b' after the date to distinguish the works.

*Examples:*

Winograd (1986, p. 204)

(Winograd 1986a, b)

(Winograd 1986; Flores et al. 1988)

(Bullen and Bennett 1990)

## Acknowledgements

Acknowledgements of people, grants, funds, etc. should be placed in a separate section before the References.

## References

### 1. Journal article:

Smith J, Jones M Jr, Houghton L et al (1999) Future of health insurance. *N Engl J Med* 341:325–329

### 2. Inclusion of issue number (optional):

Saunders DS (1976) The biological clock of insects. *Sci Am* 234(2):114–121

### 3. Journal issue with issue editor:

Smith J (ed) (1998) Rodent genes. *Mod Genomics J* 14(6):126–233

### 4. Journal issue with no issue editor:

*Mod Genomics J* (1998) Rodent genes. *Mod Genomics J* 14(6):126–233

### 5. Book chapter:

Brown B, Aaron M (2001) The politics of nature. In: Smith J (ed) *The rise of modern genomics*, 3rd edn. Wiley, New York

### 6. Book, authored:

South J, Blass B (2001) *The future of modern genomics*. Blackwell, London

### 7. Book, edited:

Smith J, Brown B (eds) (2001) *The demise of modern genomics*. Blackwell, London

### 8. Chapter in a book in a series without volume titles:

Schmidt H (1989) Testing results. In: Hutzinger O (ed) Handbook of environmental chemistry, vol 2E. Springer, Berlin Heidelberg New York, p 111

9. Chapter in a book in a series with volume title:

Smith SE (1976) Neuromuscular blocking drugs in man. In: Zaimis E (ed) Neuromuscular junction. Handbook of experimental pharmacology, vol 42. Springer, Berlin Heidelberg New York, pp593–660

10. Proceedings as a book (in a series and subseries):

Zowghi D et al (1996) A framework for reasoning about requirements in evolution. In: Foo N, Goebel R (eds) PRICAI'96: topics in artificial intelligence. 4th Pacific Rim conference on artificial intelligence, Cairns, August 1996. Lecture notes in computer science (Lecture notes in artificial intelligence), vol 1114. Springer, Berlin Heidelberg New York, p 157

11. Proceedings with an editor (without a publisher):

Aaron M (1999) The future of genomics. In: Williams H (ed) Proceedings of the genomic researchers, Boston, 1999

12. Proceedings without an editor (without a publisher):

Chung S-T, Morris RL (1978) Isolation and characterization of plasmid deoxyribonucleic acid from *Streptomyces fradiae*. In: Abstracts of the 3rd international symposium on the genetics of industrial microorganisms, University of Wisconsin, Madison, 4–9 June 1978

13. Paper presented at a conference:

Chung S-T, Morris RL (1978) Isolation and characterization of plasmid deoxyribonucleic acid from *Streptomyces fradiae*. Paper presented at the 3rd international symposium on the genetics of industrial microorganisms, University of Wisconsin, Madison, 4–9 June 1978

## 14. Patent:

Name and date of patent are optional

Norman LO (1998) Lightning rods. US Patent 4,379,752, 9 Sept 1998

## 15. Dissertation:

Trent JW (1975) Experimental acute renal failure. Dissertation, University of California

## 16. Institutional author (book):

International Anatomical Nomenclature Committee (1966) *Nomina anatomica. Excerpta Medica*, Amsterdam

## 17. Non-English publication cited in an English publication:

Wolf GH, Lehman P-F (1976) *Atlas der Anatomie*, vol 4/3, 4th edn. Fischer, Berlin.  
[NB: Use the language of the primary document, not that of the reference for "vol" etc.!]

## 18. Non-Latin alphabet publication:

The English translation is optional.

Marikhin VY, Myasnikova LP (1977) *Nadmolekulyarnaya struktura polimerov* (The supramolecular structure of polymers). Khimiya, Leningrad

## 19. Published and In press articles with or without DOI:

## 19.1 In press

Wilson M et al (2006) References. In: Wilson M (ed) *Style manual*. Springer, Berlin Heidelberg New York (in press)

## 19.2. Article by DOI (with page numbers)

Slifka MK, Whitton JL (2000) Clinical implications of dysregulated cytokine production. *J Mol Med* 78:74–80. DOI 10.1007/s001090000086

#### 19.3. Article by DOI (before issue publication with page numbers)

Slifka MK, Whitton JL (2000) Clinical implications of dysregulated cytokine production. *J Mol Med* (in press). DOI 10.1007/s001090000086

#### 19.4. Article in electronic journal by DOI (no paginated version)

Slifka MK, Whitton JL (2000) Clinical implications of dysregulated cytokine production. *Dig J Mol Med*. DOI 10.1007/s801090000086

### 20. Internet publication/Online document

Doe J (1999) Title of subordinate document. In: The dictionary of substances and their effects. Royal Society of Chemistry. Available via DIALOG. [http://www.rsc.org/dose/title of subordinate document](http://www.rsc.org/dose/title%20of%20subordinate%20document). Cited 15 Jan 1999

#### 20.1. Online database

Healthwise Knowledgebase (1998) US Pharmacopeia, Rockville. <http://www.healthwise.org>. Cited 21 Sept 1998

#### Supplementary material/private homepage

Doe J (2000) Title of supplementary material. <http://www.privatehomepage.com>. Cited 22 Feb 2000 University site

Doe J (1999) Title of preprint. <http://www.uni-heidelberg.de/mydata.html>. Cited 25 Dec 1999 FTP site

Doe J (1999) Trivial HTTP, RFC2169. <ftp://ftp.isi.edu/in-notes/rfc2169.txt>. Cited 12 Nov 1999

#### Organization site

ISSN International Centre (1999) Global ISSN database. <http://www.issn.org>. Cited 20 Feb 2000

## **Proofs**

Proofs will be sent to the corresponding author. One corrected proof, together with the original, edited manuscript, should be returned to the Publisher within three days of receipt by mail (airmail overseas).

## **Offprints**

Fifty offprints of each article will be provided free of charge. Additional offprints can be ordered by means of an offprint order form supplied with the proofs.

## **Page charges and colour figures**

No page charges are levied on authors or their institutions. Colour figures are published at the author's expense only.

## **Copyright**

Authors will be asked, upon acceptance of an article, to transfer copyright of the article to the Publisher. This will ensure the widest possible dissemination of information under copyright laws.

## **Permissions**

It is the responsibility of the author to obtain written permission for a quotation from unpublished material, or for all quotations in excess of 250 words in one extract or

500 words in total from any work still in copyright, and for the reprinting of figures, tables or poems from unpublished or copyrighted material.

### **Springer Open Choice**

In addition to the normal publication process (whereby an article is submitted to the journal and Access to that article is granted to customers who have purchased a subscription), Springer now provides an alternative publishing option: Springer Open Choice. A Springer Open Choice article receives all the benefits of a regular subscription-based article, but in addition is made available publicly through Springer's online platform SpringerLink. To publish via Springer Open Choice, upon acceptance please visit [www.springer.com/openchoice](http://www.springer.com/openchoice) to complete the relevant order form and provide the required payment information. Payment must be received in full before publication or articles will publish as regular subscription-model articles. We regret that Springer Open Choice cannot be ordered for published articles.

### **Additional information**

Additional information can be obtained from:

*Biodiversity and Conservation*

Springer

P.O. Box 17

3300 AA Dordrecht

The Netherlands

Fax: 78-6576254

Internet: <http://www.springer.com>

Página: 1

[AMP1]Rever!!! Todo o parágrafo!



# Livros Grátis

( <http://www.livrosgratis.com.br> )

Milhares de Livros para Download:

[Baixar livros de Administração](#)

[Baixar livros de Agronomia](#)

[Baixar livros de Arquitetura](#)

[Baixar livros de Artes](#)

[Baixar livros de Astronomia](#)

[Baixar livros de Biologia Geral](#)

[Baixar livros de Ciência da Computação](#)

[Baixar livros de Ciência da Informação](#)

[Baixar livros de Ciência Política](#)

[Baixar livros de Ciências da Saúde](#)

[Baixar livros de Comunicação](#)

[Baixar livros do Conselho Nacional de Educação - CNE](#)

[Baixar livros de Defesa civil](#)

[Baixar livros de Direito](#)

[Baixar livros de Direitos humanos](#)

[Baixar livros de Economia](#)

[Baixar livros de Economia Doméstica](#)

[Baixar livros de Educação](#)

[Baixar livros de Educação - Trânsito](#)

[Baixar livros de Educação Física](#)

[Baixar livros de Engenharia Aeroespacial](#)

[Baixar livros de Farmácia](#)

[Baixar livros de Filosofia](#)

[Baixar livros de Física](#)

[Baixar livros de Geociências](#)

[Baixar livros de Geografia](#)

[Baixar livros de História](#)

[Baixar livros de Línguas](#)

[Baixar livros de Literatura](#)  
[Baixar livros de Literatura de Cordel](#)  
[Baixar livros de Literatura Infantil](#)  
[Baixar livros de Matemática](#)  
[Baixar livros de Medicina](#)  
[Baixar livros de Medicina Veterinária](#)  
[Baixar livros de Meio Ambiente](#)  
[Baixar livros de Meteorologia](#)  
[Baixar Monografias e TCC](#)  
[Baixar livros Multidisciplinar](#)  
[Baixar livros de Música](#)  
[Baixar livros de Psicologia](#)  
[Baixar livros de Química](#)  
[Baixar livros de Saúde Coletiva](#)  
[Baixar livros de Serviço Social](#)  
[Baixar livros de Sociologia](#)  
[Baixar livros de Teologia](#)  
[Baixar livros de Trabalho](#)  
[Baixar livros de Turismo](#)