

**UNIVERSIDADE FEDERAL DE MINAS GERAIS
INSTITUTO DE CIÊNCIAS BIOLÓGICAS**

TESE DE DOUTORADO

Farmacogenética da esquizofrenia

Renan Pedra de Souza

Orientador: Dr. Marco A. Romano-Silva

BELO HORIZONTE – AGOSTO/ 2008

Livros Grátis

<http://www.livrosgratis.com.br>

Milhares de livros grátis para download.

Renan Pedra de Souza

Farmacogenética da esquizofrenia

Tese de Doutorado submetida ao curso de Pós-Graduação em Ciências Biológicas: Farmacologia Bioquímica e Molecular do Instituto de Ciências Biológicas da Universidade Federal de Minas Gerais como requisito parcial para a obtenção do título de Doutor em Farmacologia.

Orientador: Dr. Marco A. Romano-Silva

**Belo Horizonte
Minas Gerais - Brasil
2008**

“Médias são um conceito traiçoeiro. Representam um valor obtido a partir de resultados válidos para vários indivíduos, mas que não podem ser extrapolados a nenhum indivíduo em particular” (Hélio Schwartsman)

“Dados são muito parecidos com pessoas. Nascem, amadurecem, casam com outros dados, se divorciam e ficam velhos. Só não morrem, precisam ser mortos” (Arthur Miller)

“De tudo ficaram três coisas:

a certeza de que estamos sempre começando...

a certeza de que é preciso continuar...

a certeza de que seremos interrompidos antes de terminar..

fazer da interrupção um caminho novo...

da queda um passo de dança...

do medo, uma escada...

do sonho, uma ponte...

da procura...

um encontro" (Fernando Sabino)

Sumário

Agradecimentos V

Lista de abreviaturas..... VII

Resumo IX

Abstract X

1. Introdução

1.1 Esquizofrenia 02

1.2 Etiologia da esquizofrenia 05

1.3 Estudos genéticos em esquizofrenia 08

1.4 Tratamento da esquizofrenia 11

1.4.1 Discinesia tardia induzida por antipsicóticos 16

1.4.2 Ganho de peso induzido por antipsicóticos 19

2. Objetivos..... 20

3. Artigos

3.1 Association study of GSK3 β gene polymorphisms with schizophrenia and clozapine response.....	23
3.2 Genetic association analysis of the GFR alpha genes with schizophrenia and clozapine response.....	33
3.3 Clinical involvement of oxidative stress genes polymorphisms in schizophrenia: influence on the severity of symptoms and response to clozapine treatment.....	62
3.4 Genetic association study of NALCN polymorphisms with schizophrenia and antipsychotic treatment.....	74
3.5 Association of antipsychotic induced weight gain and body mass index with GNB3 gene: a meta-analysis	95
4. Conclusões.....	110
5. Referências bibliográficas.....	112

Agradecimentos

À minha mãe, Ternísia, mais uma vez, por todos sonhos abortados em prol de um ideal.

Espero que este resultado seja mais um marco de que não fora em vão tantas noites acordadas, lágrimas escondidas e sufocos vividos. Era só de tempo que precisávamos.

Ao Marco, o pai científico, que me fez acreditar novamente na competência de um referencial masculino. Embora tenha certeza de que muitos outros Renans passarão sobre sua orientação, não sei se outros Marcos cruzarão meu caminho. Muito obrigado por esses cinco anos e meio de convivência. Meu mundo cresceu muito nestes anos e fostes um dos grandes responsáveis por isto.

Ao meu pai, Jorge, pela existência e educação. Aprendi que não devo me frustrar por não receber algo de quem não pode dar. Que a vida te faça feliz, tal como tem me feito.

Ao James Kennedy e Albert Wong, pela oportunidade e receptividade.

Ao Marcus Gomez, pelo exemplo de pesquisador.

À Wolfanga Boson, Antoniana Krettli, Marco Prado, Vânia Prado e Luís Armando que, mesmo de maneira inconsciente, tiveram um papel importante na minha formação

Aos avós, Miguel, Ternísia, Augusto e Maria, pois sem vosso esforço esse castelo não teria tido um alicerce tão forte. Educação é realmente um bem fantástico.

Ao Yuri e Breno, meus irmãos, pelo convívio e aprendizagem. Se não puder ser o melhor exemplo, saibam que a intenção é sempre essa.

À Daniela, que só palavras não seriam suficientes para mensurar a alegria convívio. Muito obrigado pela companhia, em especial naqueles momentos que eu não enxergava saídas.

Sentirei a falta do cotidiano ao teu lado.

À Adriane Aparecida pelo sorriso e fazer da rotina algo tão agradável.

Aos amigos de fora do mundo científico, os quais a sanidade ainda permanece, embora em nuances cada vez mais claras, Ana Cláudia, André Luis, Cláudia Brant, Francisco Souza, Isabela Teixeira, Samira Fontes, Viviane Ribeiro e Walney Dias, que ao longo desses anos conseguiram lidar com a falta de tempo, paciência e presença.

Aos membros dos laboratórios no Brasil, presentes e passados, Alexandre Barros, Alessandra Souza, Ana Campi, Ana Cristina Pinheiro, Ana Cristina Magalhães, André Massensini, André Sampaio, Antônio Giovanni, Antônio Lúcio, Beatriz Soares, Bernardo Motta, Bruno Pinheiro, Bruno Rezende, Célio Castro, Christopher Kushmerick, Cristina Guatimosim, Cristina Martins, Daniel Carneiro, Danusa, Dawidson Gomes, Débora Miranda, Diele Barreto, Elizabete Miranda, Evanguedes Kalapothakis, Fabíola Ribeiro, Fabiana Caetano, Fabrício Lima, Felipe Filardi, Felipe Simil, Fernando Caetano, Flávio Pimenta, Gislene Valadares, Grace Schenatto, Guilherme Pimenta, Helton José, Humberto Correa, Janice Silva, Juliana Drummond, Juliana Garcia, Karen Torres, Kléber Pinto, Letícia Silveira, Livia Paulineli, Luciene Vieira, Lucimar Teodoro, Maria Aparecida, Melissa Guimarães, Melissa Vianna, Paola Perdigão, Patrícia, Paula Vidigal, Paulo Henrique, Paulo Melillo, Píndaro Massote, Raphael Penholati, Rodrigo Nicolato, Stefany Fontes, Victor Lima e Virgínia Fernandes.

Aos membros dos laboratórios no Canadá, presentes e passados, Alessio Squassina, Andrea Smart, Anjali Rastogi, Arun Tiwari, Bernard Le Foll, Clement Zai, Daniela Hloussek, Daniel Muller, Daniela Lobo, Emmanuela Viggiano, Fahmida Nipa-Haque, Joanne Brathwhite, Laura Feldcamp, Lori Dixon, Mahawib Semeralul, Maria Tampakeras, Mario Masselis, Mary Smirniw, Natalie Bulgin, Nicole King, Olga Likhodi, Periandrea Muglia, Poonam Batra, Quinton Van Adrichem, Rudi Hwang, Sahar Ehtesham, Sajid Shaikh, Subi Tharmaligam, Tricia Sicard, Vincenzo de Luca, Xingqun Ni e Yuko Hirata.

Aos parentes e amigos, a chegada até aqui também é uma conquista vossa.

Aos indivíduos que participaram deste estudo.

À vida.

Lista de abreviaturas

5-HT _{1A}	Receptor de serotonina tipo 1A
5-HT _{2A}	Receptor de serotonina tipo 2A
5-HT _{2C}	Receptor de serotonina tipo 2C
5-HT _{3A}	Receptor de serotonina tipo 3A
5-HT _{3B}	Receptor de serotonina tipo 3B
5-HT _{5A}	Receptor de serotonina tipo 5A
5-HT ₆	Receptor de serotonina tipo 6
5-HT ₇	Receptor de serotonina tipo 7
5-HTT	Transportador de serotonina
A	Adenina
A _{2A}	Receptor de adenosina tipo 2A
Akt/PKB	Proteína quinase B
BDNF	Fator neurotrófico derivado do cérebro
C	Citosina
CART	Transcrito regulado por cocaína e anfetamina
CATIE	<i>Clinical Antipsychotic Trial of Intervention Effectiveness</i>
ChAT	Colina acetiltransferase
COMT	Catecol-orto-metiltransferase
CUtLASS	<i>Cost Utility of the Latest Antipsychotics in Severe Schizophrenia</i>
CYP	Citocromo P450
DAO	D-aminoácido oxidase
DAOA/G72	Ativador da D-aminoácido oxidase
DARPP-32	Fosfoproteína regulada por dopamina e cAMP com 32 kilodaltons
DAT	Transportador de dopamina
DISC	<i>Disrupted in schizophrenia</i>
DRD1	Receptor de dopamina tipo 1
DRD2	Receptor de dopamina tipo 2
DRD3	Receptor de dopamina tipo 3
DRD4	Receptor de dopamina tipo 4
DSM-IV	Manual Diagnóstico e Estatístico das Doenças Mentais IV
DT	Discinesia tardia
DTNBP1	Disbindina 1
ECA	Enzima conversora de angiotensina
FDA	<i>Food and Drug Administration</i>
G	Guanina
GABA	Ácido gama-aminobutírico
GABA _A	Receptor de GABA do tipo A
GDNF	Fator neurotrófico derivado de glia
GNB-3	Subunidade β3 da proteína G
GPX	Glutationa peroxidase
GRIN2B	Receptor ionotrópico de glutamato NMDA subunidade 2B
GSK-3	Glicogênio sintase quinase 3
GST	Glutationa S transferase

H ₁	Receptor de histamina tipo 1
H ₂	Receptor de histamina tipo 2
His	Histidina
HLA	Antígeno de leucócito humano
MAO	Monoamino oxidase
MDR-1/ABCB1	Resistência múltipla à drogas 1
MnSOD	Mangânes superóxido dismutase
NALCN	<i>Sodium leak channel, non-selective</i>
NEF3	Neurofilamento 3
NET	Transportador de norepinefrina
NIMH	National Institute of Mental Health
NOS	Óxido nítrico sintase
NMDA	N-metil-D-aspartato
NOS	Óxido nítrico sintase
NQO1	NAD(P)H quinona oxiredutase
NRG-1	Neuregulina 1
PGP	Glicoproteína P
PRODH	Prolina desidrogenase
RGS	Proteína reguladora da proteína G
SNAP-25	Proteína sinaptossomal associada de 25 kilodaltons
SUS	Sistema Único de Saúde
T	Timina
TDAH	Transtorno de Deficit de Atenção e Hiperatividade
TH	Tirosina hidroxilase
Tir	Tirosina
TNF- α	Fator de necrose tumoral α
TPH	Triptofano hidroxilase

Resumo

A esquizofrenia é uma das doenças neuropsiquiátricas mais prevalentes, sendo caracterizada pela presença de delírios, alucinações, apatia e disfunção cognitiva, entre outros. Os medicamentos antipsicóticos foram introduzidos na década de 1950 e atualmente estão disponíveis vários antipsicóticos, com diferentes perfis de efeitos colaterais. Há evidências da presença de um componente genético na predisposição à esquizofrenia e à eficácia de resposta ao tratamento, bem como na apresentação de efeitos colaterais. Nesse trabalho foram analisados 9 genes (glicogênio sintase quinase 3 β – GSK-3 β ; 4 receptores do fator neurotrófico derivado de células gliais – GFRA1, GFRA2, GFRA3 e GFRA4; glutathione peroxidase – GPX1; mangânese superóxido dismutase – MnSOD; canal de sódio não-seletivo – NALCN e a subunidade β 3 da proteína G – GNB-3) buscando uma associação genética com a esquizofrenia, a resposta e apresentação de efeitos colaterais durante tratamento com antipsicóticos. Os resultados sugerem uma participação do gene da GSK-3 β associado à esquizofrenia; GFRA1, GFRA2 e GFRA3 na predisposição à esquizofrenia e a resposta à clozapina; NALCN contribuindo para a predisposição à esquizofrenia, resposta ao tratamento e apresentação de efeitos colaterais e o GNB-3 associado com menor índice de massa corpórea em uma meta-análise. Embora seja necessária a replicação dos resultados em amostras maiores e independentes para confirmação das associações encontradas, este trabalho apresenta dados relevantes e originais da farmacogenética da esquizofrenia.

Abstract

Schizophrenia is one of the most common neuropsychiatric disorder whose patients present disillusions, hallucinations, apathy and cognitive impairment. Antipsychotic drugs have been used since 1950. Currently, there are several drugs available that present different therapeutic efficacy and side-effects prevalence. It has been shown some evidences regarding a genetic susceptibility component to develop schizophrenia and response to antipsychotic treatment, as well as presents some side-effect to these drugs. In this study, it was analyzed association of 9 genes (glycogen syntase kinase 3β – GSK- 3β ; 4 glial-derived neurotrophic factor α receptors – GFR α 1 – 4; glutathione peroxidase – GPX1; manganese superoxide dismutase (MnSOD); sodium leak channel, non-selective (NALCN) and β 3 subunit G protein (GNB-3) with schizophrenia, clozapine treatment response, clozapine-induced weight gain and antipsychotic-induced tardive dyskinesia. Results suggest a role for: GSK- 3β in schizophrenia susceptibility; GFR α 1, GFR α 2 e GFR α 3 in schizophrenia susceptibility and treatment response; NALCN in schizophrenia susceptibility; treatment response and side-effects manifestation and GNB-3 in lower body mass index in a meta-analysis. Although it is necessary replication of our results in independent and larger samples in order to confirm our findings, this study presents original and relevant data on pharmacogenetics of schizophrenia.

1. Introdução

1.1 – Esquizofrenia

A esquizofrenia é um transtorno psicótico maior (ou um grupo de transtornos) que usualmente aparece na fase mais tardia da adolescência ou no início da idade adulta, sendo esta uma doença relativamente comum. Sua prevalência ao longo da vida é de 0,5-1% na população geral, mas essa estimativa pode variar de acordo com a metodologia utilizada nos diferentes estudos (Freedman, 2003). Os estudos epidemiológicos realizados no Brasil revelam estimativas de incidência e prevalência compatíveis com as observadas em outros países. Dados do Ministério de Saúde indicam que aproximadamente 1% da população brasileira já sofreu um episódio psicótico e estima-se que aproximadamente 80.000 pessoas são acometidas a cada ano. As taxas médias para homens e mulheres são similares, mas a idade média de início é cerca de cinco anos maior para as mulheres do que para os homens, sendo estes mais frequentemente afetados por sintomas negativos do que mulheres (Häfner, 2003). As mulheres tendem a apresentar um curso mais brando da esquizofrenia e, portanto, um melhor prognóstico e uma melhor possibilidade de adaptação social (Austin, 2005).

A importância da pesquisa sobre esquizofrenia é diretamente proporcional a sua importância social e econômica, em especial pelo fato que os indivíduos são geralmente acometidos no auge do seu potencial produtivo, entre 16 e 30 anos de idade, gerando uma sobrecarga para os pacientes e seus familiares. Na maioria dos casos, há prejuízo das funções ocupacionais ou sociais, caracterizado por afastamento social, perda de interesse ou capacidade de agir na escola ou no trabalho, mudança nos hábitos de higiene pessoal ou comportamento incomum (Loebel e cols, 1992; Häfner, 2003). Pacientes sofrem de um estresse considerável, tem sua qualidade de vida diminuída e enfrentam incapacidades prolongadas que podem impor efeitos negativos em seus empregos, nos orçamentos pessoal ou familiar, no relacionamento afetivo e na satisfação com a vida. Após passar pela fase aguda, o transtorno pode persistir, e períodos de remissão se alternam com os períodos de exacerbação (Austin, 2005).

Além de comprometer pacientes e familiares, há ainda um grande custo para toda a sociedade (tratamento e custos indiretos, como mortalidade e redução da produtividade). Após um segundo episódio psicótico, por exemplo, a taxa de desemprego entre pacientes com esquizofrenia é superior a 65%, contribuindo para o alto custo indireto da doença (Guest e Cookson, 1999). No Brasil, a esquizofrenia ocupa 30% dos leitos psiquiátricos hospitalares, ou

cerca de 100 mil leitos-dia. Ocupa ainda o segundo lugar das primeiras consultas psiquiátricas ambulatoriais (14%) e o quinto lugar na manutenção de auxílio-doença. Os gastos anuais do Sistema Único de Saúde com a internação e tratamento de 131 mil pacientes com esquizofrenia consomem quantias elevadas de recursos (Javitt e Coyle, 2004). Na Inglaterra, cerca de 6% dos custos com pacientes internados no Serviço Nacional de Saúde são relativos à esquizofrenia (Knapp, 1997), sendo esta responsável por 2,5% dos gastos com atendimento anual de saúde nos Estados Unidos (Rupp e Keith, 1993).

A esquizofrenia apresenta sintomas diferentes em múltiplos domínios, de forma muito heterogênea em diferentes indivíduos e também variabilidade nos mesmos indivíduos ao longo do tempo. Pela observação sistemática da psicopatologia, fenômenos positivos e negativos podem ser separados (Andreasen, 1982; Crow, 1985). Os sintomas positivos, de maneira abrangente, incluem delírios e ideação delirante, alucinações, distúrbios das associações, sintomas catatônicos, agitação, vivências de influência externa e desconfiança. Os sintomas negativos referem-se ao estreitamento e à redução das expressões emocionais, com diminuição da produtividade do pensamento e da fala, retraimento social e diminuição dos comportamentos direcionados a metas. Como uma terceira categoria, os sintomas desorganizados incluem desorganização do pensamento e do comportamento associado ao comprometimento da atenção.

A avaliação neuropsicológica longitudinal mostrou que os pacientes com esquizofrenia têm disfunção cognitiva considerável nos primeiros cinco anos da doença. Após esse período, há poucas evidências de deterioração (Hoff e cols, 1999). Aproximadamente metade dos pacientes com esquizofrenia tratados em serviços convencionais irão recidivar e necessitarão de readmissão nos primeiros dois anos, chegando a até 80% de recidivas num período de cinco anos (Robinson e cols, 1999); cerca de 10% a 25% não terão admissões posteriores (Fenton e McGlashan, 1987; Hegarty e cols, 1994). Anteriormente às recidivas, sinais de aviso freqüentemente aparecem, os quais usualmente consistem em sintomas não-psicóticos seguidos por distúrbios emocionais e sintomas psicóticos leves ao longo de um período de 4 a 12 semanas (Birchwood e cols, 1999; Gaebel e cols, 1993).

Fatores preditivos associados em geral à melhor evolução são: início mais tardio em relação às faixas etárias usuais, gênero feminino, indivíduos casados, personalidade pré-mórbida sociável, bom ajuste e bom funcionamento pré-mórbido, quociente intelectual mais elevado, presença de um desencadeante na esfera emocional quando do início, início agudo, ausência de

complicações perinatais, sintomas predominantemente afetivos ou sintomas positivos e ausência de desorganização ou sintomas negativos quando do início, menor número de episódios prévios, padrão por fases de episódios e remissões, bem como ausência de histórico familiar de esquizofrenia (Hegarty e cols, 1994; Davidson e McGlashan, 1997; Bottlender e cols, 2000, 2002 e 2003; Häfner, 2003).

Outros transtornos mentais comórbidos e condições médicas gerais são freqüentemente encontrados em pacientes com esquizofrenia o que colabora para uma expectativa de vida reduzida em relação à população em geral. A elevada taxa de mortalidade observada em pacientes esquizofrênicos deve-se, principalmente, ao elevado risco de suicídio (de 4 a 15% dos pacientes cometem suicídio); aos distúrbios cardiovasculares; às doenças respiratórias e infecciosas; e às injúrias traumáticas (Brown e cols, 2000). Uma das condições comórbidas mais freqüentes é o abuso de drogas, que ocorre em 15 a 71% dos pacientes com esquizofrenia (Soyka e cols, 1993; Kovasznay e cols, 1997; Bersani e cols, 2002). Fatores que influenciam o risco de abuso de substâncias estão associados ao meio ambiente social (pregresso e atual) e à personalidade pré-mórbida (Arndt e cols, 1992). Condições comórbidas podem piorar o curso e complicar o tratamento (Linszen e cols, 1994).

1.2 – Etiologia da esquizofrenia

A esquizofrenia é uma doença de etiologia complexa. Diversos grupos de pesquisa procuraram determinar o papel de variáveis biológicas específicas, tais como os fatores genéticos e bioquímicos e alterações sutis na morfologia cerebral, mas não se encontrou ainda uma alteração suficiente para explicar a etiopatogênese da doença. As idéias para tais estudos sobre a patofisiologia da esquizofrenia basearam-se em estudos farmacológicos por muitos anos. A clássica hipótese dopaminérgica suportava a presença de estado hiperdopaminérgico decorrente da potência das drogas típicas utilizadas (Seeman e Lee, 1975), gerando uma modulação diferenciada da atividade dopaminérgica devido à produção aumentada desse neurotransmissor, ou uma hipersensibilidade dos receptores de dopamina no sistema mesolímbico resultando em hiperexcitabilidade e aparecimento de sintomas positivos, e de um estado hipodopaminérgico em regiões cerebrais frontais, associado aos sintomas negativos (Sedvall e Farde, 1995). Essa hipótese tem como suporte o tratamento bem-sucedido dos sintomas psicóticos por agentes bloqueadores dos receptores de dopamina do tipo 2 (DRD2) no sistema mesolímbico.

Dados mostram que há um aumento da transmissão dopaminérgica no gânglio basal associado com um quadro de psicose (Abi-Dargham e cols, 2000) e uma alteração na resposta dopaminérgica no córtex pré-frontal em pacientes apresentando disfunções cognitivas crônicas (Weinberger e cols, 2001). Estudos psicofarmacológicos demonstraram que, além da dopamina, outros neurotransmissores (como a serotonina e o glutamato) parecem estar envolvidos na fisiopatologia da esquizofrenia. (Meltzer e cols, 1989; Javitt e Zukin, 1991). No entanto, como todos esses sistemas interagem, essas hipóteses não são mutualmente excludentes, e, ao menos, a modulação do sistema dopaminérgico pode ser uma ação secundária de uma alteração na sinalização glutamatérgica cortical (Coyle, 2006).

Algumas das evidências para o papel do glutamato são oriundas da farmacologia, tais como o fato de fenilciclidina e a cetamina, antagonistas do receptor de glutamato N-metil-D-aspartato (NMDA), poderem causar anormalidades cognitivas e psicóticas; e pacientes com esquizofrenia parecem ser especialmente sensíveis aos efeitos psicomiméticos dessas drogas (Ross e cols, 2006). De maneira interessante, ensaios clínicos demonstraram que agentes que modulam o receptor de NMDA (glicina, D-serina, D-cicloserina e D-alanina) melhoram, em especial, sintomas negativos quando combinados com antipsicóticos (Heresco-Levy e cols, 2002;

Lane e cols, 2005; Tsai e Gleeson; 2005). Esses dados revelam que a hipofunção do receptor NMDA, que está relacionada de maneira crítica com a função de interneurônios produtores de ácido-gamaaminobutírico (GABA), poderia contribuir para a patofisiologia da esquizofrenia (Ross e cols, 2006).

O potencial papel do GABA, principal neurotransmissor inibitório no sistema nervoso de mamíferos e que é sintetizado a partir do glutamato, na patogênese da esquizofrenia é resultante, em sua maior parte, de estudos neuropatológicos (Lewis e cols, 2005). Um subtipo particular de interneurônios de GABA, conhecidos como *chandelier*, apresentam redução da imunomarcação para o transportador de GABA, possivelmente relacionada à redução da sinalização do fator neurotrófico derivado do cérebro (BDNF) ou hipofunção do receptor NMDA. Consistente com a possível redução do transporte de GABA, estudos imunocitoquímicos e de ligação demonstraram maior atividade dos receptores de GABA do tipo A (GABA_A) nessa área. Entretanto, não é clara qual a extensão da relevância desses neurotransmissores na etiopatogênese da esquizofrenia (Ross e cols, 2006).

Alguns resultados mostram relações entre alterações do neurodesenvolvimento e esquizofrenia, tal como a maior incidência de problemas motores e neuropsicológicos em crianças com maior chance de desenvolver esquizofrenia e aumento ventricular e redução do volume cortical nos pacientes com esquizofrenia (Lawrie e cols, 1999; Pantelis e cols, 2003). A redução de algumas estruturas cerebrais relatadas podem em princípio ser o resultado do neurodesenvolvimento anormal e/ou neurodegeneração. A existência de um mecanismo alterado durante o neurodesenvolvimento é suportado pela falha de se encontrar marcadores de processos neurodegenerativos (Harrison, 1999). Por sua vez, estudos neuropatológicos também falharam em estabelecer um diagnóstico claro e alguns dados são controversos. Apesar disso, há evidências apresentadas por mais de um estudo relatando: redução do tamanho neuronal, em especial no lobo temporal, córtex pré-frontal e dorso talâmico (Harrison, 1999). Estas alterações, junto com reduções vistas em marcadores sinápticos e dendríticos e anormalidades na substância branca (Akbarian e cols, 1996; Davis e cols, 2003), sugerem problemas na estrutura sináptica e na função, bem como na conectividade, dos neurônios (Arnold e cols, 2005).

Os fatores desencadeantes, independente de quais são, parecem atuar de forma mais importante durante o neurodesenvolvimento, durante os períodos pré e perinatal, do que somente uma simples manifestação imediatamente prévia ao surgimento dos sintomas (Murray e Lewis,

1987; Weinberg, 1995, Marengo e Weinberger, 2000). Os fatores que poderiam gerar tais distúrbios podem ser desde uma exposição à infecção por influenza, herpes vírus, citomegalovírus, vírus do pólio ou *Toxoplasma gondii* já no primeiro trimestre de gravidez ou outros fatores no segundo e terceiro trimestres de gestação (Brown e Susser, 2002). Entre esses fatores de risco pode-se apresentar: rubéola e infecções respiratórias; nascimentos em baixas classes sócio-econômicas; nascimentos na zona urbana; baixo peso; incompatibilidade de fator Rh; complicações durante o parto; e nascimento durante fim do inverno e início da primavera (Lewis e cols, 1987; Dohrenwend e cols, 1992; Marcelis e cols, 1999; Susser e cols, 1996; Torrey e cols, 1997; Cannon e cols, 2002; Kyle e Pichard, 2006; St Clair e cols, 2005). O quanto a doença vai se apresentar ou não pode ser o resultado da combinação entre fatores genéticos e ambientais. Segundo Gottesman e Bertelsen (1989) e Gottesman (1991) os componentes genéticos poderiam explicar apenas metade do risco de desenvolver esquizofrenia, e as complicações pré ou perinatais seriam responsáveis por aproximadamente 1% desse risco.

1.3 – Estudos genéticos em esquizofrenia

O balanço entre componentes genéticos e ambientais – estes podendo ser biológicos ou psicossociais – estão provavelmente presentes na etiologia da maioria dos transtornos mentais (Rutter e cols, 1999). Como foi enfatizado pelo *U.S. Department of Health e Human Services* no *Mental Health's Report of the Surgeon General* em 1999:

“...dois pontos importantes sobre fatores biológicos devem ser mantidos em mente. O primeiro é que influências biológicas não são necessariamente sinônimas às de origem genética ou herdáveis. Anormalidades biológicas do sistema nervoso central que influenciam o comportamento, o pensamento ou os sentimentos podem ser causadas por trauma físico, infecções, desnutrição, ou exposição à toxinas, tais como contaminação do ambiente com chumbo. Essas anormalidades não são herdáveis. Transtornos mentais que têm maior probabilidade de ter componentes genéticos incluem autismo, transtorno bipolar, esquizofrenia e transtorno de déficit de atenção e hiperatividade (TDAH). Segundo, é errôneo assumir que fatores biológicos e ambientais são independentes um do outro, quando de fato eles interagem. Por exemplo, experiências traumáticas podem induzir alterações biológicas persistentes. Em contrapartida, crianças com um comportamento anormal com base biológica podem modificar seu ambiente...”

O modelo apresentado por Dawson e Nuechterlein (1984) integra de forma interessante estes componentes, propondo que a vulnerabilidade resultará no desenvolvimento de sintomas quando estressores ambientais estiverem presentes e os mecanismos para lidar com eles falharem. Os fatores de vulnerabilidade, baseados em um componente biológico que inclui a predisposição genética, seriam capazes de interagir com fatores complexos físicos, ambientais e psicológicos de vulnerabilidade.

Algumas doenças seguem um padrão de herança mendeliana simples, tal como a doença de Huntington e fibrose cística. Essas doenças são geralmente causadas por mutações em único

gene que resulta no surgimento da doença, apresentando uma herdabilidade que pode ser facilmente traçada durante as gerações (Chakravarti e Little, 2003). A diversidade de mutações em cada locus é alta, cada mutação é rara, pode-se afirmar que estas ocorreram recentemente na história humana e cada mutação é necessária e suficiente para causar o fenótipo de interesse (Chakravarti, 1999). Desordens que seguem esse padrão são raras. No entanto, uma grande parte das doenças que tem um componente genético seguem um padrão de herança poligênica. Nestas, muitos genes envolvidos com o aparecimento do fenótipo, não sendo possível definir um gene principal. Mutações nesses genes são comuns e apresentam apenas um pequeno efeito (Chakravarti e Little, 2003). Esses genes podem agir de forma aditiva, aumentando a susceptibilidade à doença. Este modelo requer também a existência de um limiar de susceptibilidade, a partir do qual a doença passa a ocorrer. Em indivíduos acometidos, esse limiar pode ser atingido através de diferentes combinações de fatores de risco genéticos e ambientais. Dessa forma, a presença isolada de um alelo que predisponha à doença pode não ser nem necessária ou mesmo suficiente para que esta ocorra (Conneally, 2003).

Os estudos científicos de genética em esquizofrenia iniciaram em 1916 (Kendler e Zerbin-Rudin, 1996; Kendler e cols, 1996) embora na oitava edição de seu livro texto, Kraepelin (1913) já descrevera que cerca de 70% de seus pacientes com *dementia praecox* na *Heilberg Clinic* (1891 – 1899) apresentavam história familiar de psicose (Shorter, 1997). Na análise apresentada por Zerbin-Rudin (1967), o risco para filhos de pacientes com esquizofrenia desenvolverem a doença era próxima de 15 vezes maior (12,3%) que a população em geral; irmãos e parentes cerca de 10 vezes (8,5% e 8,2%, respectivamente); para tios (2%); sobrinhos (2,2%) e netos (2,8%) (Tsuang e Vandermeij, 1980). Em onze estudos em gêmeos conduzidos entre 1928 e 1972 (Hamilton, 1976), as taxas de concordância em monozigotos variaram de 35% a 69% e em dizigóticos de 0% a 26%. Dados de McGue e Gottesman (1989) mostraram uma maior concordância entre gêmeos, atingindo 80% para monozigóticos e 50% entre os dizigóticos.

Atualmente, a existência de um componente genético na etiologia da esquizofrenia é clara, sendo que estudos mostram repetidamente a presença de um maior risco de incidência entre parentes de esquizofrênicos e que isto deve-se, em sua maior parte, a um componente genético (Gottesman e Shields, 1967). Análises complexas de segregação rejeitaram os modelos monogênicos, sustentando a hipótese de herança poligênica. Dados provenientes de estudos epidemiológicos em genética também relatam que, assim como em outras doenças, esquizofrenia

tem um padrão de transmissão complexo. Entretanto, o número de *loci* de susceptibilidade, o risco da manifestação da doença para cada locus, a grande heterogeneidade genética e o grau de interação entre esses *loci* são ainda desconhecidos.

O componente genético em esquizofrenia já foi alvo de diversas revisões de literatura (Owen e cols, 2004; Owen e cols, 2005; Riley e Kendler, 2006; Craddock e cols, 2006). Desde o primeiro estudo de ligação realizado por Sherrington e cols (1988), diversos outros estudos descreveram regiões cromossômicas que poderiam abrigar genes associados com a esquizofrenia, sendo que mais de dez diferentes locos cromossômicos já foram relacionados à esquizofrenia (Mirnics e cols, 2001). Alguns dos mais relevantes, comprovados por um maior número de estudos, estão localizados nos braços cromossômicos 1q, 2p, 5q, 6p, 8p, 10p, 17p, 20q e 22q (Brzustowicz e cols, 2000; Freedman e cols, 2001; Gurling e cols, 2001; De Lisi e cols, 2002; Mimmack e cols, 2002; Straub e cols, 2002; Lerer e cols, 2003; Lewis e cols, 2003; Ekelund e cols, 2004; Sklar e cols, 2004; Hamshere e cols, 2006; Suarez e cols, 2006). Os estudos utilizando a estratégia de genes candidatos apresentam entre os genes mais analisados e com resultados positivos replicados destacam-se disbindina 1 - DTNBP1 (6p22.3) (Straub e cols, 2002), neuregulina 1 - NRG1 (8p12) (Stefansson e cols, 2002), ativador da D-aminoácido oxidase - DAOA/G72 (13q33.2-q34) (Chumakov e cols, 2002), regulador tipo 4 da proteína G sinalizadora - RGS-4 (1q23.3) (Chowdari e cols, 2002), catecol-orto-metiltransferase - COMT (22q11.21) (Glatt e cols, 2003); prolina desidrogenase - PRODH (22q11.21) (Jacquet e cols, 2002) e *disrupted in schizophrenia 1* - DISC1 (1q42.1) (Millar e cols, 2000).

1.4 – Tratamento da esquizofrenia

A esquizofrenia é geralmente tratada com uma combinação de psicoterapia e ajustes sociais, bem como administração de fármacos. As propostas terapêuticas para a esquizofrenia mudou drasticamente nos últimos 100 anos. Inicialmente, formulações como cocaína (Becker, 1921), manganês (Reed, 1929), óleo de castor (Ingham, 1930) e injeções de óleo sulfúrico para indução de febres (Croce, 1932; Lehmann, 1993) foram utilizadas. Outros tratamentos incluíam terapia do sono e coma induzido por insulina (Ban, 2001). O primeiro tratamento bem aceito e amplamente utilizado para esquizofrenia foi a clorpromazina (Ban, 2002). A clorpromazina fora sintetizada em 1950 (Charpentier e cols, 1952) e introduzida para o uso clínico em 1952 (Delay e cols, 1952) Na mesma época, enquanto iniciava-se a utilização da clorpromazina na Europa, a reserpina fora sintetizada (1952) e introduzida na prática clínica (1954) na América do Norte (Muller e cols, 1952; Delay e cols, 1954). Porém, à reserpina restaria apenas o interesse histórico e a utilidade como ferramenta farmacológica, sendo a clorpromazina considerada o primeiro dos antipsicóticos, sendo esta uma fenotiazina. Em 1958, uma nova classe foi sintetizada, a das butirofenonas, tendo como protótipo o haloperidol (Janssen, 1996), sendo introduzida na clínica em 1959 (Divry e cols, 1959). Esses fármacos foram inicialmente denominados neurolépticos, curiosamente não devido aos seus efeitos terapêuticos, mas sim devido aos seus efeitos colaterais, os efeitos extrapiramidais. Dentro desse grupo de antipsicóticos conhecidos como típicos ou de primeira geração os mais utilizados atualmente são: clorpromazina, promazina, haloperidol, tioridazina, estelazina, trifluoperazina, tiotixene e sulpirida (Kapur e Remington, 2001).

As duas últimas décadas testemunharam mudanças significativas na utilização de terapia com antipsicóticos em todo mundo. A introdução sucessiva de onze antipsicóticos atípicos ou de segunda geração (clozapina, amisulprida, zotepina, risperidona, olazapina, quetiapina, sertindol, ziprasidona, aripiprazol, perospirona e paliperidona) criou um otimismo entre clínicos e pacientes sobre o que poderia ser alcançado em relação à eficácia terapêutica desse grupo de drogas. Assim como os 51 antipsicóticos típicos ou de primeira geração (entre eles a clorpromazina) que estão comercialmente disponíveis no mundo, estes 11 princípios ativos são, ao menos, tão eficazes na redução de sintomas típicos como ilusões, alucinações e pensamento desorganizado (Tandon e cols, 2008). Os atípicos são considerados por alguns clínicos mais

interessantes que os típicos por apresentarem maior espectro de atuação (particularmente em relação aos sintomas negativos, cognitivos e relacionados ao humor) e maior segurança em relação à manifestação de efeitos colaterais motores agudos e de longa duração (Moller, 2000; Kapur e Remington, 2001; Meltzer, 2004; Tandon, 2007). Em consequência disso, há um consenso entre médicos e associações médicas recomendando o uso desses novos agentes (Kane e cols, 2003; Miller e cols, 2004; Lehman, 2004; Falkai e cols, 2005). Embora criado tamanho entusiasmo com a introdução dos atípicos, os governos se tornaram receiosos com o aumento significativo dos custos em torno dessa classe de medicação. As despesas globais com medicações antipsicóticas multiplicaram mais de 20 vezes na última década (de 0,5 bilhão para 15 bilhões por ano). Este aumento foi causado em sua maior parte pelo fato de que os atípicos são de cinco a 30 vezes mais caros que as drogas típicas (Hoenberg e Goetz, 2006).

A eficácia terapêutica da clozapina já fora comparada em relação a outras drogas atípicas. No recente estudo no Reino Unido denominado CUtLASS (do inglês “*Cost Utility of the Latest Antipsychotics in Severe Schizophrenia*”) 136 pacientes exibindo uma resposta insatisfatória a dois ou mais agentes antipsicóticos receberam aleatoriamente clozapina ou outro atípico e a qualidade de vida fora acompanhada por um ano (Lewis e cols, 2006). Os resultados mostraram que a clozapina foi mais eficaz que os outros atípicos avaliados de forma significativa com referência à redução de sintomas ($p=0,01$). Fora igualmente observado de forma quase estatisticamente significativa ($p=0,08$) uma maior melhora na qualidade de vida neste grupo de pacientes. Esse estudo corrobora com outros que igualmente suportam uma superioridade terapêutica da clozapina em relação a outros princípios atípicos (Kane e cols, 1988; Chakos e cols, 2001; Tuunainen e cols, 2002). Os resultados de outro estudo clínico recente, o americano CATIE (do inglês “*Clinical Antipsychotic Trial of Intervention Effectiveness*”) também apresentam uma melhor resposta á clozapina em pacientes refratários (McEvoy e cols, 2006). Em contraste com os dados que suportam uma superioridade clínica da clozapina em relação aos típicos e outros atípicos em pacientes refratários e naqueles com alta taxa de tentativa de suicídio (Meltzer e cols, 2003), não há evidências de uma maior eficácia da clozapina no tratamento do primeiro episódio psicótico (Lieberman e cols, 2003) ou mesmo em outras populações de pacientes.

As diferenças clínicas observadas entre típicos e atípicos podem, ao menos em parte, serem atribuídas aos mecanismos de ação dessas drogas. Enquanto os típicos apresentam alta

afinidade de ligação *in vivo* com receptores de dopamina do tipo 2 (DRD2) e esse potencial de ligação apresenta relação com a eficácia clínica de cada droga desse grupo; a clozapina, o protótipo dos atípicos, apresenta como alvos diversos receptores, não se restringindo somente ao sistema dopaminérgico. Evidências apontam que ao menos os sistemas serotoninérgico, histaminérgico, adrenérgico e colinérgico seriam modulados pelos atípicos. Há a hipótese de que o bloqueio dos receptores de serotonina 2A (5-HT_{2A}) e o bloqueio preferencial de subtipos específicos de receptores da dopamina se constituem como um importante mecanismo para a eficácia dos antipsicóticos atípicos no tratamento dos sintomas negativos (Möller, 2003). Esses dois grupos também diferem entre si em relação à incidência dos efeitos colaterais. Enquanto pacientes em uso de típicos tendem a apresentar efeitos extrapiramidais, tal como a discinesia tardia (DT); os usuários dos atípicos são mais frequentemente acometidos com desbalanços metabólicos, sendo notáveis ganho de peso e uma maior incidência de diabetes nesse grupo. Assim como na resposta a esses medicamentos, a DT e o ganho de peso induzido por antipsicóticos parecem apresentar um componente genético que influencia a incidência bem como a gravidade desses efeitos colaterais.

A história da farmacogenética inicia-se na década 1950 após Arno Motulsky enunciar que: “traços herdados poderiam explicar as diferenças tanto no efeito das drogas quanto na presença de efeitos colaterais”, e evoluiu paralelamente com a história da genética. Diversos resultados mostram que a resposta a antipsicóticos pode ter um componente genético (Arranz e De Leon, 2007; Malhotra e cols, 2007). Em 2005, o órgão regulador americano *Food e Drug Administration (FDA)* aprovou para uso clínico um *chip* com o nome comercial de AmpliChip® CYP450 fabricado pela Roche. Este chip possibilita o teste de dois genes polimórficos, o do citocromo P450 2D6 (CYP2D6) e do citocromo P450 2C19 (CYP2C19), enzimas que são responsáveis pela metabolização de várias drogas antidepressivas e antipsicóticas (De Leon e cols, 2006).

Estudos farmacogenéticos têm analisado primariamente os medicamentos atípicos, em especial a clozapina, talvez porque seja mais fácil o acesso ao sangue desses pacientes, uma vez que estes precisam ser monitorados quanto à agranulocitose ou por causa da superior eficácia clínica em populações de paciente resistentes ao tratamento. Análises farmacogenéticas da resposta à clozapina têm utilizado a estratégia de estudos de associação com genes candidatos, usando como genes candidatos os receptores dopaminérgicos e serotoninérgico (Malhotra e cols,

2004). Estes seriam candidatos com forte racionalidade biológica já que clozapina apresenta alta afinidade com estes receptores.

Arranz e colaboradores, em 1995, chamaram a atenção para o receptor 5-HT_{2A} reportando uma associação significativa com o alelo 102C e uma pior resposta à clozapina numa população de 149 pacientes com esquizofrenia. Entretanto, esse resultado não foi consistentemente replicado por uma série de outros estudos, sendo que alguns incluíram outros antipsicóticos (Masellis e cols, 1995; Burnet e Harrison, 1995). A variação T102C no 5-HT_{2A} pode ser considerada como um fraco candidato uma vez aqui a troca das bases não implica na troca de aminoácidos na proteína e nenhuma função fora até então descrita para essa mudança de bases (Masellis e cols, 1995). A variação His452Tir, que não foi encontrada em desequilíbrio de ligação com o T102C (Malhotra e cols, 1996), aparenta produzir alterações funcionais *in vitro*. Entretanto, não fora relatada nenhuma forte associação dessa variante com a resposta à antipsicóticos (Nothen e cols, 1995). A variante -1438G/A já foi analisada nesse mesmo contexto e não foram relatadas associações (Arranz e cols, 1998a).

Outros genes no sistema serotoninérgico foram avaliados, entre eles os receptores 5-HT_{1A} (Masellis e cols, 2001), 5-HT_{2C} (Sodhi e cols, 1995), 5-HT_{3A/B} (Arranz e cols, 2000b), 5-HT_{5A} (Birkett e cols, 2000), 5-HT₆ (Yu e col, 1999), 5-HT₇ (Masellis e cols, 2001), o transportador de serotonina - 5HTT (Arranz e cols, 2000a; Tsai e cols, 2000) e triptofano hidroxilase – TPH (Anttila e cols, 2007). Embora alguns resultados apresentem associações positivas, há somente uma fraca indicação de que esses genes são associados com a resposta à antipsicóticos. Genes no sistema dopaminérgico também foram extensivamente analisados, tais como receptores DRD1 (Potkin e cols, 2003), DRD2 (Arranz e cols, 1998b), DRD3 (Shaikh e cols, 1996), DRD4 (Shaikh e cols, 1993) e o transportador de dopamina – DAT (Szekeres e cols, 2004).

Diversos outros genes já foram igualmente reportados em análises de associação com a resposta aos antipsicóticos, entre eles a subunidade 2B do receptor NMDA de glutamato - GRIN2B (Hong e cols, 2001a), o receptor de histamina do tipo 1 e 2 – H₁ e H₂ (Mancama e cols, 2002), receptores α 1 e α 2 adrenérgicos (Bolonna e cols, 2000; Tsai e cols, 2001; De Luca e cols, 2005), transportador de norepinefrina – NET (Meary e cols, 2007); receptor de neurotensina (Huezo-Diaz e cols, 2004), fator neurotrófico derivado do cérebro – BDNF (Hong e cols, 2003), antígeno HLA-A1 (Lahdelma e cols, 2001), fator de necrose tumoral alfa (Tsai e cols, 2003), apolipoproteína E (Hong e cols, 2000), COMT (Yamanouchi e cols, 2003), subunidade β 3 da

proteína G – GNB-3(Muller e cols, 2005a), RGS-2 (Greenbaum e cols, 2007), RGS-4 (Kampman e cols, 2006), proteínas de ligação a fator solúvel sensível a N-etilmaleimida SNAP-25 (Muller e cols, 2005), resistência múltipla à drogas MDR-1/ABCB1 (Yasui-Furukori e cols, 2006), NEF3(Strous e cols, 2007), proteína quinase B – PKB/Akt1 (Xu e cols, 2007), glicogênio sintase quinase 3 isoforma β (Souza e cols, 2008), colina acetiltransferase – ChAT (Mancama e cols, 2007), NOTCH4 (Anttila e cols, 2004); NRG-1 (Kampman e cols, 2004), enzima conversora de angiotensina – ECA (Illi e cols, 2003) e glicoproteína P – PGP (Lin e cols, 2006).

1.4.1 – Discinesia tardia induzida por antipsicóticos

A DT é uma síndrome extrapiramidal induzida por antipsicóticos, caracterizada por movimentos involuntários, anormais e repetitivos localizados principalmente na região orofacial, tronco, extremidades inferiores e superiores, podendo acometer inclusive o sistema respiratório. O termo DT foi introduzido por Faurbye (1964), como tardia enfatizando a cinética temporal até a apresentação de movimentos involuntários, já que esse efeito colateral é causado por uma longa utilização de antipsicóticos. O diagnóstico apresentado pelo DSM-IV (American Psychiatric Association, 2000) requer pelo menos três meses de exposição a essas drogas. A DT é potencialmente irreversível com a descontinuação do uso da medicação.

Os dados de prevalência são difíceis de interpretar uma vez que apresentam estudos em populações heterogêneas e formas diagnósticas diferentes. Em uma revisão 56 estudos entre 1959 e 1979, Kane e Smith (1982) encontraram uma prevalência de 0,5 até 65%. Em 1992, Yassa e Jeste reportaram uma prevalência de 24% em 39187 pacientes de 76 estudos. Dois outros estudos utilizando mesmo critérios diagnósticos, Woerner e cols (1991) e Muscettola e cols (1993), apresentaram prevalência de 23,4% e 19,1%, respectivamente. Dados mais recentes mostram que DT acomete pelo menos 20% dos indivíduos em uso de antipsicóticos, com taxas de incidência para novos casos de aproximadamente 3 a 5% ao ano (Kane, 2001). Essa incidência parece ocorrer de maneira cumulativa e chegar a 30% entre os idosos expostos ao uso crônico de antipsicóticos. Além dos antipsicóticos, outros fatores que têm sido relacionados ao aparecimento e prognóstico da discinesia tardia são: idade, gênero feminino, co-morbidade psiquiátrica, presença de outros transtornos extrapiramidais na fase aguda do tratamento com antipsicóticos e diabetes (Kane, 2001).

Em função da complexidade dessa síndrome, diversas teorias fisiopatológicas têm sido sugeridas, de maneira geral implicando um ou mais neurotransmissores no aparecimento dos sintomas de DT. No passado, acreditava-se que o uso crônico de antipsicóticos provocasse uma hipersensibilidade dopaminérgica na região nigro-estriatal que levaria ao surgimento dos sintomas de DT. Entretanto, essa hipersensibilidade não explica a susceptibilidade individual para desenvolver discinesia tardia e passou-se então a investigar a ocorrência de possíveis alterações concomitantes nos neurotransmissores colinérgicos, GABAérgicos e serotoninérgicos. Além disso, tem-se sugerido que o uso crônico desses medicamentos causaria uma

superprodução de radicais livres de oxigênio com conseqüente degeneração neuronal (Cadet e Lohr, 1989). Essa degeneração, a princípio reversível, resultaria em lesões irreversíveis dos neurônios nigroestriatais, com conseqüente morte celular, e o aparecimento da DT.

Há uma forte concordância em relação a incidência de DT entre parentes de primeiro-grau em uso de antipsicóticos (Youssef e cols, 1989; Müller e cols, 2001). Em função de que as primeiras explicações para o surgimento de DT estarem vinculadas a uma maior atividade do sistema dopaminérgico, em especial no gânglio basal, e as drogas típicas (mais frequentemente associadas a esse efeito colateral) serem antagonista de receptores dentro desse sistema, genes que codificam moléculas envolvidas com a propagação do sinal dopaminérgico foram primeiramente analisadas em estudos de associação (Ozdemir e cols, 2001). Confirmando a plausibilidade biológica, o DRD2 apresentou resultados positivos, especialmente com a variante C939T (Chen e cols, 1997). Da mesma forma, sugere-se associação significativa com outros genes do sistema dopaminérgico: DRD1 (Srivastava e cols, 2006); DRD3 (Rietschel e cols, 1993). DRD4 (Rietschel e cols, 1996) e o DAT (Segman e cols, 2003) já foram igualmente analisados entretanto não apresentaram resultados significativos.

Genes em outros sistemas de neurotransmissão, tais como serotoninérgico, e genes envolvidos com o processamento do estresse oxidativo (Thelma e cols, 2007) foram também reportados. Entre eles encontra-se resultados com: 5-HT_{2A} (Segman e cols, 2001); 5-HT_{2C} (Segman e cols, 2000), 5-HT₆ (Ohmori e cols, 2002), 5-HTT (Chong e cols, 2000), TPH (Segman e cols, 2003), COMT (Herken e cols, 2003); monoamino oxidase – MAO (Matsumoto e cols, 2004), GRIN2B (Liou e cols, 2007), receptor de adenosina 2A – A_{2A} (Hong e cols, 2005), receptor opióide μ e Δ (Ohmori e cols, 2001), receptor de estrógeno (Lai e cols, 2002), GNB-3 (Lee e cols, 2007), PGP (de Leon e cols, 2005), BDNF (Liou e cols, 2004), mangânes superóxido dismutase – MnSOD (Hori e cols, 2000), NAD(P)H quinona oxireductase - NQO1 (Pae e cols, 2004); glutationa peroxidase – GPX1 (Shinkai e cols, 2006); óxido nítrico sintase – NOS (Shinkai e cols, 2004), fenilalanina hidroxilase (Richardson e cols, 2006); glutationa-S-transferases – GSTM1 e GSTT1 (Pae e cols, 2004), ECA (Segman e cols, 2002), CYP1A2 (Basile e cols, 2000); CYP2D6 (Arthur e cols, 1995), CYP3A4 (Tiwari e cols, 2005) e CYP3A5 (de Leon e cols, 2005).

1.4.2 – Ganho de peso induzido por antipsicóticos

O ganho de peso é um sério problema em pacientes usando antipsicóticos (Malhotra e cols, 2004; Müller e cols, 2006). Evidências mostram que os antipsicóticos interagem no sistema neuroendócrino, levando a efeitos colaterais como aumento do apetite, obesidade, hiperglicemia e diabetes (Bernstein, 1987; Henderson e cols, 2000). O excesso de peso é um evento comum nesses pacientes, tendo sido demonstrado que os mesmos apresentam um índice de massa corporal significativamente maior do que os pacientes psiquiátricos sem o diagnóstico de esquizofrenia e do que a população geral (Allison e Casey, 2001). A magnitude do ganho de peso varia conforme os medicamentos e a dosagem, sendo que alguns se destacam por ganhos de peso de 1,5 a 8,8 kg em períodos de 6 meses (Allison e cols, 1999; Goudie e cols, 2005). Vários estudos convergem e sugerem que alguns antipsicóticos atípicos implicam em ganho de peso significativamente maior após a administração em curto e em longo prazo, quando comparados com antipsicóticos típicos (Henderson, 2007).

O excesso de peso corporal aumenta intensamente o risco de mortalidade e morbidade de vários transtornos clínicos, incluindo hipertensão, dislipidemia, diabetes melito tipo II, doenças cardíacas, doenças da vesícula biliar, osteoartrite, apnéia do sono, problemas respiratórios e cânceres de endométrio, mama, próstata e cólon, reduzindo ainda mais a sobrevida e a qualidade de vida dos portadores de esquizofrenia (NIH, 1998). Desta forma, há uma atenção para se adequar a prescrição ao perfil do paciente, questionando sobre outros fatores de risco, como hipertensão, diabetes prévia, idade maior que 50 anos, raça e história familiar. Além disso, o monitoramento dos níveis glicêmicos e ponderais deve ser feito para orientar estratégias no tratamento destas alterações (Meltzer, 2001). Apesar de recente e pouco explorado, Wehmeier e colaboradores e Theisen e colaboradores, ambos em 2005, apresentaram dados que mostram concordância de ganho de peso induzida por antipsicótico em gêmeos monozigóticos e/ou pares de irmãos do mesmo sexo.

A maioria dos estudos genéticos que avaliaram a indução de ganho de peso por antipsicóticos examinaram genes que codificam proteínas do sistema nervoso central, especialmente receptores de neurotransmissores. A seleção de genes candidatos a serem analisados nesse contexto baseia-se primariamente nas conhecidas bases neurobiológicas da saciedade. Por outro lado, é possível pensar que a medicação possa também operar em

mecanismos periféricos, tais como o controle do metabolismo e tônus muscular modulando assim a queima de calorias e/ou diretamente alterando a lipogênese (Basile e cols, 2001). Grande parte dos estudos tem se focado no sistema serotoninérgico que é conhecido como controlador da saciedade. Os sinais de controle de saciedade convergem no hipotálamo provenientes de diversas áreas do corpo, incluindo receptores gustativos, olfatórios, gástricos, intestinais e hepáticos. Outros pontos que fortalecem a idéia de um papel da serotonina nesse contexto surgem em torno da capacidade dessa amina modular o comportamento alimentar em diversos modelos analisados. De uma forma geral, o aumento da concentração de serotonina estaria relacionado a uma diminuição do comportamento alimentar, sendo que a relação inversa é igualmente verdadeira (Davis e Faulds, 1996). Alguns receptores e o transportador de serotonina já foram analisados nesse contexto: 5-HT_{1A} (Basile e cols, 2001), 5-HT_{2A} (Hong e cols, 2001b), 5-HT_{2C} (Rietschel e cols, 1997), 5-HT₆ (Hong e cols, 2001b), 5-HTT (Hong e cols, 2001b). Estudos exploratórios com: DRD1 (Lane e cols, 2006), DRD2 (Lane e cols, 2006), DRD3 (Lane e cols, 2006), DRD4 (Rietschel e cols, 1996), H₁ (Basile e cols, 2001), H₂ (Basile e cols, 2001), SNAP-25 (Müller e cols, 2005b), CYP2D6 (Ellingrod e cols, 2002), CYP1A2 (Basile e cols, 2001), receptores adrenérgicos (Basile e cols, 2001), leptina e seu receptor (Zhang e cols, 2003), neuropeptídeo Y e seus receptores (Ruaño e cols, 2007), paraoxonase 1 (Ruaño e cols, 2007), apolipoproteínas A4 e E (Ruaño e cols, 2007), PGP (Lin e cols, 2006), TNF- α (Zai e cols, 2006), BDNF (Lane e cols, 2006) e GNB-3 (Tsai e cols, 2004) foram reportados.

2. Objetivos

- Determinar se há participação genética dos receptores α do fator neurotrófico derivado de glia (GDNF) na esquizofrenia e na resposta ao tratamento de clozapina.
- Estudar a predisposição genética à esquizofrenia e à resposta a clozapina associada a marcadores no gene da GSK-3.
- Avaliar o papel do gene NALCN na genética de esquizofrenia, resposta ao tratamento, discinesia tardia induzida por antipsicóticos e ganho de peso induzido por clozapina.
- Analisar se a variante C825T no gene da GNB-3 está associado com o ganho de peso induzido por antipsicóticos e no índice de massa corpórea

3. Artigos

3.1. Association study of GSK3 gene polymorphisms with schizophrenia and clozapine response

Psychopharmacology
DOI 10.1007/s00213-008-1193-9

ORIGINAL INVESTIGATION

Association study of GSK3 gene polymorphisms with schizophrenia and clozapine response

Renan P. Souza · Marco A. Romano-Silva ·
Jeffrey A. Lieberman · Herbert Y. Meltzer ·
Albert H. C. Wong · James L. Kennedy

Received: 29 February 2008 / Accepted: 27 April 2008
© Springer-Verlag 2008

Abstract

Rationale A number of human and animal studies implicate GSK3 in the pathophysiology and genetics of schizophrenia. In general, the data suggest that phosphorylation levels of GSK3 β are reduced in schizophrenia, resulting in increased GSK3 β activity. Since GSK3 β regulation is altered in schizophrenia, polymorphic variation in this gene may affect susceptibility to schizophrenia or treatment response.

Objective To analyze GSK3 β genetic variants for association with schizophrenia and clozapine response.

Materials and methods We examined GSK3 β markers in 185 matched case-control subjects, 85 small nuclear

families, and 150 schizophrenia patients treated with clozapine for 6 months.

Results Three markers (rs7624540, rs4072520, and rs6779828) showed genotypic association with schizophrenia in the case-control sample. We did not observe any family and clozapine response association with a specific allele, genotype, or haplotype.

Conclusions Our results suggest that GSK3 β polymorphisms might be involved in schizophrenia risk but do not appear to play a significant role in clozapine response.

Keywords Schizophrenia · Clozapine response · GSK3 β · Wnt signaling · Genetic association · Family-based association test

R. P. Souza (✉) · M. A. Romano-Silva
Grupo de Pesquisa em Neuropsiquiatria Clínica e Molecular,
UFMG,
Belo Horizonte, Brazil
e-mail: renanps@yahoo.com.br

R. P. Souza · M. A. Romano-Silva
Saude Mental, UFMG,
Belo Horizonte, Brazil

R. P. Souza · A. H. C. Wong · J. L. Kennedy
Neurogenetics Section, CAMH,
Toronto, ON, Canada

J. L. Kennedy
Department of Psychiatry, University of Toronto,
Toronto, ON, Canada

J. A. Lieberman
Department of Psychiatry, University of North Carolina,
Chapel Hill, NC, USA

H. Y. Meltzer
Psychiatric Hospital, Vanderbilt University,
Nashville, TN, USA

Introduction

Schizophrenia is a devastating psychiatric disorder that affects 1% of the population, and pharmacological treatment is only partially effective for the majority of patients. There are currently no clear criteria by which to predict treatment response or to select the optimal drug that will have the best efficacy and minimal side effects (Seeman 2002; Ananth et al. 2004). Approximately 10–20% of patients do not respond to antipsychotic drug treatment, and an additional 20–30% do respond early on eventually relapse; others develop serious side effects that cause them to discontinue medication altogether (Malhotra et al. 1993, 2004; Basile et al. 2002). Pharmacogenetic studies may help to identify genetic variants that modulate treatment response or side-effect profiles, with the eventual objective of applying this knowledge to clinical practice.

Glycogen synthase kinase 3 (GSK3) phosphorylates, and thereby regulates, a large number of substrates, with over

50 known to date (Jope et al. 2007). Four GSK3 regulatory mechanisms have been described: phosphorylation, protein complexes, localization, and substrate phosphorylation; and in combination, these can provide substrate-specific regulation of GSK3 activity. The most well-defined regulatory mechanism is inhibition of GSK3 activity by phosphorylation of serine-9 in the isoform GSK3 β or serine-21 in GSK3 α (Woodgett 1990; Sutherland et al. 1993). The phosphatidylinositol 3-kinase/Akt signaling pathway is activated by insulin and many growth factors, and this pathway can also regulate GSK3 through Akt phosphorylation of GSK3 on these same serine residues. Protein kinase C and protein kinase A are also able to phosphorylate these regulatory serines (Shin et al. 2002; Jope and Johnson 2004).

There is support for the hypothesis that alterations in GSK3 are connected with schizophrenia and its treatment, mainly in the areas of altered dopaminergic activity and disrupted neurodevelopment. All antipsychotics bind to dopamine D2 receptors with an affinity inversely proportional to the clinically effective dose (Seeman and Lee 1975). Some antipsychotics with very high D2 receptor affinity such as haloperidol bind with low affinity to other related G-protein-coupled neurotransmitter receptors (Kapur and Seeman 2001). In contrast, some antipsychotics such as clozapine have relatively low D2 receptor affinity but bind strongly to 5-HT₂ receptors (Meltzer 1989). Therefore, typical and atypical antipsychotics may regulate GSK3 activity via different signaling pathways. Previously, Kang et al. (2004) demonstrated that although both Dvl and Akt are activated by clozapine, only Dvl is responsible for Ser9-phosphorylation of GSK3 in SH-SY5Y cells. More recently, Roh et al. (2007) compared the phosphorylation at Ser21/9 residues of GSK3 in vivo after acute treatment with haloperidol or clozapine. Acute treatments with both drugs increased pSer21/9-GSK3 in vivo, but they affected upstream regulators differently. Furthermore, Li et al. (2007) recently have reported that clozapine increases pSer9-GSK3 β .

Low immunoreactivity and activity of GSK3 β was demonstrated post mortem in the frontal cortex of schizophrenia patients (Kozlovsky et al. 2000, 2001, 2004; Beasley et al. 2001; Nadri et al. 2003), but there is at least one negative study on a different brain collection (Nadri et al. 2004). Previous genetic studies have not found significant association between GSK3 β and schizophrenia in case-control samples (Scassellati et al. 2004; Ikeda et al. 2005; Lee et al. 2006; Szczepankiewicz et al. 2006). However, the potential effect of genetic variation in GSK3 β on clozapine response has not yet been explored. Thus, our objectives were to examine genetic association between GSK3 β and schizophrenia (using case-control and nuclear family samples), as well as clozapine response.

Materials and methods

Clinical sample

All recruitment and clinical assessments were conducted with written informed consent and with the explicit approval of our institutional ethics review board. Clinical data and DNA samples were obtained from the probands of 85 small nuclear families, as well as 180 Caucasian patients with a DSM-III-R or DSM-IV diagnosis of schizophrenia. The healthy controls ($N=185$) were matched for age (± 5 years) and sex, 146 male and 73 female cases and the same number of controls, mean age 36 ± 8). The Structured Clinical Interview for DSM-IV Axis I Disorders was administered by trained research assistants to each patient, and diagnosis was supplemented by a review of medical records. The diagnosis was established via consensus procedures incorporating two of the investigators. The controls were screened for current or past history of major psychiatric disorders.

For the clozapine response sample, clinical data from 140 patients with a DSM-III-R diagnosis of schizophrenia (114 Caucasians), almost all of whom were treatment refractory or intolerant of typical antipsychotic therapy (Kane et al. 1988), were obtained at two research clinics: Case Western Reserve University in Cleveland, OH, USA ($n=90$; 63 males and 27 females, mean age 36 ± 8) and Hillside Hospital in Glen Oaks, NY, USA ($n=50$; 33 males and 17 females, mean age 35 ± 8). After informed consent was obtained, patients underwent a washout period of 2 to 4 weeks during which, unless clinically necessary, they received no medications before starting clozapine. Clozapine treatment was continued for a minimum of 6 months during which patients were evaluated prospectively. Clozapine blood levels were monitored during the course of treatment to ascertain compliance. Treatment response was evaluated as the percentage score change on the 18-item Brief Psychiatric Rating Scale (BPRS). Treatment response was expressed as a dichotomous variable at 6 months using criteria based on those of Kane (Kane et al. 1988): a reduction of $\geq 20\%$ on the overall score of the BPRS from the baseline score at enrolment. There were no differences observed between the sites in terms of gender ratio, mean age, mean age of onset, or response ratio. Caucasians and African-American subjects were not significantly different in terms of gender ratio, mean age, mean age of onset, or response ratio (data not shown; Hwang et al. 2005).

Genetic analyses

Genomic DNA was extracted using the high salt method of Lahiri and Numberger (1991). We analyzed 12 markers across GSK3 β (rs6805251, rs4688043, rs7624540,

rs13319151, rs6438552, rs4072520, rs9878473, rs4491944, rs6772172, rs11919783, rs11923196, rs6779828, rs9846422, rs3755557) and two markers after the 3'-untranslated region of the gene (rs9846422, rs3755557). Genotyping was performed by GoldenGate assay (Illumina, San Diego, CA, USA) at The Centre for Applied Genomics at the Hospital for Sick Children in Toronto, ON, Canada. Rs4072520 was re-genotyped in the whole sample using TaqMan® SNP Genotyping Assays from Applied Biosystems as a quality check procedure.

Statistical analyses

Individual markers analyses of responder (control)/non-responder (case) data were performed using χ^2 tests. The statistical program used was the Statistical Package for the Social Sciences, version 10.0.7. We applied the family-based association test (FBAT, version 1.0, Laird et al. 2000) under the assumption of an additive model and PEDSTATS (Wigginton and Abecasis 2005) for Hardy-Weinberg equilibrium in the family data. Linkage disequilibrium (LD) was assessed using Haploview, version 4.0 (Barrett et al. 2005). Haplotype analyses were performed using UNPHASED 3.0.10 (Dudbridge 2003) and Haploview, version 4.0.

Results

Linkage disequilibrium analysis

Pairwise LD between the markers is presented for each sample (Fig. 1). In this study, we defined a haplotype block as a region over which less than 5% of pairwise comparisons among informative markers showed strong evidence of historical recombination (upper confidence bound on D' less than 0.9; Gabriel et al. 2002). We found the same block in each of the case-control, family, and response samples composed of 12 of the 14 markers.

Genotype data

Significant deviations from Hardy-Weinberg equilibrium were observed for rs7624540 ($p=0.007$), rs4075250 ($p=0.004$), and rs6779828 ($p=0.023$) in cases from the case-control sample (Fig. 2). Only two markers in cases showed deviation from Hardy-Weinberg equilibrium at a significance threshold of $p<0.01$. Cases and controls were compared for genotype and allele frequencies across the markers (see Table 1). The following markers were genotypic distribution associated with schizophrenia: rs7624540 ($\chi^2=8.28$, genotype $p=0.016$), rs4072520 ($\chi^2=9.94$, genotype $p=0.007$), and rs6779828 ($\chi^2=6.83$, genotype $p=0.033$).

In the family samples, no significant deviation from Hardy-Weinberg equilibrium was observed among 81 unrelated individuals (data not shown). Allele frequencies are shown in Table 2. Family-based association results showed no association with schizophrenia. In the clozapine response sample, there were no significant deviations from





Fig. 1 LD plot for the analyzed markers in case-control, family, and response samples (a, b, c). Values presented are the D'

Fig. 2 Hardy–Weinberg equilibrium p values for case–control and responders (cases)–non-responders (controls). The solid line represents $p=0.05$



Hardy–Weinberg equilibrium (Fig. 2). Responder/non-responder groups were compared for genotype and allele frequencies across the markers, and no significant differences were observed (see Table 3), including when Caucasian or African-American subjects were analyzed separately (data not shown).

Haplotype analyses

Cases and controls were compared for haplotype frequencies across the *GSK3 β* markers. We performed a three-marker window (Durrant et al. 2004), and results did not show association. Allelic occurrence in haplotypes constructed by rs7624540, rs4072520, and rs6779828 (markers that showed single genotypic association) showed an overall trend ($\chi^2=9.22$, $p=0.05$), as well as the A–A–G haplotype ($\chi^2=3.09$, $p=0.078$; after 1,000 permutations $p=0.223$). Furthermore, genotypic occurrence in haplotypes constructed by rs7624540, rs4072520, and rs6779828 showed an overall significant association ($\chi^2=23.79$, $p=0.001$), and the A/A–A/A–A/A haplotype presented the strongest association ($\chi^2=7.98$, $p=0.004$; after 1,000 permutations $p=0.011$). Considering haplotypes in the main LD block, no association was found. In the family sample, no haplotype associations were observed. Haplotype frequencies in clozapine responders and non-responders were compared, and no associations were observed in the whole sample, nor were there associations for specific haplotype

blocks when Caucasian or African-American subjects were analyzed separately.

Discussion

This exploratory study examined the association of 14 markers spanning the *GSK3 β* gene and two phenotypes: schizophrenia (case–control and nuclear family samples) and clozapine response. Three individual markers (rs7624540, rs4072520, and rs6779828) and one haplotype (rs7624540, rs4072520, and rs6779828) showed nominally significant association with schizophrenia in our case–control sample. The matched case–control sample has 43.2% power to detect a significant effect with a genotypic relative risk of 2.0 (Purcell et al. 2003). Although gene variants of small effect or those that are rare major determinants of risk for developing schizophrenia would likely not be detected because very large sample size would be required, we have found three positive associations. Additional studies in larger samples are needed to resolve whether *GSK3 β* is truly playing a role in schizophrenia, especially after our significant findings. No associations were detected in the nuclear family samples.

Corrections for multiple testing have been a controversial issue (Aickin 1999; Bender and Lange 1999; Perneger 1998), and considering the exploratory nature of this study,

Table 1 Allelic and genotypic frequencies for the case-control association

SNP	Allele frequency						Genotype frequency						Association			
	Control			Case			Control			Case			Allele		Genotype	
	Samples	Percent	Case	Samples	Percent	Case	Samples	Percent	Case	Samples	Percent	Case	Chi square	p value	Chi square	p value
rs6805251	A	141	38.1	A	138	38.3	AA	27	14.6	AA	21	11.7	0.01	0.950	1.60	0.449
	C	229	61.9	C	222	61.7	AC	87	47.0	AC	96	53.3				
rs4688043	A	346	93.5	A	335	93.1	AA	161	87.0	AA	155	86.0	0.06	0.804	0.66	0.797
	G	24	6.5	G	25	6.9	AG	24	13.0	AG	25	14.0				
rs7624540	A	70	18.9	A	60	16.7	GG	00	00.0	GG	00	00.0	0.63	0.426	8.28	0.016
	G	300	81.1	G	300	83.3	AA	08	04.3	AA	00	00.0				
rs13319151	A	00	00.0	A	01	00.3	AG	54	29.2	AG	60	33.3	1.03	0.867	1.03	0.867
	G	359	100.0	G	359	99.7	GG	123	66.5	GG	120	66.7				
rs6438552	A	229	61.9	A	222	61.7	AA	00	00.0	AA	00	00.0	0.01	0.950	2.20	0.333
	G	141	38.1	G	138	38.3	AG	85	45.9	AG	96	53.3				
rs4072520	A	71	19.2	A	63	17.5	GG	28	15.1	GG	21	11.7	0.35	0.555	9.94	0.007
	C	299	80.8	C	297	82.5	AA	09	05.0	AA	00	00.0				
rs9878473	A	229	61.9	A	222	61.7	AC	53	29.0	AC	63	35.0	0.01	0.950	2.20	0.333
	T	141	38.1	T	138	38.3	CC	123	65.0	CC	117	65.0				
rs4491944	A	01	0.03	A	01	0.03	AA	00	00.0	AA	00	00.0	0.01	0.999	0.01	0.984
	T	369	99.7	T	359	99.7	AT	01	00.5	AT	01	00.6				
rs6772172	A	359	100.0	A	359	99.7	TT	184	99.5	TT	179	99.4	1.03	0.867	1.03	0.867
	G	00	00.0	G	01	00.3	AG	00	00.0	AG	01	00.6				
rs11919783	A	44	11.9	A	32	8.9	GG	00	00.0	GG	00	00.0	1.76	0.184	5.01	0.082
	G	326	88.1	G	328	91.1	AA	05	02.7	AA	00	00.0				
rs11923196	A	00	00.0	A	01	00.3	AG	34	18.4	AG	32	17.8	1.03	0.867	1.03	0.867
	T	359	100.0	T	359	99.7	GG	146	78.9	GG	148	82.2				
rs6779828	A	71	19.2	A	62	17.2	AA	00	00.0	AA	00	00.0	0.47	0.491	6.83	0.033
	G	299	80.8	G	298	82.8	AT	00	00.0	AT	01	06.6				
rs9846422	A	359	100.0	A	359	99.7	TT	185	100.0	TT	179	99.4	1.03	0.867	1.03	0.867
	G	00	00.0	G	01	00.3	AG	00	00.0	AG	01	06.6				
rs3755557	A	312	84.3	A	311	86.4	AA	133	71.9	AA	135	75.0	0.62	0.430	0.63	0.728
	C	58	15.7	C	49	13.6	AC	46	24.9	AC	41	22.8				
						CC	06	03.2	CC	04	02.2					

Table 2 Family-based association test results

SNP	Allele	Frequency	Family	S	$E(S)$	$Var(S)$	Z	p
rs11919783	A	0.125	13	10	7.333	3.222	1.486	0.137
rs11923196	A	0.003	00	–	–	–	–	–
rs13319151	A	0.005	01	–	–	–	–	–
rs3755557	A	0.819	17	22	23.833	4.694	–0.846	0.397
rs4072520	A	0.238	29	19	17.783	7.764	0.437	0.662
rs4491944	A	0.010	02	–	–	–	–	–
rs4688043	A	0.885	13	20	18.217	3.820	0.912	0.362
rs6438552	A	0.561	34	41	41.950	10.264	–0.297	0.767
rs6772172	A	0.995	01	–	–	–	–	–
rs6779828	A	0.225	27	17	16.117	7.042	0.333	0.739
rs6805251	A	0.436	34	29	28.050	10.264	0.297	0.767
rs7624540	A	0.226	27	18	16.783	7.264	0.451	0.652
rs9846422	A	0.995	01	–	–	–	–	–
rs9878473	A	0.562	34	41	41.950	10.264	–0.297	0.767

S Test statistic for observed number of alleles, E expected value of S under null hypothesis, $Var(S)$ variance between the observed and expected transmission

without prespecified hypotheses for most of our markers, the individually significant associations require independent confirmation (Bender and Lange 1999). No associations remain significant after Bonferroni correction (Bonferroni-corrected threshold $p < 0.003$). Bonferroni correction is based on the assumption that tests are independent. This is a very conservative adjustment for these data due to the high levels of LD. Applying the Nyholt (2004) correction (Nyholt-corrected threshold $p < 0.006$), a nearly significant result for rs4072520 (genotype $p = 0.007$) is found. Nyholt showed that significant results occurring after calculating the adjusted alpha value that are not significant with standard Bonferroni or Sidak correction indicate substantial non-independence (i.e., strong intermarker LD) among the tested markers. Therefore, permutation testing was applied as strong LD can be seen (Fig. 1). After 1,000 permutations, the rs4072520 association remains significant ($p < 0.008$).

All markers that showed nominal individual significant association with schizophrenia deviated ($p < 0.05$) from Hardy–Weinberg equilibrium among the cases. Although the standard χ^2 test often has been used, the Fischer exact test is preferred when genotype counts are low since it does not rely on the χ^2 null distribution approximation (Guo and Thompson 1992; Wigginton et al. 2005). With the Fisher exact test, significant deviations from Hardy–Weinberg equilibrium were observed in cases (rs7624540 $p = 0.005$; rs4072520 $p = 0.002$; rs6779828 $p = 0.032$). At a significance threshold of $p < 0.01$, just two markers in the cases show deviation from the Hardy–Weinberg equilibrium (HWE). Trikalinos et al. (2006) concluded that HWE should be routinely and transparently assessed in gene-disease association studies. An outlier marker with extreme HWE disequilibrium usually results from genotyping error, so it has also been used to detect such errors (Hoh et al.

2001). None of the markers appears to be an outlier according to the HWE statistics, so HWE resulting from genotyping error is not a likely explanation for the observed results. In our sample, it is more likely to be an association finding in a recessive genetic model. The detection of HWE disequilibrium has recently been shown to be useful as a method for detecting gene–phenotype association (Feder et al. 1996; Nielsen et al. 1998; Hoh et al. 2001; Lee 2003; Hao et al. 2004; Wittke-Thompson et al. 2005; Luo et al. 2005, 2006; Balding 2006).

Three other studies analyzing *GSK3 β* and schizophrenia have been published. Using a sample of 147 patients and 212 healthy individuals, Scassellati et al. (2004) analyzed two markers at positions –1727 A/T and –50 C/T and a (CAA) (n) repeat polymorphism localized in intron 1 of the gene. Their results showed no overall associations. However, in schizophrenia patients with paranoid subtype, they found that heterozygous (CAA) (three)/(CAA) (five) subjects were over-represented. Ikeda et al. (2005) analyzed seven markers and the (CAA) (n) repeat in 381 cases and 352 controls and detected no association with schizophrenia in a Japanese population. Lee et al. (2006) analyzed the same two markers in 138 cases and 350 controls and did not find any association in a Korean population. Furthermore, Szczepankiewicz et al. (2006) evaluated T50C in 432 cases and 408 controls and find no association with schizophrenia. Our results are the first analysis of genetic association the *GSK3 β* gene with schizophrenia based on both matched case–control and family-based samples.

A number of human and animal studies have emerged that implicate GSK3 in the pathophysiology and genetics of schizophrenia (Cotter et al. 1998; Beaulieu et al. 2004; Emamian et al. 2004). In general, the data suggests that phosphorylation levels of GSK3 β are reduced in schizo-

Table 3 Allelic and genotypic frequencies for the clozapine response association

SNP	Allele frequency			Genotype frequency						Association						
	Non-responder			Responder			Non-responder			Responder			Allele		Genotype	
	Samples	Percent		Samples	Percent		Samples	Percent		Samples	Percent		Chi square	p value	Chi square	p value
rs6805251	A	65	44.5	A	72	53.7	AA	14	19.2	AA	22	32.8	2.37	0.123	3.41	0.181
	C	81	55.5	C	62	46.3	AC	37	50.7	AC	28	41.8				
rs4688043	A	133	91.1	A	120	89.6	AA	61	83.6	AA	54	80.6	0.19	0.662	0.21	0.899
	G	13	8.9	G	14	10.4	AG	11	15.1	AG	12	17.9				
rs7624540	A	28	19.2	A	27	20.1	AA	2	2.7	AA	2	3.0	0.04	0.838	0.04	0.978
	G	118	80.8	G	107	79.9	AG	24	32.9	AG	23	31.3				
rs13319151	A	1	0.7	A	3	2.2	AA	0	0.0	AA	0	0.0	1.20	0.370	1.21	0.270
	G	145	99.3	G	131	97.8	AG	1	1.4	AG	3	4.5				
rs6438552	A	82	56.2	A	63	47.0	AA	22	30.1	AA	17	25.4	2.34	0.125	3.48	0.175
	G	64	43.8	G	71	53.0	AG	38	52.1	AG	29	43.3				
rs4072520	A	30	20.5	A	33	24.6	AA	2	2.7	AA	4	6.0	0.67	0.414	1.02	0.600
	C	116	79.5	C	101	75.4	AC	26	35.6	AC	25	37.3				
rs9878473	A	28	19.2	A	27	20.1	AA	22	30.1	AA	17	25.4	2.34	0.125	3.48	0.175
	T	118	80.8	T	107	79.9	AT	38	52.1	AT	29	43.3				
rs4491944	A	1	0.7	A	3	2.2	AA	0	0.0	AA	0	0.0	1.20	0.370	1.21	0.270
	T	145	99.3	T	131	97.8	AT	1	1.4	AT	3	4.5				
rs6772172	A	145	99.3	A	131	97.8	AA	72	98.6	AA	64	95.5	1.20	0.370	1.21	0.270
	G	1	0.07	G	1	2.2	AG	1	1.4	AG	3	4.5				
rs11919783	A	13	8.9	A	9	6.7	AA	0	0.0	AA	1	1.5	0.46	0.496	2.55	0.279
	G	133	91.1	G	125	93.3	AG	13	17.8	AG	7	10.4				
rs11923196	A	2	1.4	A	2	1.5	AA	0	0.0	AA	0	0.0	0.01	0.999	0.01	0.931
	T	144	98.6	T	132	98.5	AT	2	2.7	AT	2	3.0				
rs6779828	A	30	20.8	A	34	25.4	AA	2	2.8	AA	4	6.0	0.81	0.368	1.09	0.579
	G	114	79.2	G	100	74.6	AG	26	36.1	AG	26	38.8				
rs9846422	A	145	99.3	A	131	97.8	AA	72	98.6	AA	64	95.5	1.20	0.370	1.21	0.270
	G	1	0.07	G	1	2.2	AG	1	1.4	AG	3	4.5				
rs3755557	A	125	85.6	A	117	87.3	AA	53	72.6	AA	51	76.1	0.17	0.678	0.25	0.881
	C	21	14.4	C	17	12.7	AC	19	26.0	AC	15	22.4				
						CC	1	1.4	CC	1	1.5					

phrenia, resulting in increased GSK3 β activity. If GSK3 β regulation is altered in schizophrenia, it is possible that psychoactive compounds might also alter GSK3 signaling. Recent data suggest that antipsychotics have the opposite effect on Akt and GSK3 seen in schizophrenia, and this could be part of the mechanism of their therapeutic effects. In animal models, chronic antipsychotic treatment may increase the total levels of GSK3 (Alimohamad et al. 2005a, b) or Akt-induced serine phosphorylation of GSK3 (Alimohamad et al. 2005a; Emamian et al. 2004). Drugs that induce psychosis might also affect GSK3 signaling. Svenningsson et al. (2003) compared the short-term (15 min) effects of three psychomimetic drugs: amphetamine, D-lysergic acid (LSD; a serotonin agonist), and phencyclidine (PCP; a glutamate receptor antagonist) and found that in mice, all three increased pSer9-GSK3 β in the frontal cortex and striatum but not in the hippocampus or cerebellum.

Recent evidence indicates that the expression and phosphorylation of GSK3 can be regulated by neurotransmitter receptors such as dopamine D2 and serotonin 5HT1/5HT2. Acute stimulation of dopamine D2 receptors elicits dephosphorylation and activation of GSK3 (Beaulieu et al. 2004, 2005). D2 blockade-mediated phosphorylation of GSK3 was accompanied with the activation of Akt, suggesting that Akt may be an upstream regulator of GSK3 in D2 receptor-mediated signaling (Beaulieu et al. 2007). On the other hand, the acute stimulation of serotonin 5HT1 receptors or blockade of 5HT2 receptors produces phosphorylation-induced inhibition of GSK3 in the rodent brain (Li et al. 2004). However, the mediators of the 5HT receptor-coupled regulation of GSK3 have not been well established; a recent report stated that they were not mediated by Akt (Li et al. 2007). The first report of a possible involvement of GSK3 in schizophrenia was by Yang et al. (1995), where they showed that both cellular activity and protein levels of kinase FA/GSK3 α in the lymphocytes of schizophrenia patients were much lower than in normal controls.

We did not observe any association between GSK3 β genetic variants and clozapine response. The finding that GSK3 is inhibited by both haloperidol and clozapine suggests that GSK3 may be a common target of both typical and atypical antipsychotics. In vivo results suggest that clozapine increases pSer9-GSK3 β and pSer473-Akt (Kozlovsky et al. 2006; Roh et al. 2007), although Kang et al. (2004) did not find this effect in cultured cells SH-SY5Y that do not express D2 or 5HT2 receptors. Because of the links between GSK3 and both the pathophysiology of psychotic symptoms and antipsychotic medication effects (Jope and Roh 2006), further studies are required to understand how the associated markers reported here may affect these processes. Studies in larger, independent samples are also required to strengthen our findings.

Acknowledgments Funding and grants were provided by (1) CNPq—Brazil (#202447/2006-5; #140950/2005-2; #554496/2005-4), (2) National Institutes of Health (NIH), (3) Canadian Institutes of Health Research (CIHR) #940595, (4) Ontario Mental Health Foundation (OMHF), and (5) Fapemig—Brazil. The authors would like to thank Daniela VF Rosa, Mary Smirniw, Nicole King and Arun Tiwari for their help and support throughout the manuscript preparation.

Disclosure/conflicts of interest Mr. Souza, Dr. Romano-Silva, and Dr. Wong have nothing to declare. Dr. Lieberman has served as a consultant/advisor or grantee of Acadia, Astra Zeneca, Bristol-Myers Squibb, Eli Lilly, GlaxoSmithKline, Janssen Pharmaceutica, Lundbeck, Merck, Organon, Pfizer, and Wyeth and holds a patent from Repligen. Dr. Meltzer declares that he is a consultant or grantee of Abbott, Acadia, ARYx, Astra Zeneca, Bristol-Myers Squibb, Eli Lilly, Janssen, Memory, Minster, Organon, Pfizer, Solvay, Wyeth, and Vanda. Dr. Kennedy declares that he is a consultant for GlaxoSmithKline.

References

- Aickin M (1999) Other method for adjustment of multiple testing exists. *BMJ* 318:127–128
- Alimohamad H, Rajakumar N, Seah YH, Rushlow W (2005a) Antipsychotics alter the protein expression levels of beta-catenin and GSK-3 in the rat medial prefrontal cortex and striatum. *Biol Psychiatry* 57:533–542
- Alimohamad H, Sutton L, Mouyal J, Rajakumar N, Rushlow WJ (2005b) The effects of antipsychotics on beta-catenin, glycogen synthase kinase-3 and dishevelled in the ventral midbrain of rats. *J Neurochem* 95:513–525
- Ananth J, Parameswaran S, Gunatilake S (2004) Side effects of atypical antipsychotic drugs. *Curr Pharm Des* 10:2219–2229
- Balding DJ (2006) A tutorial on statistical methods for population association studies. *Nat Rev Genet* 7:781–791
- Barrett JC, Fry B, Maller J, Daly MJ (2005) Haploview: analysis and visualization of LD and haplotype maps. *Bioinformatics* 21:263–265
- Basile VS, Masellis M, Potkin SG, Kennedy JL (2002) Pharmacogenomics in schizophrenia: the quest for individualized therapy. *Hum Mol Genet* 11:2517–2530
- Beasley C, Cotter D, Khan N, Pollard C, Sheppard P, Varndell I, Lovestone S, Anderton B, Everall I (2001) Glycogen synthase kinase-3beta immunoreactivity is reduced in the prefrontal cortex in schizophrenia. *Neurosci Lett* 302:117–120
- Beaulieu JM, Sotnikova TD, Marion S, Lefkowitz RJ, Gainetdinov RR, Caron MG (2005) An Akt/beta-arrestin 2/PP2A signaling complex mediates dopaminergic neurotransmission and behavior. *Cell* 122:261–273
- Beaulieu JM, Sotnikova TD, Yao WD, Kockeritz L, Woodgett JR, Gainetdinov RR, Caron MG (2004) Lithium antagonizes dopamine-dependent behaviors mediated by an AKT/glycogen synthase kinase 3 signaling cascade. *Proc Natl Acad Sci U S A* 101:5099–5104
- Beaulieu JM, Tirota E, Sotnikova TD, Masri B, Salahpour A, Gainetdinov RR, Borrelli E, Caron MG (2007) Regulation of Akt signaling by D2 and D3 dopamine receptors in vivo. *J Neurosci* 27:881–885
- Bender R, Lange S (1999) Multiple test procedures other than Bonferroni's deserve wider use. *BMJ* 318:600–601
- Cotter D, Kerwin R, Al-Sarraj S, Brion JP, Chadwick A, Lovestone S, Anderton B, Everall I (1998) Abnormalities of Wnt signalling in schizophrenia—evidence for neurodevelopmental abnormality. *Neuroreport* 9:1379–1383

- Dudbridge F (2003) Pedigree disequilibrium tests for multilocus haplotypes. *Genet Epidemiol* 25:115–121
- Durrant C, Zondervan KT, Cardon LR, Hunt S, Deloukas P, Morris AP (2004) Linkage disequilibrium mapping via cladistic analysis of single-nucleotide polymorphism haplotypes. *Am J Hum Genet* 75:35–43
- Emamian ES, Hall D, Bimbaum MJ, Karayiorgou M, Gogos JA (2004) Convergent evidence for impaired AKT1-GSK3beta signaling in schizophrenia. *Nat Genet* 36:131–137
- Feder JN, Gnirke A, Thomas W, Tsuchihashi Z, Ruddy DA, Basava A, Dormishian F, Domingo R Jr, Ellis MC, Fullan A, Hinton LM, Jones NL, Kimmel BE, Kronmal GS, Lauer P, Lee VK, Loeb DB, Mapa FA, McClelland E, Meyer NC, Mintier GA, Moeller N, Moore T, Morikang E, Prass CE, Quintana L, Starnes SM, Schatzman RC, Brunke KJ, Drayna DT, Risch NJ, Bacon BR, Wolff RK (1996) A novel MHC class I-like gene is mutated in patients with hereditary haemochromatosis. *Nat Genet* 13:399–408
- Gabriel SB, Schaffner SF, Nguyen H, Moore JM, Roy J, Blumenstiel B, Higgins J, DeFelice M, Lochner A, Faggart M, Liu-Cordero SN, Rotimi C, Adeyemo A, Cooper R, Ward R, Lander ES, Daly MJ, Altshuler D (2002) The structure of haplotype blocks in the human genome. *Science* 296:2225–2229
- Guo SW, Thompson EA (1992) Performing the exact test of Hardy-Weinberg proportion for multiple alleles. *Biometrics* 48:361–372
- Hao K, Xu X, Laird N, Wang X, Xu X (2004) Power estimation of multiple SNP association test of case-control study and application. *Genet Epidemiol* 26:22–30
- Hoh J, Wille A, Ott J (2001) Trimming, weighting, and grouping SNPs in human case-control association studies. *Genome Res* 11:2115–2119
- Hwang R, Shinkai T, De LV, Muller DJ, Ni X, Macciardi F, Potkin S, Lieberman JA, Meltzer HY, Kennedy JL (2005) Association study of 12 polymorphisms spanning the dopamine D(2) receptor gene and clozapine treatment response in two treatment refractory/intolerant populations. *Psychopharmacology (Berl)* 181:179–187
- Ikedo M, Iwata N, Suzuki T, Kitajima T, Yamanouchi Y, Kinoshita Y, Ozaki N (2005) No association of GSK3beta gene (GSK3B) with Japanese schizophrenia. *Am J Med Genet B Neuropsychiatr Genet* 134:90–92
- Jope RS, Johnson GV (2004) The glamour and gloom of glycogen synthase kinase-3. *Trends Biochem Sci* 29:95–102
- Jope RS, Roh MS (2006) Glycogen synthase kinase-3 (GSK3) in psychiatric diseases and therapeutic interventions. *Curr Drug Targets* 7:1421–1434
- Jope RS, Yuskaitis CJ, Beurel E (2007) Glycogen synthase kinase-3 (GSK3): inflammation, diseases, and therapeutics. *Neurochem Res* 32:577–595
- Kane J, Honigfeld G, Singer J, Meltzer H (1988) Clozapine for the treatment-resistant schizophrenic. A double-blind comparison with chlorpromazine. *Arch Gen Psychiatry* 45:789–796
- Kang UG, Seo MS, Roh MS, Kim Y, Yoon SC, Kim YS (2004) The effects of clozapine on the GSK-3-mediated signaling pathway. *FEBS Lett* 560:115–119
- Kapur S, Seeman P (2001) Does fast dissociation from the dopamine d (2) receptor explain the action of atypical antipsychotics?: a new hypothesis. *Am J Psychiatry* 158:360–369
- Kozlovsky N, Amar S, Belmaker RH, Agam G (2006) Psychotropic drugs affect Ser9-phosphorylated GSK-3 beta protein levels in rodent frontal cortex. *Int J Neuropsychopharmacol* 9:337–342
- Kozlovsky N, Belmaker RH, Agam G (2000) Low GSK-3beta immunoreactivity in postmortem frontal cortex of schizophrenic patients. *Am J Psychiatry* 157:831–833
- Kozlovsky N, Belmaker RH, Agam G (2001) Low GSK-3 activity in frontal cortex of schizophrenic patients. *Schizophr Res* 52:101–105
- Kozlovsky N, Shanon-Weickert C, Tomaskovic-Crook E, Kleinman JE, Belmaker RH, Agam G (2004) Reduced GSK-3beta mRNA levels in postmortem dorsolateral prefrontal cortex of schizophrenic patients. *J Neural Transm* 111:1583–1592
- Lahiri DK, Nurnberger JI (1991) A rapid non-enzymatic method for the preparation of HMW DNA from blood for RFLP studies. *Nucleic Acids Res* 19:5444
- Laird NM, Horvath S, Xu X (2000) Implementing a unified approach to family-based tests of association. *Genet Epidemiol* 19(Suppl 1):S36–S42
- Lee WC (2003) Searching for disease-susceptibility loci by testing for Hardy-Weinberg disequilibrium in a gene bank of affected individuals. *Am J Epidemiol* 158:397–400
- Lee KY, Ahn YM, Joo EJ, Jeong SH, Chang JS, Kim SC, Kim YS (2006) No association of two common SNPs at position -1727 A/T, -50 C/T of GSK-3 beta polymorphisms with schizophrenia and bipolar disorder of Korean population. *Neurosci Lett* 395:175–178
- Li X, Zhu W, Roh MS, Friedman AB, Rosborough K, Jope RS (2004) In vivo regulation of glycogen synthase kinase-3beta (GSK3beta) by serotonergic activity in mouse brain. *Neuropsychopharmacology* 29:1426–1431
- Li X, Rosborough KM, Friedman AB, Zhu W, Roth KA (2007) Regulation of mouse brain glycogen synthase kinase-3 by atypical antipsychotics. *Int J Neuropsychopharmacol* 10:7–19
- Luo X, Kranzler HR, Zuo L, Lappalainen J, Yang BZ, Gelernter J (2006) ADH4 gene variation is associated with alcohol dependence and drug dependence in European Americans: results from HWD tests and case-control association studies. *Neuropsychopharmacology* 31:1085–1095
- Luo X, Kranzler HR, Zuo L, Wang S, Blumberg HP, Gelernter J (2005) CHRM2 gene predisposes to alcohol, drug dependence and affective disorder: results from an extended population structured association study. *Hum Mol Genet* 14:2421–2434
- Malhotra AK, Litman RE, Pickar D (1993) Adverse effects of antipsychotic drugs. *Drug Saf* 9:429–436
- Malhotra AK, Murphy GM, Kennedy JL (2004) Pharmacogenetics of psychotropic drug response. *Am J Psychiatry* 161:780–796
- Meltzer HY (1989) Clinical studies in the mechanism of action of clozapine: the dopamine serotonin hypothesis of schizophrenia. *Psychopharmacology* 99:518–527
- Nadri C, Lipska BK, Kozlovsky N, Weinberger DR, Belmaker RH, Agam G (2003) Glycogen synthase kinase (GSK)-3beta levels and activity in a neurodevelopmental rat model of schizophrenia. *Brain Res Dev Brain Res* 141:33–37
- Nadri C, Dean B, Scarr E, Agam G (2004) GSK-3 parameters in postmortem frontal cortex and hippocampus of schizophrenic patients. *Schizophr Res* 71:377–382
- Nielsen DM, Ehm MG, Weir BS (1998) Detecting marker-disease association by testing for Hardy-Weinberg disequilibrium at a marker locus. *Am J Hum Genet* 63:1531–1540
- Nyholt DR (2004) A simple correction for multiple testing for single-nucleotide polymorphisms in linkage disequilibrium with each other. *Am J Hum Genet* 74:765–769
- Pemeger TV (1998) What's wrong with Bonferroni adjustments. *BMJ* 316:1236–1238
- Purcell S, Cherny SS, Sham PC (2003) Genetic Power Calculator: design of linkage and association genetic mapping studies of complex traits. *Bioinformatics* 19:149–150
- Roh MS, Seo MS, Kim Y, Kim SH, Jeon WJ, Ahn YM, Kang UG, Juhn YS, Kim YS (2007) Haloperidol and clozapine differentially regulate signals upstream of glycogen synthase kinase 3 in the rat frontal cortex. *Exp Mol Med* 39:353–360
- Scassellati C, Bonvicini C, Perez J, Bocchio-Chiavetto L, Tura GB, Rossi G, Racagni G, Gennarelli M (2004) Association study of -1727 A/T, -50 C/T and (CAA)_n repeat GSK-3beta gene polymorphisms with schizophrenia. *Neuropsychobiology* 50:16–20

- Seeman P (2002) Atypical antipsychotics: mechanism of action. *Can J Psychiatry* 47:27–38
- Seeman P, Lee T (1975) Antipsychotic drugs: direct correlation between clinical potency and presynaptic action on dopamine neurons. *Science* 188:1217–1219
- Shin SY, Yoon SC, Kim YH, Kim YS, Lee YH (2002) Phosphorylation of glycogen synthase kinase-3beta at serine-9 by phospholipase Cgamma1 through protein kinase C in rat 3Y1 fibroblasts. *Exp Mol Med* 34:444–450
- Sutherland C, Leighton IA, Cohen P (1993) Inactivation of glycogen synthase kinase-3 beta by phosphorylation: new kinase connections in insulin and growth-factor signalling. *Biochem J* 296(Pt 1):15–19
- Svenningsson P, Tzavara ET, Carruthers R, Rachleff I, Wattler S, Nehls M, McKinzie DL, Fienberg AA, Nomikos GG, Greengard P (2003) Diverse psychotomimetics act through a common signaling pathway. *Science* 302:1412–1415
- Szczepankiewicz A, Skibinska M, Hauser J, Slopian A, Leszczynska-Rodziewicz A, Kapelski P, Dmitrzak-Weglaz M, Czerski PM, Rybakowski JK (2006) Association analysis of the GSK-3beta T-50C gene polymorphism with schizophrenia and bipolar disorder. *Neuropsychobiology* 53:51–56
- Trikalinos TA, Salanti G, Khoury MJ, Ioannidis JP (2006) Impact of violations and deviations in Hardy–Weinberg equilibrium on postulated gene–disease associations. *Am J Epidemiol* 163:300–309
- Wigginton JE, Abecasis GR (2005) PEDSTATS: descriptive statistics, graphics and quality assessment for gene mapping data. *Bioinformatics* 21:3445–3447
- Wigginton JE, Cutler DJ, Abecasis GR (2005) A note on exact tests of Hardy–Weinberg equilibrium. *Am J Hum Genet* 76:887–893
- Witke-Thompson JK, Pluzhnikov A, Cox NJ (2005) Rational inferences about departures from Hardy–Weinberg equilibrium. *Am J Hum Genet* 76:967–986
- Woodgett JR (1990) Molecular cloning and expression of glycogen synthase kinase-3/factor A. *EMBO J* 9:2431–2438
- Yang SD, Yu JS, Lee TT, Yang CC, Ni MH, Yang YY (1995) Dysfunction of protein kinase FA/GSK-3 alpha in lymphocytes of patients with schizophrenic disorder. *J Cell Biochem* 59:108–116

3.2 - Genetic association analysis of the GFR alpha genes with schizophrenia and clozapine response

Genetic association analysis of the GFR alpha genes with schizophrenia and clozapine response

Renan P. Souza^{a,b,c}, Marco A. Romano-Silva^{a,b}, Jeffrey A. Lieberman^e, Herbert Y Meltzer^f,
Leslie MacNeil^g, Joseph G. Culotti^g, James L. Kennedy^{c,d}, Albert H.C. Wong^{c,d}.

^aGrupo de Pesquisa em Neuropsiquiatria Clínica e Molecular, UFMG, Belo Horizonte, Brazil; ^bLaboratório de Neurociência, Dept. Saúde Mental, Faculdade de Medicina, UFMG, Brazil; ^cNeurogenetics Section, CAMH, Toronto, ON, Canada ^dDepartment of Psychiatry, University of Toronto, ON, Canada; ^eDepartment of Psychiatry, University of North Carolina, Chapel Hill, NC, USA; ^fPsychiatric Hospital, Vanderbilt University, Nashville, TN, USA; ^gSamuel Lunenfeld Research Institute, Mount Sinai Hospital, Toronto, ON, Canada.

Abstract

GDNF (glial-cell-line derived neurotrophic factor) is a potent neurotrophic factor for dopaminergic neurons. Neuropsychiatric diseases and their treatments are associated with alterations in the levels of both GDNF and its receptor family (GDNF family receptor alpha or GFRA). *GFRA1*, *GFRA2* and *GFRA3* are located in chromosomal regions with suggestive linkage to schizophrenia. In this study we analyzed polymorphisms located in all four known GFRA genes and examined association with schizophrenia and clozapine response. We examined SNPs across the genes *GFRA1 - 4* in 219 matched case-control subjects, 85 small nuclear families and 140 schizophrenia patients taking clozapine for 6 months. *GFRA1* rs11197557 was associated with schizophrenia; *GFRA1* rs730357 and some haplotypes showed a significantly different transmission pattern, and two haplotypes (rs11197612-rs3781514 and rs12413585-rs730057-rs1197612) were associated with clozapine response. In *GFRA2*, three individual SNPs (rs1128397, rs13250096 and rs4567028) and several haplotypes showed association with response. *GFRA3* rs11242417 SNP and a haplotype containing all markers analyzed were associated with schizophrenia. None of the *GFRA4* markers evaluated had a significant association. We also found evidence for interactions between *GFRA1*, 2 and 3 associated with schizophrenia and clozapine response, consistent with the locations of these three genes within linkage regions for schizophrenia. *GFRA4*, which is not located in such a region, showed no interactions with the other genes in this regard. Our results presented nominally significant evidence that the GFRA genes may affect susceptibility to schizophrenia and response to clozapine treatment.

Keywords: schizophrenia, clozapine response. GDNF, GFR alpha, genetic association, family-based association test.

1 - Introduction

Schizophrenia is a serious, complex genetic neuropsychiatric disorder with a life-time prevalence of 0.5-1% in the population [1]. Family, twin and adoption studies convincingly demonstrate that relatives of affected individuals have a higher risk for schizophrenia and that this is largely the result of genetic factors [2]. A variety of different genes, each with small or moderate effect, are thought to be involved in the etiology of schizophrenia and the strongest findings to date include neuregulin-1, dysbindin and disrupted-in-schizophrenia-1 [3-6]. These genes share several important features: genetic association of specific SNPs and haplotypes with schizophrenia, chromosomal localization within linkage regions for schizophrenia and evidence for modulating neurodevelopment [7, 8].

It has been postulated that targeting the synthesis and secretion of neurotrophic factors, such as nerve growth factor (NGF), brain-derived neurotrophic factor (BDNF) and GDNF, might be a new approach to treating neurodegenerative and depressive disorders [9, 10], but this possibility has only recently been considered for antipsychotic drugs [11-13]. NGF and BDNF have been implicated in the neuroprotective actions of antipsychotic drugs [14, 15]. GDNF is a distantly-related member of the transforming growth factor-beta (TGF-beta) family that was isolated from a glial cell line [16]. Other members of the GDNF family were subsequently identified as neurturin, persephin and artemin. GDNF is synthesized in striatal cells and undergoes retrograde transport to dopaminergic cell bodies in the midbrain [17, 18]. GDNF enhances the survival of dopaminergic neurons [17, 19-21] and it has been postulated as the most potent neurotrophic factor for dopaminergic neurons [19].

GFRA proteins are non-signaling extracellular molecules that act as co-receptors for the binding of GDNF family proteins to the RET receptor. GFRA proteins are glycoproteins anchored to the cell surface by a C-terminal glycosylphosphatidylinositol-linkage [23]. Four GFRA family members have been recognized with similar structures and 30%–45% sequence identity. However each has a distinct expression pattern and affinity for GDNF-family ligands. GFRA1 has high affinity for GDNF and RET [24, 25], GFRA2 with neurturin [26, 27], GFRA3 with artemin [28] and GFRA4 with persephin [29, 30]. *GFRA1*, *GFRA2* and *GFRA3* are located within chromosomal regions that have been reported to be in linkage with schizophrenia [7]. Furthermore, TGF-beta genes are crucial regulators of neuron migration in *C. elegans* [31-33], a major component of cortical neurodevelopment that is hypothesized to be abnormal in schizophrenia. It has been suggested that GDNF plays a role in mammalian neuronal development [8, 34]. Based on the above rationale, we hypothesized that the GFRA genes may affect susceptibility to schizophrenia.

Variation in individual clinical response to antipsychotic treatment remains a critical problem in the management of serious mental illness. Treatment often proceeds by trial and error in order to determine the medication and dose that maximizes response and minimizes toxicity. Although a minority of patients may experience complete remission, a large proportion of patients continue to experience significant symptoms. In addition, a subset of patients develops drug-induced adverse events that range from troublesome to life-threatening. In spite of the wide array of medicines available, 10–20% of patients do not respond to treatment with an antipsychotic. An additional 20–30% who do respond early on eventually relapse and some develop serious side effects that cause them to discontinue medication [35, 36].

Some reports have shown GDNF system changes after administration of psychotropic drugs. Antidepressant and atypical antipsychotic drugs, but not typical drugs haloperidol, have been reported to increase GDNF release from C6 cells [37, 38]. GDNF levels are also altered by lithium and valproic acid administration [39, 40]. Furthermore, electroconvulsive seizure increased *GFRA1* and 2 mRNA levels in hippocampus; although *GDNF* and *c-ret* mRNAs were not significantly changed [41]. Given that this collection of psychiatric treatments affect components of the GDNF system, we hypothesized that genetic variation in the GFRA genes could affect response to clozapine treatment in schizophrenia.

2 - Methods

Clinical sample

All recruitment and clinical assessments were conducted with written informed consent, with the explicit approval of our institutional ethics review board and in accordance to Declaration of Helsinki. Clinical data and DNA samples were obtained from the probands of 85 small nuclear families, as well as 219 patients with a DSM-III-R or DSM-IV diagnosis of schizophrenia. The healthy controls (N=219) were matched for age (± 5 years), sex, and ethnicity (146 male and 73 female cases and the same number of controls, mean age 36 ± 8). The Structured Clinical Interview for DSM-IV Axis I Disorders (SCID-I) was administered by trained research assistants to each patient and diagnosis was supplemented by a review of medical records. The diagnosis was established via consensus procedures incorporating two of the investigators. The controls were screened for current or past history of major psychiatric disorders or substance misuse, and excluded if either was detected.

For the clozapine response sample, clinical data from 140 patients with a DSM-III-R or DSM-IV diagnosis of schizophrenia, almost all of whom were treatment refractory or intolerant of typical antipsychotic therapy [42], were obtained at the two following research clinics: Case Western Reserve University in Cleveland, OH (n=90) (63 males and 27 females in each group, mean age 36 ± 8) and Hillside Hospital in Glen Oaks, NY (n=50) (33 males and 17 females in each group, mean age 35 ± 8). After informed consent was obtained, patients underwent a washout period of 2 to 4 weeks during which, unless clinically necessary, they received no medications before starting clozapine. Clozapine treatment was continued for a minimum of 6 months during which patients were evaluated prospectively. Clozapine blood levels were monitored throughout the course of treatment to ascertain compliance. Treatment response was evaluated as the percentage score change on the 18 item Brief Psychiatric Rating Scale (BPRS). Treatment response was expressed as a dichotomous variable in the whole sample at 6 months using criteria based on those of Kane [42]: a reduction of $\geq 20\%$ on the overall score of the BPRS from the baseline score at enrolment. There were no differences observed between the sites in terms of gender ratio, mean age, mean age of onset or response ratio. Caucasians and African-American subjects were not significantly different in terms of gender ratio, mean age, mean age of onset or response ratio (data not shown) [43].

Genetic analyses

Genomic DNA was extracted using the high salt method. We analyzed 26 SNPs in the *GFRA1* (rs1078080, rs11598215, rs3781514, rs2694783, rs2694801, rs3824840, rs12776813, rs7085306, rs10787627, rs9787429, rs12775655, rs11197557, rs10749189, rs11197567, rs3781539, rs7903297, rs17094340, rs7920934, rs10885877, rs4751956, rs11812459, rs10885888, rs12413585, rs730357, rs11197612), 17 in *GFRA2* (rs15881, rs4567027,

rs7813735, rs10088105, rs1128397, rs6988470, rs4237073, rs10283397, rs4739217, rs6587002, rs7014143, rs4567028, rs4739286, rs11993990, rs4078157, rs4739285, rs13250096), 4 in *GFRA3* (rs10036665, rs10952, rs11242417, rs7726580) and 2 in *GFRA4* (rs633924, rs6084432). Genotyping was performed by GoldenGate assay (Illumina, San Diego, CA, USA) at The Centre for Applied Genomics (TCAG) at the Hospital for Sick Children in Toronto, ON, Canada. Quality check procedure has been performed in this sample as described at [44].

Statistical analyses

Individual SNP analyses of clozapine responder (case) and non-responder (control) data and Hardy–Weinberg equilibrium assessment were performed using χ^2 tests. The statistical program used was the Statistical Package for the Social Sciences, version 10.0.7 for genotypic association and Haploview 4.0 [45] for allelic association. We applied the family-based association test (FBAT, version 1.0 [46]) under the assumption of an additive model, and PEDSTATS [47] for Hardy–Weinberg equilibrium in the family data. Linkage disequilibrium (LD) was assessed using Haploview, version 4.0. Haplotype analyses were performed using UNPHASED 3.0.10 [48], Haploview version 4.0 and FBAT. Gene–gene interactions were examined using the multifactor dimensionality reduction (MDR) method version 1.1.0 [49-51]. A detailed explanation of MDR has been published elsewhere [52].

3 - Results

Linkage disequilibrium analysis

Case-control sample: Pairwise LD between the SNPs is presented for each gene (Figure S1). In this study, we defined a haplotype block as a region over which < 5% of pairwise comparisons among informative SNPs showed strong evidence of historical recombination (upper confidence bound on D' less than 0.9; [53]). Based on this definition we found 6 LD blocks in *GFRA1*, 5 in *GFRA2* and 1 block in each of *GFRA3* and *GFRA4*.

Family sample: Following the same criteria used for case-control sample, *GFRA1* showed 6 blocks, 3 blocks in *GFRA2*, and 1 block in each of *GFRA3* and *GFRA4* (FigureS2).

Response sample: we observed 6 LD blocks in *GFRA1*, 4 in *GFRA2* and 1 block in *GFRA4* (FigureS3). No blocks were observed in *GFRA3*.

Single-marker analysis

Case-control sample: Significant deviations from Hardy–Weinberg equilibrium were observed for *GFRA1* rs3824840 ($p = 0.043$), *GFRA1* rs9787429 ($p = 0.032$), *GFRA1* rs10749189 ($p = 0.030$) and *GFRA2* rs4739217 ($p = 0.008$) in the control group. The following markers deviated from the Hardy-Weinberg equilibrium in cases: *GFRA1* rs1078080 ($p = 0.016$), *GFRA1* rs10885888 ($p = 0.004$), *GFRA4* rs633924 ($p = 0.007$) and *GFRA4* rs6084432 ($p = 0.041$). Only one SNP in the controls and two in the cases showed deviation from Hardy-Weinberg equilibrium at a significance threshold of $p < 0.01$. The cases and controls were compared for genotype and allele frequencies across the markers (see Table S1 and Figure 3). In our population, no significant differences in genotype or allele frequency were found between cases and controls for any SNPs in *GFRA2* and *GFRA4*. *GFRA1* rs11197557 (allele $p = 0.020$, $X^2 = 5.34$; genotype $p = 0.017$, $X^2 = 8.09$) and *GFRA3* rs11242417 (allele $p = 0.010$, $X^2 = 6.56$;

genotype $p = 0.014$, $X^2 = 8.50$) were associated with schizophrenia. All markers that showed significant association were in Hardy-Weinberg equilibrium.

Family sample: Allele frequencies are presented in Table S2. *GFRA1* rs730357 A allele was overtransmitted to patients with schizophrenia (allele frequency = 0.742, $z = 2.112$, $p = 0.035$).

Response sample: Significant deviation from Hardy-Weinberg equilibrium was observed for *GFRA2* rs4739217 ($p = 0.008$) in the non-responder group. The following markers deviated significantly from Hardy-Weinberg equilibrium in the responder group: *GFRA1* rs3824840 ($p = 0.030$), *GFRA4* rs6084432 ($p = 0.041$) and *GFRA4* rs633924 ($p = 0.007$). Responder/non-responder groups were compared for genotype and allele frequencies across the markers (see Table S3 and Figure 4). In our population, no significant differences in genotype or allele frequency were seen between responders and non-responders for any SNPs in *GFRA1*, *GFRA3* and *GFRA4*. The following *GFRA2* SNPs showed significant association with treatment response: rs1128397 (allele $p = 0.009$, $X^2 = 6.70$; genotype $p = 0.022$, $X^2 = 7.59$); rs13250096 (allele $p = 0.019$, $X^2 = 5.50$; genotype $p = 0.064$, $X^2 = 5.51$) and rs4567028 (allele $p = 0.047$, $X^2 = 3.92$; genotype $p = 0.068$, $X^2 = 5.39$). All markers that showed significant association were in Hardy-Weinberg equilibrium.

Haplotype analysis

Case-control sample: Cases and controls were compared for haplotype frequencies across the GFRA markers. A small marker size (two and three markers) was chosen since high levels of haplotype diversity were expected due to the moderate LD observed in our samples [54]. *GFRA1*, *GFRA2* and *GFRA4* genes did not show any significant haplotypic association with two or three marker windows or haplotypes located in the same LD block. Considering haplotypes in the same LD block, the haplotype within *GFRA3* block 1 (rs10036665, rs10952, rs11242417 and rs7726580) T-T-C-G showed significant association with schizophrenia (control frequency = 0.173, case frequency = 0.109, $p = 0.006$; $X^2 = 7.33$; after 1 000 permutations $p = 0.027$).

Family sample: Analyzing haplotypes composed of *GFRA1* SNPs that reached lowest p -values (rs1078080 $p = 0.113$; rs7920934 $p = 0.053$ and rs730357 $p = 0.035$) the rs1078080-rs7920934 A-A haplotype was overtransmitted (frequency = 0.677, $z = 2.166$, $p = 0.030$; after 1 000 permutations $p = 0.111$); rs7920934-rs730357 G-G (frequency = 0.119, $z = -2.361$, $p = 0.018$; after 1,000 permutations $p = 0.043$) and rs1078080-rs7920934 G-G (frequency = 0.045, $z = -2.614$, $p = 0.008$; after 1,000 permutations $p = 0.026$) were undertransmitted. Considering haplotypes in the same LD block, inside *GFRA1* block 5 (rs7920934, rs10885877 and rs4751956) the G-G-G haplotype was undertransmitted to patients with schizophrenia (frequency = 0.223, transmitted: untransmitted ratio = 9 : 21, $p = 0.028$, $X^2 = 4.80$; after 1,000 permutations $p = 0.290$). In addition, in *GFRA1* block 6 (rs12413585, rs730357 and rs11197612) the G-A-G (frequency = 0.258; transmitted: untransmitted ratio = 23 : 11, $p = 0.028$, $X^2 = 4.82$, after 1,000 permutations $p = 0.284$) and G-G-G haplotypes showed significant different transmission pattern (frequency = 0.247, transmitted: untransmitted ratio = 9 : 22, $p = 0.019$, $X^2 = 5.45$; after 1 000 permutations $p = 0.186$). *GFRA2*, *GFRA3* and *GFRA4* genes did not show any significant haplotypic association with two or three marker windows or haplotypes located in the same LD block.

Response sample: Clozapine responders and non-responders were compared for haplotype frequencies across the GFRA markers. *GFRA1* markers showed significant haplotypic

association for rs11197612-rs3781514 (global $p = 0.044$, $X^2 = 8.09$). Considering haplotypes in the same LD block, inside *GFRA1* block 6 (rs12413585, rs730057 and rs1197612) the haplotype G-A-G showed association (non-responder frequency = 0.208, responder frequency = 0.330, $p = 0.021$, $X^2 = 5.30$; after 1,000 permutations $p = 0.289$). Two window *GFRA2* marker analysis found associations, the strongest of which consisted of rs1128397 and rs13250096 (global $p = 0.005$, $X^2 = 12.6$). The A-C (non-responder frequency = 0.073, responder frequency = 0.010, $p = 0.0002$, $X^2 = 13.66$) and T-G (non-responder frequency = 0.073, responder frequency = 0.010, $p = 0.0002$, $X^2 = 13.66$) haplotypes were associated with better response in our sample. The strongest *GFRA2* three-maker haplotype association was with rs1128397, rs4567028 and rs13250096 (global $p = 0.012$, $X^2 = 16.2$). The A-A-C (non-responder frequency = 0.070, responder frequency = 0.011, $p = 0.001$, $X^2 = 10.55$) and T-A-G (non-responder frequency = 0.348, responder frequency = 0.554, $p = 0.0006$, $X^2 = 11.59$) haplotypes were associated with better response in our sample. Considering haplotypes in the same LD block, the *GFRA2* block 1 (rs15881 and rs1128397) haplotype G-C showed association (non-responder frequency = 0.473, responder frequency = 0.321, $p = 0.009$, $X^2 = 6.70$; after 1 000 permutations $p = 0.093$). No haplotypic association was found in the *GFRA3* and *GFRA4*.

Gene-gene interactions

Case-control sample: Multi dimension reduction (MDR) analyses showed a significant association between combinations of the *GFRA* genes and schizophrenia. The best two-locus model contained *GFRA1* rs11197557 and *GFRA3* rs11242417 with a maximum cross-validation (CV) consistency of 10/10 and a maximum prediction accuracy of 56.6% ($p = 0.001$; after 1 000 permutations $p = 0.377$). The best three-locus model contained *GFRA1* rs11197557, *GFRA1* rs1078080 and *GFRA3* rs11242417, with a maximum CV consistency of 10/10 and a maximum prediction accuracy of 58.0% ($p = 0.001$; after 1 000 permutations $p = 0.054$). None of the interactions showed synergy (Figure S2).

Response sample: There were significant associations between gene-gene interactions and treatment response. The best two-locus model contained *GFRA1* rs1078080 and *GFRA2* rs15881 with a maximum cross-validation (CV) consistency of 4/10 and a maximum prediction accuracy of 46.6% ($p = 0.37$; after 1 000 permutations $p = 0.623$). The best three-locus model contained *GFRA1* rs10885888, *GFRA2* rs4237073 and *GFRA3* rs7726580, with a maximum CV consistency of 10/10 and a maximum prediction accuracy of 71.7% ($p = 0.001$; after 1 000 permutations $p = 0.054$). Likewise, the interaction dendrogram (Figure S2) placed *GFRA1* rs10885888, *GFRA2* rs4237073 and *GFRA3* rs7726580 on the same branch. Their position in the diagram indicates that this is the strongest interaction, which is consistent with location of these three genes within suggestive chromosomal linkage regions for schizophrenia. *GFRA4* is not located in a schizophrenia linkage region, and no interaction with the other genes was detected in this analysis.

4 - Discussion

This exploratory study examined the association of 26 SNPs in *GFRA1*, 17 in *GFRA2*, 4 in *GFRA3* and 2 in *GFRA4* and two phenotypes: diagnosis of schizophrenia, and clozapine response in schizophrenia patients. In the case-control analyses, the *GFRA1* rs11197557 and *GFRA3* rs11242417 markers showed nominally significant association with schizophrenia. One haplotype including *GFRA3* markers (rs10036665, rs10952, rs11242417, rs7726580) also showed association with schizophrenia. In the family-based sample, *GFRA1* gene showed allelic

and haplotypic association with schizophrenia. The rs730357 A allele; rs1078080-rs7920934 A-A and rs12413585-rs730357 -rs11197612 G-A-G haplotypes were overtransmitted. Haplotypes rs7920934-rs730357 G-G; rs1078080-rs7920934 G-G; rs7920934-rs10885877-rs4751956 G-G-G and rs12413585-rs730357 -rs11197612 G-G-G were undertransmitted. In the treatment response sample, *GFRA1* showed two protective haplotypes: rs11197612-rs3781514 and rs12413585-rs730057-rs1197612. *GFRA2* gene showed association with three individual SNPs (rs1128397, rs13250096 and rs4567028) and treatment response. Interestingly these polymorphisms did not show LD in the Haploview analysis, but some haplotypes showed association with treatment response, in particular a haplotype located in the first LD block (rs15881 and rs1128397). In *GFRA3* and *GFRA4* no genetic associations were observed.

All SNPs that presented allelic, genotypic or haplotypic nominal significant association were in Hardy-Weinberg equilibrium. We performed Bonferroni correction and permutation analysis ($n = 1\ 000$) for multiple testing for all of our individual SNP associations. After applying thresholds created after these corrections to our single SNP association findings, none remained significant (Bonferroni corrected $p < 0.001$). Dealing with multiple testing is a controversial issue and has been intensely debated [55-57]. Considering the exploratory nature of this study, without prespecified hypotheses for most of our SNPs, there is no clear structure in the multiple tests [56]. “Significant” findings are therefore labeled as “exploratory” with confirmatory studies needed. Suggestion of potential gene-gene interaction was observed three times in analyses using non-parametric statistical models. The recently developed MDR method improves power by data reduction to efficiently identify potential gene-gene interactions in relatively small samples [49]. After cross-validation and permutation testing procedures were performed we found a trend ($p = 0.054$) for the interaction within *GFRA1* rs11197557, *GFRA1* rs1078080 and *GFRA3* rs11242417 in the case-control sample and *GFRA1* rs10885888, *GFRA2* rs4237073 and *GFRA3* rs7726580 in the response sample. *GFRA1* rs1078080, used in our best three-locus model in case-control sample, deviated from Hardy-Weinberg equilibrium in cases ($p = 0.016$) at a significance threshold of $p < 0.05$. Trikalinos et al (2006) concluded that Hardy-Weinberg equilibrium should be routinely and transparently assessed in gene-disease association studies [58]. Discrepant results in these analyzes do not necessarily mean that postulated association should be dismissed, but they point to the need for more evidence and validation [58], especially considering that deviations can be a symptom of disease association, the implications of which are often under-exploited [59-61].

Of considerable interest, all significant findings were with markers in *GFRA1*, *GFRA2* and *GFRA3* that are located in chromosomal linkage regions for schizophrenia, while no significant results were found for *GFRA4*, which is not located in a linkage region. The interpretation of genetic linkage results is controversial and some degree of subjectivity enters into the determination of which regions of the genome should be considered to have truly significant evidence for linkage to schizophrenia. Two recent meta-analyses have summarized these findings [62, 63]. Most of the identified regions of linkage did not overlap across the two studies; however a region on chromosome 8p where *GFRA2* is located was supported by both investigations. The putative susceptibility genes for which the most follow-up genetic association data are available are those encoding dysbindin, neuregulin 1, D-amino-acid oxidase, D-amino-acid oxidase activator (formerly known as G72) and regulator of G-protein signalling 4 [64, 65]. Many of these candidate genes share a putative role in neurodevelopment, as does the GDNF pathway. The neurodevelopmental hypothesis for schizophrenia has strong support from the

effects of prenatal and perinatal insults, premorbid cognitive and neurological abnormalities, and the nature of histopathological abnormalities in brain tissue from schizophrenia patients [66].

GDNF was initially described as a trophic factor for dopaminergic neurons. Although considered one of the most potent neurotrophic factors for these neurons, GDNF is widely expressed throughout the brain, and exerts neuroprotective effects in several central and peripheral neuronal populations. Changes in the expression of other classes of neurotrophic factors and their receptors have been reported as a consequence of increased neural activity, injury and degeneration. Similarly, the expression of members of the GDNF family and their receptors are also affected by neural insults and degeneration. Studies have shown that expression of GFRA and their ligands can be altered in the rodent following peripheral nerve injury, ischemia and seizures [67]. Dopamine D2 receptor (DRD2) null mutant mice have altered GDNF levels, although *GFRA1* mRNA expression was unchanged [68]. Moreover, GDNF^{+/-} mutant mice have abnormal hippocampal synaptic transmission and impaired spatial learning [69, 70]. Rosa et al (2006) reported increased serum GDNF levels in bipolar patients during acute manic and depressive episodes when compared with matched healthy controls [71]. Medication treatments used for psychiatric disorders also alter GFRA receptors and ligand levels [37-39, 41]. Taken together, these findings suggest that alterations in GDNF signalling may play a role in neuropsychiatric disease and associated treatment effects. Our results, in combination with these other observations, suggest that the GFRA receptors may be involved in a pathway that affects neurodevelopment in schizophrenia, but further work is clearly required to strengthen this hypothesis.

Disclosure/Conflicts of interest

Souza, Romano-Silva, Culotti, MacNeil and Wong have nothing to declare. Lieberman has served as a consultant/ advisor or grantee of Acadia, Astra Zeneca, Bristol-Myers Squibb, Eli Lilly, GlaxoSmithKline, Janssen Pharmaceutica, Lundbeck, Merck, Organon, Pfizer and Wyeth; and holds a patent from Repligen. Meltzer declares that he is a consultant or grantee of Abbott, Acadia, ARYx, Astra Zeneca, Bristol Myers Squibb, Eli Lilly, Janssen, Memory, Minster, Organon, Pfizer, Solvay, Wyeth, and Vanda. Kennedy declares that he is a consultant for GlaxoSmithKline.

Acknowledgements

Authors would like to thank Daniela VF Rosa, Mary Smirniw and Nicole King for their help and support throughout manuscript preparation

Support

Funding and grants: 1) CNPq – Brazil (#202447/2006-5; #140950/2005-2; #554496/2005-4); 2) National Institutes of Health (NIH); 3) Canadian Institutes of Health Research (CIHR) #940595, 4) Ontario Mental Health Foundation (OMHF) and 5) Fapemig – Brazil. Funding agencies had no further role in study design; in the collection, analysis and interpretation of data; in the writing of the report; and in the decision to submit the paper for publication.

References

- [1] Jablensky A. Epidemiology of schizophrenia: the global burden of disease and disability. *Eur Arch Psychiatry Clin Neurosci* 2000; 250: 274-285.
- [2] Gottesman II. Schizophrenia genesis: The origins of madness. New York: Freeman; 1991.
- [3] Stefansson H, Thorgeirsson TE, Gulcher JR, Stefansson K. Neuregulin 1 in schizophrenia: out of Iceland. *Mol Psychiatry* 2003; 8: 639-640.
- [4] Straub RE, Jiang Y, MacLean CJ, Ma Y, Webb BT, Myakishev MV et al.. Genetic variation in the 6p22.3 gene DTNBP1, the human ortholog of the mouse dysbindin gene, is associated with schizophrenia. *Am J Hum Genet* 2002; 71: 337-348.
- [5] Schwab SG, Knapp M, Mondabon S, Hallmayer J, Borrmann-Hassenbach M, Albus M et al.. Support for association of schizophrenia with genetic variation in the 6p22.3 gene, dysbindin, in sib-pair families with linkage and in an additional sample of triad families. *Am J Hum Genet* 2003; 72: 185-190.
- [6] Millar JK, Wilson-Annan JC, Anderson S, Christie S, Taylor MS, Semple CA et al. Disruption of two novel genes by a translocation co-segregating with schizophrenia. *Hum Mol Genet* 2000; 9: 1415-1423.
- [7] Owen MJ, Craddock N, O'Donovan MC. Schizophrenia: genes at last? *Trends Genet* 2005; 21: 518-525.
- [8] Ross CA, Margolis RL, Reading SA, Pletnikov M, Coyle JT. Neurobiology of schizophrenia. *Neuron* 2006; 52: 139-153.
- [9] Duman RS. Synaptic plasticity and mood disorders. *Mol Psychiatry* 2002; 7: S29-S34.
- [10] Obara Y, Nakahata N. The Signaling Pathway of Neurotrophic Factor Biosynthesis. *Drug News Perspect* 2002; 15: 290-298.
- [11] Chlan-Fourney J, Ashe P, Nysten K, Juorio AV, Li XM. Differential regulation of hippocampal BDNF mRNA by typical and atypical antipsychotic administration. *Brain Res* 2002; 954: 11-20.
- [12] Xu H, Qing H, Lu W, Keegan D, Richardson JS, Chlan-Fourney J et al. Quetiapine attenuates the immobilization stress-induced decrease of brain-derived neurotrophic factor expression in rat hippocampus. *Neurosci Lett* 2002; 321: 65-68.
- [13] Bai O, Chlan-Fourney J, Bowen R, Keegan D, Li XM. Expression of brain-derived neurotrophic factor mRNA in rat hippocampus after treatment with antipsychotic drugs. *J Neurosci Res* 2003; 71: 127-131.

- [14] Angelucci F, Aloe L, Gruber SH, Fiore M, Mathe AA. Chronic antipsychotic treatment selectively alters nerve growth factor and neuropeptide Y immunoreactivity and the distribution of choline acetyl transferase in rat brain regions. *Int J Neuropsychopharmacol* 2000; 3: 13-25.
- [15] Takahashi M, Shirakawa O, Toyooka K, Kitamura N, Hashimoto T, Maeda K et al.. Abnormal expression of brain-derived neurotrophic factor and its receptor in the corticolimbic system of schizophrenic patients. *Mol Psychiatry* 2000; 5: 293-300.
- [16] Lin LF, Doherty DH, Lile JD, Bektesh S, Collins F. GDNF: a glial cell line-derived neurotrophic factor for midbrain dopaminergic neurons. *Science* 1993; 260: 1130-1132.
- [17] Tomac A, Widenfalk J, Lin LF, Kohno T, Ebendal T, Hoffer BJ et al.. Retrograde axonal transport of glial cell line-derived neurotrophic factor in the adult nigrostriatal system suggests a trophic role in the adult. *Proc Natl Acad Sci U S A* 1995; 92: 8274-8278.
- [18] Trupp M, Belluardo N, Funakoshi H, Ibanez CF. Complementary and overlapping expression of glial cell line-derived neurotrophic factor (GDNF), c-ret proto-oncogene, and GDNF receptor-alpha indicates multiple mechanisms of trophic actions in the adult rat CNS. *J Neurosci* 1997; 17: 3554-3567.
- [19] Henderson CE, Phillips HS, Pollock RA, Davies AM, Lemeulle C, Armanini M et al. GDNF: a potent survival factor for motoneurons present in peripheral nerve and muscle. *Science* 1994; 266: 1062-1064.
- [20] Beck KD, Valverde J, Alexi T, Poulsen K, Moffat B, Vandlen RA et al. Mesencephalic dopaminergic neurons protected by GDNF from axotomy-induced degeneration in the adult brain. *Nature* 1995; 373: 339-341.
- [21] Bowenkamp KE, David D, Lapchak PL, Henry MA, Granholm AC, Hoffer BJ et al. 6-hydroxydopamine induces the loss of the dopaminergic phenotype in substantia nigra neurons of the rat. A possible mechanism for restoration of the nigrostriatal circuit mediated by glial cell line-derived neurotrophic factor. *Exp Brain Res* 1996; 111: 1-7.
- [22] Choi-Lundberg DL, Lin Q, Chang YN, Chiang YL, Hay CM, Mohajeri H et al. Dopaminergic neurons protected from degeneration by GDNF gene therapy. *Science* 1997; 275: 838-841.
- [23] Airaksinen MS, Titievsky A, Saarma M. GDNF family neurotrophic factor signaling: four masters, one servant? *Mol Cell Neurosci* 1999; 13: 313-325.
- [24] Jing S, Wen D, Yu Y, Holst PL, Luo Y, Fang M et al. GDNF-induced activation of the ret protein tyrosine kinase is mediated by GDNFR-alpha, a novel receptor for GDNF. *Cell* 1996; 85: 1113-1124.
- [25] Treanor JJ, Goodman L, de SF, Stone DM, Poulsen KT, Beck CD et al.. Characterization of a multicomponent receptor for GDNF. *Nature* 1996; 382: 80-83.

- [26] Klein RD, Sherman D, Ho WH, Stone D, Bennett GL, Moffat B et al. A GPI-linked protein that interacts with Ret to form a candidate neurturin receptor. *Nature* 1997; 387: 717-721.
- [27] Sanicola M, Hession C, Worley D, Carmillo P, Ehrenfels C, Walus L et al. Glial cell line-derived neurotrophic factor-dependent RET activation can be mediated by two different cell-surface accessory proteins. *Proc Natl Acad Sci U S A* 1997; 94: 6238-6243.
- [28] Baloh RH, Tansey MG, Lampe PA, Fahrner TJ, Enomoto H, Simburger KS et al. Artemin, a novel member of the GDNF ligand family, supports peripheral and central neurons and signals through the GFRalpha3-RET receptor complex. *Neuron* 1998; 21: 1291-1302.
- [29] Enokido Y, de SF, Hongo JA, Ninkina N, Rosenthal A, Buchman VL et al. GFR alpha-4 and the tyrosine kinase Ret form a functional receptor complex for persephin. *Curr Biol* 1998; 8: 1019-1022.
- [30] Masure S, Cik M, Hoefnagel E, Nosrat CA, Van dL, I, Scott R et al. Mammalian GFRalpha-4, a divergent member of the GFRalpha family of coreceptors for glial cell line-derived neurotrophic factor family ligands, is a receptor for the neurotrophic factor persephin. *J Biol Chem* 2000; 275: 39427-39434.
- [31] Colavita A, Krishna S, Zheng H, Padgett RW, Culotti JG. Pioneer axon guidance by UNC-129, a *C. elegans* TGF-beta. *Science* 1998; 281: 706-709.
- [32] Nash B, Colavita A, Zheng H, Roy PJ, Culotti JG. The forkhead transcription factor UNC-130 is required for the graded spatial expression of the UNC-129 TGF-beta guidance factor in *C. elegans*. *Genes Dev* 2000; 14: 2486-2500.
- [33] Merz DC, Alves G, Kawano T, Zheng H, Culotti JG. UNC-52/perlecan affects gonadal leader cell migrations in *C. elegans* hermaphrodites through alterations in growth factor signaling. *Dev Biol* 2003; 256: 173-186.
- [34] Burke RE. GDNF as a candidate striatal target-derived neurotrophic factor for the development of substantia nigra dopamine neurons. *J Neural Transm* 2006 Suppl 41-45.
- [35] Basile VS, Masellis M, Potkin SG, Kennedy JL. Pharmacogenomics in schizophrenia: the quest for individualized therapy. *Hum Mol Genet* 2002; 11: 2517-2530.
- [36] Malhotra AK, Murphy GM, Jr., Kennedy JL. Pharmacogenetics of psychotropic drug response. *Am J Psychiatry* 2004; 161: 780-796.
- [37] Hisaoka K, Nishida A, Koda T, Miyata M, Zensho H, Morinobu S et al. Antidepressant drug treatments induce glial cell line-derived neurotrophic factor (GDNF) synthesis and release in rat C6 glioblastoma cells. *J Neurochem* 2001; 79: 25-34.
- [38] Shao Z, Dyck LE, Wang H, Li XM. Antipsychotic drugs cause glial cell line-derived neurotrophic factor secretion from C6 glioma cells. *J Psychiatry Neurosci* 2006; 31: 32-37.

- [39] Angelucci F, Aloe L, Jimenez-Vasquez P, Mathe AA. Lithium treatment alters brain concentrations of nerve growth factor, brain-derived neurotrophic factor and glial cell line-derived neurotrophic factor in a rat model of depression. *Int J Neuropsychopharmacol* 2003; 6: 225-231.
- [40] Castro LM, Gallant M, Niles LP. Novel targets for valproic acid: up-regulation of melatonin receptors and neurotrophic factors in C6 glioma cells. *J Neurochem* 2005; 95: 1227-1236.
- [41] Chen AC, Eisch AJ, Sakai N, Takahashi M, Nestler EJ, Duman RS. Regulation of GFRalpha-1 and GFRalpha-2 mRNAs in rat brain by electroconvulsive seizure. *Synapse* 2001; 39: 42-50.
- [42] Kane J, Honigfeld G, Singer J, Meltzer H. Clozapine for the treatment-resistant schizophrenic. A double-blind comparison with chlorpromazine. *Arch Gen Psychiatry* 1988 45: 789-796.
- [43] Hwang R, Shinkai T, De Luca, V, Muller DJ, Ni X, Macciardi F et al. Association study of 12 polymorphisms spanning the dopamine D(2) receptor gene and clozapine treatment response in two treatment refractory/intolerant populations. *Psychopharmacology (Berl)* 2005; 181: 179-187.
- [44] Souza RP, Romano-Silva MA, Lieberman JA, Meltzer HY, Wong AH, Kennedy JL. Association study of GSK3 gene polymorphisms with schizophrenia and clozapine response. *Psychopharmacology (Berl)* 2008;
- [45] Barrett JC, Fry B, Maller J, Daly MJ. Haploview: analysis and visualization of LD and haplotype maps. *Bioinformatics* 2005; 21: 263-265.
- [46] Laird NM, Horvath S, Xu X. Implementing a unified approach to family-based tests of association. *Genet Epidemiol* 2000; 19: S36-S42.
- [47] Wigginton JE, Abecasis GR. PEDSTATS: descriptive statistics, graphics and quality assessment for gene mapping data. *Bioinformatics* 2005; 21: 3445-3447.
- [48] Dudbridge F. Pedigree disequilibrium tests for multilocus haplotypes. *Genet Epidemiol* 2003; 25: 115-121.
- [49] Ritchie MD, Hahn LW, Roodi N, Bailey LR, Dupont WD, Parl FF et al. Multifactor-dimensionality reduction reveals high-order interactions among estrogen-metabolism genes in sporadic breast cancer. *Am J Hum Genet* 2001; 69: 138-147.
- [50] Ritchie MD, Hahn LW, Moore JH. Power of multifactor dimensionality reduction for detecting gene-gene interactions in the presence of genotyping error, missing data, phenocopy, and genetic heterogeneity. *Genet Epidemiol* 2003;24: 150-157.
- [51] Moore JH, Gilbert JC, Tsai CT, Chiang FT, Holden T, Barney N et al.. A flexible computational framework for detecting, characterizing, and interpreting statistical patterns of epistasis in genetic studies of human disease susceptibility. *J Theor Biol* 2006; 241: 252-261.

- [52] Ritchie MD, Motsinger AA (2005). Multifactor dimensionality reduction for detecting gene-gene and gene-environment interactions in pharmacogenomics studies. *Pharmacogenomics* 2005; 6: 823-834.
- [53] Gabriel SB, Schaffner SF, Nguyen H, Moore JM, Roy J, Blumenstiel B et al. The structure of haplotype blocks in the human genome. *Science* 2002; 296: 2225-2229.
- [54] Durrant C, Zondervan KT, Cardon LR, Hunt S, Deloukas P, Morris AP. Linkage disequilibrium mapping via cladistic analysis of single-nucleotide polymorphism haplotypes. *Am J Hum Genet* 2004; 75: 35-43.
- [55] Aickin M. Other method for adjustment of multiple testing exists. *BMJ* 1999; 318: 127-128.
- [56] Bender R, Lange S. Multiple test procedures other than Bonferroni's deserve wider use. *BMJ* 1999; 318: 600-601.
- [57] Perneger TV. What's wrong with Bonferroni adjustments. *BMJ* 1998; 316: 1236-1238.
- [58] Trikalinos TA, Salanti G, Khoury MJ, Ioannidis JP. Impact of violations and deviations in Hardy-Weinberg equilibrium on postulated gene-disease associations. *Am J Epidemiol* 2006; 163: 300-309.
- [59] Nielsen DM, Ehm MG, Weir BS. Detecting marker-disease association by testing for Hardy-Weinberg disequilibrium at a marker locus. *Am J Hum Genet* 1998; 63: 1531-1540.
- [60] Wittke-Thompson JK, Pluzhnikov A, Cox NJ. Rational inferences about departures from Hardy-Weinberg equilibrium. *Am J Hum Genet* 2005; 76: 967-986.
- [61] Balding DJ. A tutorial on statistical methods for population association studies. *Nat Rev Genet* 2006; 7: 781-791.
- [62] Badner JA, Gershon ES. Meta-analysis of whole-genome linkage scans of bipolar disorder and schizophrenia. *Mol Psychiatry* 2002; 7: 405-411.
- [63] Lewis CM, Levinson DF, Wise LH, DeLisi LE, Straub RE, Hovatta I et al. Genome scan meta-analysis of schizophrenia and bipolar disorder, part II: Schizophrenia. *Am J Hum Genet* 2003; 73: 34-48.
- [64] Harrison PJ, Weinberger DR. Schizophrenia genes, gene expression, and neuropathology: on the matter of their convergence. *Mol Psychiatry* 2005; 10: 40-68.
- [65] Arranz MJ, de Leon J. Pharmacogenetics and pharmacogenomics of schizophrenia: a review of last decade of research. *Mol Psychiatry* 2007; 12: 707-747.
- [66] Lewis DA, Levitt P. Schizophrenia as a disorder of neurodevelopment. *Annu Rev Neurosci* 2002; 25: 409-432.

- [67] Kozłowski DA, Miljan EA, Bremer EG, Harrod CG, Gerin C, Connor B et al. Quantitative analyses of GFRalpha-1 and GFRalpha-2 mRNAs and tyrosine hydroxylase protein in the nigrostriatal system reveal bilateral compensatory changes following unilateral 6-OHDA lesions in the rat. *Brain Res* 2004 1016: 170-181.
- [68] Bozzi Y, Borrelli E. Absence of the dopamine D2 receptor leads to a decreased expression of GDNF and NT-4 mRNAs in restricted brain areas. *Eur J Neurosci* 1999; 11: 1275-1284.
- [69] Nanobashvili A, Airaksinen MS, Kokaia M, Rossi J, Asztely F, Olofsdotter K et al. Development and persistence of kindling epilepsy are impaired in mice lacking glial cell line-derived neurotrophic factor family receptor alpha 2. *Proc Natl Acad Sci U S A* 2000; 97: 12312-12317.
- [70] Gerlai R, McNamara A, Choi-Lundberg DL, Armanini M, Ross J, Powell-Braxton L et al. Impaired water maze learning performance without altered dopaminergic function in mice heterozygous for the GDNF mutation. *Eur J Neurosci* 2001; 14 : 1153-1163.
- [71] Rosa AR, Frey BN, Andreazza AC, Ceresér KM, Cunha AB et al. Increased serum glial cell line-derived neurotrophic factor immunocontent during manic and depressive episodes in individuals with bipolar disorder. *Neurosci Lett* 2006; 407: 146-150.

Table 1:

rs	Control				Case				rs	Control				Case				rs	Control				Case				
	n	%	n	%	n	%	n	%		n	%	n	%	n	%	n	%		n	%	n	%					
GFR1 rs1078080	A	271	61.9	A	271	64.2	GFR1 rs11598215	A	105	24.0	A	86	20.4	GFR1 rs3781514	A	70	16.0	A	69	16.4	GFR1 rs2694783	A	250	57.1	A	247	58.5
	G	167	38.1	G	151	35.8		G	333	76.0	G	336	79.6		G	368	84.0	G	353	83.6		T	188	42.9	T	175	41.5
	AA	87	39.7	AA	95	45.0		AA	14	06.4	AA	09	04.3		AA	04	01.8	AA	07	03.3		AA	68	31.1	AA	70	33.2
	AG	97	44.3	AG	81	38.4		AG	77	35.2	AG	68	32.2		AG	62	28.3	AG	55	26.1		AT	114	52.1	AT	107	50.7
	GG	35	16.0	GG	35	16.6		GG	128	58.4	GG	134	63.5		GG	153	69.9	GG	149	70.6		TT	37	16.9	TT	34	16.1
GFR1 rs2694801	A	28	19.2	A	19	14.2	GFR1 rs3824840	A	108	24.9	A	100	23.7	GFR1 rs12776813	A	67	15.3	A	67	15.9	GFR1 rs7085306	A	263	60.0	A	255	60.4
	G	118	80.8	G	115	85.8		G	326	75.1	G	322	76.3		G	371	84.7	G	355	84.1		G	175	40.0	G	167	39.6
	AA	04	05.5	AA	01	01.5		AA	19	08.8	AA	09	04.3		AA	05	02.3	AA	04	01.9		AA	80	36.5	AA	71	33.6
	AG	20	27.4	AG	17	25.4		AG	70	32.3	AG	82	38.9		AG	57	26.0	AG	59	28.0		AG	103	47.0	AG	113	53.6
	GG	49	67.1	GG	49	73.1		GG	128	59.0	GG	120	56.9		GG	157	71.7	GG	148	70.1		GG	36	16.4	GG	27	12.8
GFR1 rs10787627	A	202	46.1	A	195	46.2	GFR1 rs9787429	A	95	21.9	A	87	20.6	GFR1 rs12775655	A	324	74.0	A	330	78.2	GFR1 rs11197557	A	99	22.6	A	69	16.4
	G	236	53.9	G	227	53.8		G	339	78.1	G	335	79.4		C	114	26.0	C	92	21.8		G	339	77.4	G	353	83.6
	AA	43	19.6	AA	45	21.3		AA	05	02.3	AA	06	02.8		AA	119	54.3	AA	128	60.7		AA	04	01.8	AA	00	0.00
	AG	116	53.0	AG	105	49.8		AG	85	39.2	AG	75	35.5		AC	86	39.3	AC	74	35.1		AG	91	41.6	AG	69	32.7
	GG	60	27.0	GG	61	27.4		GG	127	58.5	GG	130	61.6		CC	14	06.4	CC	09	04.3		GG	124	56.6	GG	142	67.3
GFR1 rs10749189	A	269	61.4	A	272	64.5	GFR1 rs11197567	A	102	23.3	A	115	27.3	GFR1 rs3781539	A	375	85.6	A	358	84.8	GFR1 rs7903297	A	205	46.3	A	188	44.5
	G	169	38.6	G	150	35.5		G	336	76.7	G	307	72.7		G	63	14.4	G	64	15.2		C	235	53.7	C	234	55.5
	AA	75	34.2	AA	85	40.3		AA	15	6.8	AA	15	7.1		AA	161	73.5	AA	152	72.0		AA	50	22.8	AA	41	19.4
	AG	119	54.3	AG	102	48.3		AG	72	32.9	AG	85	40.3		AG	53	24.2	AG	54	25.6		AC	103	47.0	AC	106	50.2
	GG	25	11.4	GG	24	11.4		GG	132	60.3	GG	111	52.6		GG	05	02.3	GG	05	02.4		CC	66	30.1	CC	64	30.3
GFR1 rs17094340	A	358	81.7	A	363	86.0	GFR1 rs7920934	A	328	74.9	A	318	75.7	GFR1 rs10885877	A	106	24.2	A	115	27.3	GFR1 rs4751956	A	203	46.3	A	185	43.8
	G	80	18.3	G	59	14.0		G	110	25.1	G	102	24.3		G	332	75.8	G	307	72.7		G	235	53.7	G	237	56.2
	AA	147	67.1	AA	153	72.5		AA	123	56.2	AA	122	58.1		AA	14	06.4	AA	13	06.2		AA	49	22.4	AA	40	19.0
	AG	64	29.2	AG	57	27.0		AG	82	37.4	AG	74	35.2		AG	78	35.6	AG	89	42.2		AG	105	47.9	AG	105	49.8
	GG	08	03.7	GG	01	00.5		GG	14	06.4	GG	14	06.7		GG	127	58.0	GG	109	51.7		GG	65	29.7	GG	66	31.3
GFR1 rs11812459	A	175	40.1	A	191	45.3	GFR1 rs10885888	A	97	22.4	A	92	21.9	GFR1 rs12413585	A	226	51.6	A	210	49.8	GFR1 rs730357	A	312	71.2	A	312	73.9
	G	261	59.9	G	231	54.7		G	337	77.6	G	328	78.1		G	212	48.4	G	212	50.2		G	126	28.8	G	110	26.1
	AA	35	16.1	AA	46	21.8		AA	10	04.6	AA	03	01.4		AA	57	26.0	AA	47	22.3		AA	112	51.1	AA	115	54.5
	AG	105	48.2	AG	99	46.9		AG	77	35.2	AG	86	40.8		AG	112	51.1	AG	116	55.0		AG	88	40.2	AG	82	38.9
	GG	78	35.8	GG	66	31.3		GG	130	59.4	GG	121	59.4		GG	50	22.8	GG	48	22.7		GG	19	8.7	GG	14	6.6

rs	Control				Case				rs	Control				Case				rs	Control				Case																																																																																																																
	n	%	n	%	n	%	n	%		n	%	n	%	n	%	n	%		n	%	n	%																																																																																																																	
GFRA1 rs11197612	A	45	30.8	A	38	28.4	GFRA2 rs15881	A	62	42.5	A	72	53.7	GFRA2 rs10283397	C	66	45.2	C	62	46.3	GFRA2 rs11993990	A	71	48.6	A	67	50.0	G	101	69.2	G	96	71.6	C	84	57.5	C	62	46.3	G	75	51.4	G	67	50.0	AA	09	12.3	AA	07	10.4	AA	12	16.4	AA	21	31.3	CC	14	19.2	CC	17	25.4	AA	15	20.5	AA	20	29.9	AG	27	37.0	AG	24	35.8	AC	38	52.1	AC	30	44.8	CG	38	52.1	CG	28	41.8	AG	41	56.2	AG	27	40.3	GG	37	50.7	GG	36	53.7	CC	23	31.5	CC	16	23.9	GG	21	28.8	GG	22	32.8	GG	17	23.3	GG	20	29.9																		
	GFRA2 rs7813735	A	76	52.1	A	73		54.5	GFRA2 rs10088105	A	22	15.1	A		13	09.7	GFRA2 rs4567027	C	104	71.2		C	102	76.1	GFRA2 rs4237073	A	62	42.5	A	71	53.0	G	70	47.9	G	61	45.5	G	42	28.8	G	32	23.9	G	42	57.5	G	63	47.0	AA	18	24.7	AA	20	29.9	AA	02	02.7	AA	02	03.0	CC	36	49.3	CC	39	58.2	AA	14	19.2	AA	20	29.9	AG	40	54.8	AG	33	49.3	AG	18	24.7	AG	09	13.4	CG	32	43.8	CG	24	35.8	AG	34	46.6	AG	31	46.3	GG	15	20.5	GG	14	20.9	GG	53	72.6	GG	56	83.6	GG	05	06.8	GG	04	06.0	GG	25	34.2	GG	16	23.9														
		GFRA2 rs4078157	A	61	41.8	A		47		35.1	GFRA2 rs13250096	C	18		12.3	C		06	4.5	GFRA2 rs4739285		A	85	58.2		A	64	47.8	GFRA2 rs4739286	A	43	29.5	A	33	24.6	G	85	58.2	G	87	64.9	G	61	41.8	G	70	52.2	G	103	70.5	G	101	75.4	AA	13	17.8	AA	8	11.9	CC	01	01.4	CC	00	0.0	AA	04	05.5	AA	06	09.0	AG	35	47.9	AG	31	46.3	CG	16	21.9	CG	06	9.0	AG	35	47.9	AG	21	33.3	GG	25	34.2	GG	28	41.8	GG	56	76.7	GG	61	91.0	GG	34	46.6	GG	40	59.7																												
			GFRA2 rs4567028	A	104	71.2		A		109		81.3	GFRA2 rs6587002		A	99		67.8	A			91	67.9	GFRA2 rs4739217		C	79	55.6		C	62	46.3	GFRA2 rs7014143	A	53	36.3	A	50	37.3	G	42	28.8	G	25	18.7	G	63	44.4	G	72	53.7	C	93	63.7	C	84	62.7	AA	35	47.9	AA	45	67.2	AA	36	49.3	AA	33	49.3	CC	24	32.9	CC	19	28.4	AA	9	12.3	AA	10	14.9	AG	34	46.6	AG	19	28.4	AG	27	37.0	AG	25	37.3	CG	31	42.5	CG	24	35.8	AC	35	47.9	AC	30	44.8	GG	04	05.5	GG	03	04.5	GG	10	13.7	GG	09	13.4	CC	29	39.7	CC	27	40.3												
				GFRA2 rs1128397	A	69		47.3		A		43			32.1	GFRA2 rs6988470		A	77			52.7	A			70	52.2	GFRA3 rs10036665		A	117	80.1		A	105	78.4	GFRA3 rs10952	A	43	29.5	A	36	26.9	T	77	52.7	T	91	67.9	G	69	47.3	G	64	47.8	T	29	19.9	T	29	21.6	T	103	70.5	T	98	73.1	AA	14	19.2	AA	7	10.4	AA	20	27.4	AA	19	28.4	AA	47	64.4	AA	41	61.2	AA	06	08.2	AA	05	07.5	AT	41	56.2	AT	29	43.3	AG	37	50.7	AG	32	47.8	AT	23	31.5	AT	23	34.3	AT	31	42.5	AT	26	38.8	TT	18	24.7	TT	31	46.3	GG	16	21.9	GG	16	23.9	TT	03	04.1	TT	03	04.5	TT	36
GFRA3 rs11242417					A	124	84.9	A		113		84.3		GFRA3 rs7726580	A			50	34.2		A	45	33.6			GFRA4 rs6084432	A			23	15.8	A		23	17.2	GFRA4 rs633924		A	89	61.0	A	79	59.0	C	22	15.1	C	21	15.7	G	96	65.8	G	89	66.4	G	123	84.2	G	111	82.8	G	87	39.0	G	55	41.0	AA	52	71.2	AA	48	71.6	AA	8	11.0	AA	10	14.9	AA	01	01.4	AA	02	03.0	AA	27	37.0	AA	24	35.8	AC	20	27.4	AC	17	25.4	AG	34	46.6	AG	25	37.3	AG	21	28.8	AG	19	28.4	AG	35	47.9	AG	31	46.3	CC	01	01.4	CC	02	03.0	GG	31	42.5	GG	32	47.8	GG	51	69.9	GG	46	68.7	GG	11

Table 2

Gene	rs	Allele	Frequency	Family	S	E(S)	Var(S)	Z	P
GFRA1	rs1078080	A	0.696	32	48	42.500	12.028	1.586	0.113
	rs11598215	A	0.190	25	13	17.000	8.778	-1.350	0.177
	rs3781514	A	0.198	30	18	17.733	8.462	0.092	0.927
	rs2694783	A	0.564	41	41	44.900	14.712	-1.017	0.309
	rs2694801	A	0.233	27	22	20.833	8.194	0.408	0.684
	rs3824840	A	0.281	29	25	23.433	7.934	0.556	0.578
	rs12776813	A	0.183	30	15	17.333	8.222	-0.814	0.416
	rs7085306	A	0.573	40	46	45.067	13.240	0.257	0.798
	rs10787627	A	0.461	40	40	39.600	13.462	0.109	0.913
	rs9787429	A	0.248	24	21	21.833	7.194	-0.311	0.756
	rs12775655	A	0.772	32	45	44.100	9.212	0.297	0.767
	rs11197557	A	0.216	33	19	19.833	9.417	-0.272	0.786
	rs10749189	A	0.617	40	48	45.900	13.712	0.567	0.571
	rs11197567	A	0.273	26	20	20.633	8.354	-0.219	0.827
	rs3781539	A	0.818	21	32	29.867	6.104	0.863	0.388
	rs7903297	A	0.412	32	34	31.833	11.417	0.641	0.521
	rs17094340	A	0.845	20	28	29.833	5.194	-0.804	0.421
	rs7920934	A	0.764	34	53	46.667	10.722	1.934	0.053
	rs10885877	A	0.311	25	25	21.333	7.667	1.324	0.185
	rs4751956	A	0.472	35	39	35.333	12.167	1.051	0.293
rs11812459	A	0.450	29	30	30.667	10.953	-0.201	0.840	
rs10885888	A	0.241	26	14	16.583	7.632	-0.935	0.350	
rs12413585	A	0.499	35	34	33.333	12.444	0.189	0.850	
rs730357	A	0.742	30	45	38.167	10.472	2.112	0.035	
rs11197612	A	0.388	30	25	25.333	10.167	-0.105	0.917	
GFRA2	rs15881	A	0.500	29	32	31.000	10.614	0.307	0.759
	rs4567027	C	0.713	33	47	44.667	10.558	0.718	0.473
	rs7813735	A	0.485	33	29	31.333	12.558	-0.658	0.510
	rs10088105	A	0.151	18	09	10.333	5.222	-0.583	0.560
	rs1128397	A	0.432	37	29	33.167	13.086	-1.152	0.249
	rs6988470	A	0.548	30	32	31.167	10.972	0.252	0.801
	rs4237073	A	0.519	37	43	39.700	12.632	0.928	0.353
	rs4739217	C	0.483	33	35	35.333	12.500	-0.094	0.925
	rs10283397	C	0.483	30	31	30.200	10.827	0.243	0.808
	rs6587002	A	0.662	27	34	33.833	8.972	0.056	0.956
	rs7014143	A	0.418	29	27	24.367	9.632	0.848	0.396
	rs4567028	A	0.777	29	45	42.500	9.528	0.810	0.418
	rs4739286	A	0.315	30	24	24.000	10.722	0.000	1.000
	rs11993990	A	0.554	37	49	46.667	12.953	0.648	0.517
	rs4078157	A	0.350	33	26	27.833	10.808	-0.558	0.577
rs4739285	A	0.503	35	37	38.167	12.250	-0.333	0.739	
rs13250096	C	0.157	18	11	12.333	5.722	-0.557	0.577	
GFRA3	rs10036665	A	0.830	16	28	25.167	4.972	1.271	0.204
	rs10952	A	0.365	34	25	24.167	10.250	0.260	0.795
	rs11242417	A	0.856	14	24	20.667	4.222	1.622	0.105
	rs7726580	A	0.413	36	26	27.667	11.336	-0.495	0.621
GFRA4	rs633924	A	0.647	33	45	42.167	11.472	0.837	0.403
	rs6084432	A	0.200	22	10	13.667	6.944	-1.391	0.164

Table 3

rs	Non-responder				Responder				rs	Non-responder				Responder				rs	Non-responder				Responder			
	n	%	n	%	n	%	n	%		n	%	n	%	n	%	n	%		n	%	n	%				
GFR1 rs1078080	A	87	59.6	A	77	57.5	GFR1 rs11598215	34	23.3	A	37	27.6	GFR1 rs3781514	A	14	09.6	A	17	12.7	GFR1 rs2694783	A	86	58.9	A	82	61.2
	G	59	40.4	G	57	42.5		112	76.7	G	97	72.4		G	132	90.4	G	117	87.3		T	60	41.1	T	52	38.8
	AA	28	38.4	AA	25	37.3		04	05.5	AA	08	11.9		AA	01	01.4	AA	01	01.5		AA	28	38.4	AA	26	38.8
	AG	31	42.5	AG	27	40.3		26	35.6	AG	21	31.3		AG	12	16.4	AG	15	22.4		AT	30	41.1	AT	30	44.8
	GG	14	19.2	GG	15	22.4		43	58.9	GG	38	56.7		GG	60	82.2	GG	51	82.2		TT	15	20.5	TT	11	16.4
GFR1 rs2694801	A	28	19.2	A	19	14.2	GFR1 rs3824840	31	21.2	A	28	20.9	GFR1 rs12776813	A	21	14.4	A	15	11.2	GFR1 rs7085306	A	93	63.7	A	89	66.4
	G	118	80.8	G	115	85.8		115	78.8	G	106	79.1		G	125	85.6	G	119	88.8		G	53	36.3	G	45	33.6
	AA	04	05.5	AA	01	01.5		03	4.1	AA	00	0.00		AA	01	01.4	AA	01	01.5		AA	30	41.1	AA	27	40.3
	AG	20	27.4	AG	17	25.4		25	34.2	AG	28	41.8		AG	19	26.0	AG	13	19.4		AG	33	45.2	AG	35	52.2
	GG	49	67.1	GG	49	73.1		45	61.6	GG	39	58.2		GG	53	72.6	GG	53	79.1		GG	10	13.7	GG	05	07.5
GFR1 rs10787627	A	68	46.6	A	62	46.3	GFR1 rs9787429	28	19.2	A	25	18.7	GFR1 rs12775655	A	119	81.5	A	103	76.9	GFR1 rs11197557	A	22	15.1	A	20	14.9
	G	78	53.4	G	72	53.7		118	80.8	G	109	81.3		C	27	18.5	G	31	23.1		G	124	84.9	G	114	85.1
	AA	20	27.4	AA	16	27.4		01	01.4	AA	00	0.00		AA	48	65.8	AA	40	59.7		AA	01	1.4	AA	00	0.00
	AG	28	38.4	AG	30	38.4		26	35.6	AG	25	37.3		AC	23	31.5	AG	23	34.3		AG	20	27.4	AG	20	29.9
	GG	25	34.2	GG	21	34.2		46	63.0	GG	42	62.7		CC	02	02.7	GG	04	06.0		GG	52	21.2	GG	47	70.1
GFR1 rs10749189	A	102	69.9	A	82	61.2	GFR1 rs11197567	32	21.9	A	35	26.1	GFR1 rs3781539	A	131	89.7	A	120	89.6	GFR1 rs7903297	A	59	40.4	A	57	42.5
	G	44	30.1	G	52	38.8		114	78.1	G	99	73.9		G	15	10.3	G	14	10.4		C	87	59.6	C	77	57.5
	AA	33	45.2	AA	24	35.8		02	02.7	AA	03	04.5		AA	59	80.8	AA	53	79.1		AA	08	11.0	AA	10	14.9
	AG	36	49.3	AG	34	50.7		28	38.4	AG	29	43.3		AG	13	17.8	AG	14	20.9		AC	43	58.9	AC	37	55.2
	GG	04	05.5	GG	09	13.4		43	58.9	GG	35	52.2		GG	01	1.4	GG	00	00.0		CC	22	30.1	CC	20	29.9
GFR1 rs17094340	A	130	89.0	A	118	88.1	GFR1 rs7920934	100	68.5	A	100	74.6	GFR1 rs10885877	A	31	21.2	A	41	30.6	GFR1 rs4751956	A	61	41.8	A	47	35.1
	G	16	11.0	G	16	11.9		46	31.5	G	34	25.4		G	115	78.8	G	93	69.4		G	85	58.2	G	87	64.9
	AA	57	78.1	AA	51	76.1		35	47.9	AA	39	58.2		AA	04	05.5	AA	05	07.5		AA	14	19.2	AA	11	16.4
	AG	16	21.9	AG	16	23.9		30	41.1	AG	22	32.8		AG	23	31.5	AG	31	46.3		AG	33	45.2	AG	25	37.3
	GG	00	00.0	GG	00	00.0		08	11.0	GG	06	09.0		GG	46	63.0	GG	31	46.3		GG	26	35.6	GG	31	46.3
GFR1 rs11812459	A	64	43.8	A	65	48.5	GFR1 rs10885888	45	30.8	A	33	24.6	GFR1 rs12413585	A	77	52.7	A	60	44.8	GFR1 rs730357	A	105	77.6	A	104	77.6
	G	82	56.2	G	69	51.5		101	69.2	G	101	75.4		G	69	47.3	G	74	55.2		G	41	22.4	G	30	22.4
	AA	13	17.8	AA	17	25.4		06	08.2	AA	04	06.0		AA	20	27.4	AA	13	19.4		AA	38	52.1	AA	39	58.2
	AG	38	52.1	AG	31	46.3		33	45.2	AG	25	37.3		AG	37	50.7	AG	34	50.7		AG	29	39.7	AG	26	38.8
	GG	22	30.1	GG	19	28.4		34	46.6	GG	38	56.7		GG	16	21.9	GG	20	29.9		GG	06	08.2	GG	02	03.0

rs	Non-responder				Responder				rs	Non-responder				Responder				rs	Non-responder				Responder																																																																																																																																																		
	n	%	n	%	n	%	n	%		n	%	n	%	n	%	n	%		n	%	n	%																																																																																																																																																			
GFRA1 rs11197612	A	45	30.8	A	38	28.4	GFRA2 rs15881	A	62	42.5	A	72	53.7	GFRA2 rs10283397	C	66	45.2	C	62	46.3	GFRA2 rs11993990	A	71	48.6	A	67	50.0	GFRA2 rs11993990	G	101	69.2	G	96	71.6	GFRA2 rs15881	C	84	57.5	C	62	46.3	GFRA2 rs10283397	G	80	54.8	G	72	53.7	GFRA2 rs11993990	G	75	51.4	G	67	50.0	GFRA2 rs15881	AA	09	12.3	AA	07	10.4	GFRA2 rs10283397	CC	14	19.2	CC	17	25.4	GFRA2 rs11993990	AA	15	20.5	AA	20	29.9	GFRA2 rs15881	AG	27	37.0	AG	24	35.8	GFRA2 rs10283397	CG	38	52.1	CG	28	41.8	GFRA2 rs11993990	AG	41	56.2	AG	27	40.3	GFRA2 rs15881	GG	37	50.7	GG	36	53.7	GFRA2 rs10283397	GG	21	28.8	GG	22	32.8	GFRA2 rs11993990	GG	17	23.3	GG	20	29.9																																																			
	GFRA2 rs7813735	A	76	52.1	A	73		54.5	GFRA2 rs10088105	A	22	15.1	A		13	09.7	GFRA2 rs4567027	C	104	71.2		C	102	76.1	GFRA2 rs4237073	A	62		42.5	A	71	53.0	GFRA2 rs4237073	G		70	47.9	G	61	45.5	GFRA2 rs10088105		G	124	84.9	G	121	90.3		GFRA2 rs4567027	G	42	28.8	G	32		23.9	GFRA2 rs4237073	G	42	57.5	G		63	47.0	GFRA2 rs10088105	AA	18	24.7		AA	20	29.9	GFRA2 rs4567027	CC	36		49.3	CC	39	58.2	GFRA2 rs4237073	AA		14	19.2	AA	20	29.9	GFRA2 rs10088105		AG	40	54.8	AG	33	49.3		GFRA2 rs4567027	CG	32	43.8	CG	24		35.8	GFRA2 rs4237073	AG	34	46.6	AG		31	46.3	GFRA2 rs10088105	GG	15	20.5	GG	14	20.9	GFRA2 rs4567027	GG	05	06.8	GG	04	06.0	GFRA2 rs4237073	GG	25	34.2	GG	16	23.9																																		
		GFRA2 rs4078157	A	61	41.8	A		47		35.1	GFRA2 rs13250096	C	18		12.3	C		06	4.5	GFRA2 rs4739285		A	85	58.2		A	64		47.8	GFRA2 rs4739286	A	43		29.5		A	33	24.6	GFRA2 rs4739286	G			85	58.2	G	87	64.9	GFRA2 rs13250096			G	128	87.7	G	128		95.5		GFRA2 rs4739285	G	61	41.8		G	70		52.2	GFRA2 rs4739286	G		103	70.5	G		101	75.4		GFRA2 rs13250096	AA	13	17.8		AA		8	11.9	GFRA2 rs4739285	AA	24			32.9	AA	19	28.4	GFRA2 rs4739286	AA			04	05.5	AA	06	09.0		GFRA2 rs13250096		AG	35	47.9	AG		31	46.3		GFRA2 rs4739285	AG	37	50.7	AG	26		38.8	GFRA2 rs4739286	AG	35	47.9	AG		21	33.3	GFRA2 rs13250096	GG	25	34.2	GG	28	41.8	GFRA2 rs4739285	GG	12	16.4	GG	22	32.8	GFRA2 rs4739286	GG	34	46.6	GG	40	59.7																	
			GFRA2 rs4567028	A	104	71.2		A		109		81.3	GFRA2 rs6587002		A	99		67.8	A			91	67.9	GFRA2 rs4739217		C	79		55.6		C	62		46.3		GFRA2 rs7014143	A	53		36.3			A	50	37.3	GFRA2 rs7014143	G				42	28.8	G	25	18.7		GFRA2 rs6587002			G	47	32.2		G	43		32.1		GFRA2 rs4739217		G	63	44.4		G	72			53.7	GFRA2 rs7014143	C		93		63.7	C		84	62.7			GFRA2 rs6587002	AA	35	47.9		AA			45	67.2	GFRA2 rs4739217	CC	24				32.9	CC	19	28.4		GFRA2 rs7014143	AA			9	12.3	AA	10	14.9		GFRA2 rs6587002		AG	34	46.6	AG		19	28.4		GFRA2 rs4739217	CG	31	42.5	CG	24		35.8	GFRA2 rs7014143	AC	35	47.9	AC		30	44.8	GFRA2 rs6587002	GG	04	05.5	GG	03	04.5	GFRA2 rs4739217	GG	16	21.9	GG	24	35.8	GFRA2 rs7014143	CC	29	39.7	CC	27	40.3
				GFRA2 rs1128397	A	69		47.3		A		43			32.1	GFRA2 rs6988470		A	77			52.7	A			70	52.2		GFRA3 rs10036665		A	117		80.1			A	105		78.4			GFRA3 rs10952	A	43		29.5				A	36	26.9	GFRA3 rs10952	T					77	52.7	T		91	67.9		GFRA2 rs6988470				G	69	47.3		G	64			47.8		GFRA3 rs10036665		T		29	19.9		T	29				21.6	GFRA3 rs10952	T		103			70.5	T		98	73.1				GFRA2 rs6988470	AA	14	19.2			AA			7	10.4	GFRA3 rs10036665	AA	47				64.4	AA	41	61.2		GFRA3 rs10952	AA			06	08.2	AA	05	07.5		GFRA2 rs6988470		AT	41	56.2	AT		29	43.3		GFRA3 rs10036665	AT	23	31.5	AT	23		34.3	GFRA3 rs10952	AT	31	42.5	AT		26	38.8	GFRA2 rs6988470	TT	18	24.7
GFRA3 rs11242417					A	124	84.9	A		113		84.3		GFRA3 rs7726580	A			50	34.2		A	45	33.6			GFRA4 rs6084432	A	23			15.8	A		23	17.2		GFRA4 rs633924	A		89		61.0		A	79		59.0		GFRA4 rs633924		C	22	15.1		C	21				15.7	GFRA3 rs7726580	G	96	65.8	G					89	66.4	GFRA4 rs6084432	G		123	84.2	G		111				82.8	GFRA4 rs633924	G	87		39.0	G		55		41.0		GFRA3 rs7726580		AA	52		71.2	AA		48	71.6	GFRA4 rs6084432				AA	01	01.4	AA		02			03.0	GFRA4 rs633924		AA	27				37.0	AA	24	35.8			GFRA3 rs7726580			AG	20	27.4	AG	17				25.4	GFRA4 rs6084432	AG	21		28.8	AG			19	28.4	GFRA4 rs633924	AG	35		47.9		AG	31	46.3	GFRA3 rs7726580		GG	01		01.4	CC	02

Figure 1

Hardy-Weinberg equilibrium in case-control sample




Figure 2:

Hardy-Weinberg equilibrium in treatment response sample




Figure 3




Figure 4




Figure S\$






Figure S1









Figure S4



Legends

Table 1: Allelic and genotypic frequencies for the case-control association

Table 1: (continued)

Table 2: Family –based association test. S represents the test statistic for observed number of alleles, E represents the expected value of S under null hypothesis and $\text{Var}(S)$ represents variance between the observed and expected transmission.

Table 3: Allelic and genotypic frequencies for the clozapine response association

Table 3:(continued)

Figure 1: Hardy-Weinberg equilibrium p-values for cases and controls. The solid line represents $p = 0.05$

Figure 2: Hardy-Weinberg equilibrium p-values for the clozapine non-responders and responders. The solid line represents $p = 0.05$

Figure 3: Allelic and genotypic p-values for the case-control association. The solid line represents $p = 0.05$

Figure 4: Allelic and genotypic p-values for the clozapine response association. The solid line represents $p = 0.05$

Figure S1: LD plot for the analyzed markers in GFRA1, GFRA2, GFRA3 and GFRA4 in case-control (A, B, C, D); respectively. Values presented are the D' .

Figure S2: LD plot for the analyzed markers in GFRA1, GFRA2, GFRA3 and GFRA4 in family sample (A, B, C, D) respectively. Values presented are the D' .

Figure S3: LD plot for the analyzed markers in GFRA1, GFRA2, GFRA3 and GFRA4 in response sample (A, B, C, D) respectively. Values presented are the D' .

Figure S4: Interaction dendrogram in case-control (A) and clozapine response (B) samples.

3.3 - Clinical involvement of oxidative stress genes polymorphisms in schizophrenia: influence on the severity of symptoms and response to clozapine treatment

Clinical involvement of oxidative stress genes polymorphisms in schizophrenia:
influence on the severity of symptoms and response to clozapine treatment

RP Souza^{1,2,3}, V Basile¹, T Shinkai⁵, M Tampakeras¹, S Potkin⁶, HY Meltzer⁷, JA Lieberman⁸, MA Romano-Silva^{2,3}, JL Kennedy¹

¹ Neurogenetics Section, CAMH and Dept. of Psychiatry, Univ. of Toronto, Canada; ² Grupo de Pesquisa em Neuropsiquiatria Clínica e Molecular, UFMG, Brazil; ³ Departamento de Saude Mental, Faculdade de Medicina, UFMG, Brazil; ⁴ Department of Psychiatry, University of Occupational and Environmental Health, Japan; ⁵ Brain Imaging Center, University of California, USA; ⁶ Psychiatric Hospital at Vanderbilt University, USA; ⁷ University of North Carolina at Chapel Hill, Department of Psychiatry, USA

Abstract

Abnormal activities of critical antioxidant enzymes and other indices of lipid peroxidation in plasma and red blood cells have been detected in patients with schizophrenia. Other results have shown that oxidative stress may be modulated by clozapine. Based on that and some studies already found different clinical relations between reactive oxygen species and negative and positive symptoms, in the present study, it was studied the association between clinical response and the polymorphism in the GPX1 (Pro197Leu) and MNSOD (Ala16Val) gene in 216 clozapine-treated patients with schizophrenia. No association was found with these two functional polymorphisms and clozapine response change after six months, not even using a gene-gene interaction model. No correlations were found between positive/negative symptoms score and both polymorphisms. Our results present that GPX1 (Pro197Leu) and MNSOD (Ala16Val) polymorphisms seem do not play a central role in the clozapine response, although studies in larger and independent samples are necessary to confirm our findings.

Introduction

Schizophrenia is a devastating psychiatric disorder that affects around 1% of the population. It has been shown to be a complex multifactorial disease with both genetic and environmental influences. Antipsychotic drugs are the best means available for symptomatically treating individuals suffering from schizophrenia; however, there is a significant variability in clinical response to these psychotropic medications. Take, for example, clozapine, the prototype atypical antipsychotic, where only 30–60% of individuals resistant to typical antipsychotics may demonstrate a beneficial clinical response with respect to positive and negative symptomatology (Malhotra et al. 2004)

Human glutathione peroxidase (GPX1; OMIM#138320) is a selenium-dependent enzyme ubiquitously expressed and is found in cytoplasm and mitochondria which plays an important role in detoxification of free radicals (Ursini et al., 1985). GPX1 knockout mice show increased susceptibility to oxidative stress-inducing agents (paraquat and hydrogen peroxide) (de Haan et al., 1998). The human GPX1 gene has been located on

chromosome 3p21.3 (Kiss et al., 1997), and it is composed of two exons within a 1.42 Kb region (Ishida et al., 1987). A single nucleotide polymorphism (SNP) in the GPX1 gene has been reported at nucleotide 593, C to T substitution which causes a proline (Pro) to leucine (Leu) substitution at codon 197 (Pro197Leu) (Forsberg et al., 1999). The effect of the Pro197Leu polymorphism on the function of GPX1 enzyme is considerable. Although one study reported that erythrocyte GPX1 activity showed no significant difference between the genotypes (Forsberg et al., 2000), a more recent study using human transfected cells, which exclusively express either the Pro- or Leu-containing GPX1 allele, showed functional differences between the two alleles (Leu-containing allele was less responsive to the stimulation of GPX1 enzyme activity). (Hu and Diamond, 2003).

Superoxide dismutases (SOD) are the only enzymes that convert superoxide radicals to hydrogen peroxide. The genes encoding these enzymes are located in different chromosomes and in all of them polymorphisms have been described. Copper-zinc SOD (CuZnSOD, SOD1) is encoded on 21q22.1 (OMIM#147450), manganese SOD (MnSOD, SOD2) on 6q25.3 (OMIM#147460), and extracellular SOD (ECSOD, SOD3) on 4p16.3–q21 (OMIM#185490). MnSOD is synthesized in the cytoplasm as a precursor molecule containing a leader signal that is later removed during the transport of the molecule to the mitochondria (Weisiger and Fridovich., 1973ab, Shimoda-Matsubayashi et al., 1996). Ala16Val SNP is common (Val allele frequency approximately 48%) and it has been suggested that it may change the secondary structure and mitochondrial targeting of the protein (Wang et al., 2001).

Abnormal activities of critical antioxidant enzymes (Reddy et al. 1991; Vaiva et al. 1994; Altuntas et al. 2000) and other lipid peroxidation parameters (Mahadik et al. 1995; Kuloghi et al. 2002; Arvindakshan et al., 2003a) in plasma and red blood cells have been detected in patients with schizophrenia. Mahadik found increased lipid peroxidation products and altered defence system in both chronic and drug-naive first episode schizophrenia patients (Mahadik et al., 1996). Zhang performed analyses in some enzymes related with oxidative stress in subjects with schizophrenia including paranoid, disorganized and residual subtypes. They found that activities of SOD and GPX were decreased but levels of malondialdehyde were elevated in patients with a chronic form of schizophrenia as compared with normal controls. SOD and GPX activities were found to be significantly lower in paranoid and residual subtypes compared to both disorganized subtype and the control group. Malondialdehyde levels were significantly higher in all subtypes compared to control group (Zhang et al., 2006).

Although classic antipsychotic drugs such as haloperidol produce a marked reduction in positive symptoms of schizophrenia, they do not improve the negative symptoms such as apathy, confusion, and social withdrawal, nor do they alter the progressive deterioration in the mental abilities of the patient. Meanwhile, atypical antipsychotics have been shown to improve both positive and negative symptoms of schizophrenia, and seem to prevent further worsening of psychotic symptoms (Buckley, 1997; Blin, 1999). The mechanisms by which clozapine exerts its antipsychotic actions in schizophrenia likely involve the blockade of dopamine and serotonin receptors; however, the molecular mechanisms by which clozapine and the other atypical antipsychotics prevent symptom progression remain to be determined. Some results indicate that oxidative stress is integral to this disease and not the result of neuroleptic treatment although antipsychotic-induced oxidative stress results in rat or cell lines are not

conclusive (Polydoro et al., 2004; Reinke et al., 2004; Agostinho et al., 2007; Pillai et al., 2007; Streck et al., 2007).

Several clinical studies that analyzed oxidative parameters and antipsychotic drugs tried to relate these features with tardive dyskinesia. Just some studies had been published relating oxidative stress with clinical response or psychiatric scales scores during the treatment with antipsychotics. Arvindakshan reported reduction in brief psychiatric rating scale (BPRS) and positive and negative syndrome scale (PANSS) score after supplementation with antioxidants agents, such as omega-3 fatty acids, vitamin C, and vitamin E. It was shown that red blood cells SOD is increased in positive schizophrenia (Crow's type I), but not in Crow's type II. This finding means that patients with schizophrenia with positive symptoms are faced with increased oxidative stress indicating that maybe response of oxidative stress could be differently related with positive and negative symptoms (Arvindakshan et al., 2003b). Based on the that oxidative stress could be modulated by clozapine and some studies already found different clinical relations between ROS and negative and positive symptoms, in the present study, we studied the association between clinical response and the polymorphism in the GPX1 (Pro197Leu) and MNSOD (Ala16Val) gene in clozapine-treated patients with schizophrenia.

Methods

Clinical sample

Clinical data from 216 patients with DSM-III-R or DSM-IV diagnoses of schizophrenia, almost all of whom met criteria for treatment refractoriness or intolerance to typical antipsychotic therapy (Souza et al. 2008), were obtained at the following research clinics: Case Western Reserve University in Cleveland, OH (Meltzer, n=100); Hillside Hospital in Glen Oaks, NY (Lieberman, n=87); University of California at Irvine (Potkin, n=29). After informed consent was obtained, patients underwent a washout period of 2 to 4 weeks during which, unless clinically necessary, they received no medications before starting clozapine. Clozapine treatment was continued for a minimum of 6 months during which patients were evaluated prospectively. Clozapine blood levels were monitored through- out the course of treatment to ascertain compliance. Treatment response was evaluated as a % score change using the 18 item Brief Psychiatric Rating Scale (BPRS), a four item (conceptual disorganization, suspiciousness, hallucinations, unusual thought content) positive symptom subscale (BPOS) and a three item (emotional withdrawal, motor retardation, blunted affect) negative symptom subscale (BNEG) after 6 months of clozapine treatment from enrolment into the study (baseline) (with a negative value indicating an improvement in symptoms): % Score Change=(6 Months Score Baseline Score) / (Baseline Score). When treatment response was evaluated as a dichotomous variable in the whole sample at 6 months using criteria based on those of Kane et al. (1988): a reduction of $\geq 20\%$ on the overall score of the BPRS from the baseline score taken at enrolment into the study.

Genetic analyses

Genomic DNA was extracted using the high salt method of Lahiri and Nurnberger (1991). GPX1 genotypes were assessed by the TaqMan allele specific assay method (Applied Biosystems, Foster City, CA) according to the manufacturer's protocols. The Pro197Leu polymorphism site was amplified by polymerase chain reaction (PCR) using the following primers: 5'-CATCGAAGCCCTGCTGTCT-3' (forward) and 5'-CACTGCAACTGCCAAGCA-3' (reverse). Genotyping was performed by 5'-exonuclease fluorescence assay. All genotypes were reported with the allelic discrimination program using the ABI software and confirmed by two experienced researchers. Samples which gave ambiguous calls were genotyped again. MnSOD genotypes were assessed by restriction fragment length polymorphism (RFLP) using the following primers 5'-AGCCCAGCCGTGCGTAGAC-3' and 5'-TACTTCTCCTCGGTGACG-3' and the PCR product was digested with BsaWI enzyme.

Statistical analyses

Individual SNP analyses of responder (control)/non-responder (case) data were performed using χ^2 tests. Individual SNP analyses of % score changes (continuous data) were performed using Analysis of Variance (ANOVA). The statistical program used was the Statistical Package for the Social Sciences, version 10.0.7. The nonparametric Multifactor Dimensionality Reduction (MDR) approach was selected for the analysis of gene-gene interaction (Moore et al., 2006).

Results

Demographic data

Demographic distribution of clinical sites is presented in Table 1. There were no differences observed between the sites in terms of gender ratio, mean age or response ratio. When compared Caucasians and African-American population were not found significant differences in terms of gender ratio, mean age or response ratio.

Genotype data

Significant deviation from Hardy-Weinberg equilibrium was observed for GPX1 polymorphism (non-responders $p=0.002$ and responders $p=0.029$). No deviations were observed for MnSOD polymorphism (non-responders $p=0.606$ and responders $p=0.433$). Responder/non-responder groups were compared for genotype and allele frequencies (see Table 2). In our population, no significant differences were observed for genotype or allele frequency comparisons between responders and non-responders for any of the studied SNPs. No significant association was found either in just the Caucasian or African-American population (data not shown).

Scale score data

BPRS, BPOS and BNEG basal and percentage change score distributions of genotype groups were compared against each other for each SNP (see Table 3).

Lieberman sample scale score data was not available so the results are relative of Meltzer and Potkin samples. No significant associations were observed in any of the scores. No associations were also found either in just the Caucasian or African-American population (data not shown).

Gene-gene interaction

After we did not find any association with single SNP and clozapine response we checked if together these two genes would be able to predispose clozapine response. MDR analyzes showed that when both genes are together there is a synergic effect (see Figure 1) but this did not reach significant association level ($p=0.179$).

Discussion

Over 50 years ago, Hoffer, Osmond and Smythies proposed that schizophrenia may be associated with free radical (i.e., reactive oxygen species, ROS) mediated pathology (Hoffer et al., 1954). Contemporary knowledge in neurochemistry increasingly emphasises the role of free radicals in the genesis of structural and functional changes of neuronal membrane that could be responsible for the beginning or aggravation of some diseases. The nervous system possess high potentials for the initiation of free radical reactions (large amount of unsaturated fatty acids, catecholamines and monoamines), which, relative to other tissues, can cause more damage in the brain and nervous system due to insufficient antioxidative protection and existing intensive aerobic metabolism accompanied with oxygen radical production.

Mahadik found increased lipid peroxidation products in both chronic and drug-naive first episode schizophrenia patients (Mahadik et al., 1996). The effect of oxidative modification of neuronal phospholipids, DNA, and proteins on their function (i.e. membrane transport, loss of mitochondrial energy production, gene expression and, therefore, receptor-mediated phospholipid-dependent signal transduction) may explain altered information processing in schizophrenia and changes in these oxidative process could be attributed to antipsychotic drugs. Although exact mechanism is not known, direct effect of drugs on lipid peroxidation or indirect effect through alteration in superoxide and hydroxy radical formation could not be ruled out. Atypical antipsychotic like clozapine seems to increase 5-hydroxyindol acetic acid (5-HIAA), which is excellent scavenger of hydroxyl and superoxide radicals (Blakely et al. 1984; Liu and Mori 1993). Some *in vitro* and *in vivo* studies in animals indicate that treatment with some atypical antipsychotics may be neuroprotective against oxidative cell injury by inducing antioxidant protection (Li et al., 1999; Parikh et al., 2003; Pillai et al., 2007). Our results could not show any association with GPX1 and MnSOD polymorphisms with clozapine response either isolated or after gene-gene interaction analyze.

BPRS score is significantly reduced after supplementation of vitamin C compared to placebo as substantiated by significant and negative correlation between BPRS score and plasma ascorbic acid levels (Dakhale et al., 2005). Recently, one study demonstrated reduction in BPRS and PANSS and increase in Henrich's quality of life scale score after supplementation with omega-3 fatty acids, vitamin C, and vitamin E (Arvindakshan et al. 2003b). Ascorbic acid is a water-soluble ketoacetone. It plays a crucial role in the

suppression of superoxide radicals by blocking catecholamine autooxidation (Cadet and Brannock, 1997) thereby inhibiting formation of potential toxic by-products such as 6-hydroxy dopamine (6-OHDA), semiquinone, hydrogen peroxide, and hydroxyl radical, eventually leading to neuronal damage in the brain and development of defect symptoms (Cadet and Lohr, 1987). This could be one of the reasons for reduction in BPRS score after vitamin C supplementation. We did not report any significant association connecting BPRS score and percentage change after 6 months of clozapine treatment and GPX1/MnSOD.

It has been shown that patients with schizophrenia with positive symptoms are faced with increased oxidative stress (Pavlovic et al., 2002), however in our sample we could not find any association with these two functional polymorphisms in key oxidative stress enzymes and the positive symptoms or with its change after six months. Sirota related a positive correlation between superoxide generation and negative symptoms in patients with schizophrenia supporting the hypothesis that superoxide anion may participate in the pathogenesis of schizophrenia, as an excess of free radicals could contribute to the deterioration phase of the disease (Sirota et al., 2003). We also did not show correlation with the negative symptoms and these two oxidative stress-related genes. In summary, although some studies reported associations among oxidative stress and schizophrenia clinical features/ antipsychotic activity, in this present study we did not show any correlation linking GPX1 and MnSOD gene polymorphisms and these features. Further studies with a larger sample are necessary to confirm these negative findings.

Acknowledgements

Authors would like to thank Daniela VF Rosa, Mary Smirniw and Nicole King for their help and support throughout manuscript preparation

Reference list

1. Agostinho FR, Jornada LK, Schroder N, Roesler R, Dal-Pizzol F, Quevedo J (2007) Effects of chronic haloperidol and/or clozapine on oxidative stress parameters in rat brain. *Neurochem.Res.*, 32:1343-1350.
2. Altuntas I, Aksoy H, Coskun I, Caykoylu A, Akcay F (2000) Erythrocyte superoxide dismutase and glutathione peroxidase activities, and malondialdehyde and reduced glutathione levels in schizophrenic patients. *Clin.Chem.Lab Med.*, 38:1277-1281.
3. Arvindakshan M, Ghate M, Ranjekar PK, Evans DR, Mahadik SP (2003) Supplementation with a combination of omega-3 fatty acids and antioxidants (vitamins E and C) improves the outcome of schizophrenia. *Schizophr.Res.*, 62:195-204.
4. Arvindakshan M, Sitasawad S, Debsikdar V, Ghate M, Evans D, Horrobin DF, Bennett C, Ranjekar PK, Mahadik SP (2003) Essential polyunsaturated fatty acid

- and lipid peroxide levels in never-medicated and medicated schizophrenia patients. *Biol.Psychiatry*, 53:56-64.
5. Blakely RD, Wages SA, Justice JB, Herndon JG, Neill DB (1984) Neuroleptics increase striatal catecholamine metabolites but not ascorbic acid in dialyzed perfusate. *Brain Res.*, 308:1-8.
 6. Blin O (1999) A comparative review of new antipsychotics. *Can.J.Psychiatry*, 44:235-244.
 7. Buckley PF (1997) New dimensions in the pharmacologic treatment of schizophrenia and related psychoses. *J.Clin.Pharmacol.*, 37:363-378.
 8. Cadet JL, Brannock C (1997) Invited review-free radicals and pathobiology of brain dopamine system. *Neurochem Int* 32:117-131
 9. Cadet JL, Lohr JB (1987) Free radicals and developmental biopathology of schizophrenic burnout. *Integr Psychiatr* 51:40-48
 10. Dakhale GN, Khanzode SD, Khanzode SS, Saoji A (2005) Supplementation of vitamin C with atypical antipsychotics reduces oxidative stress and improves the outcome of schizophrenia. *Psychopharmacology (Berl)*, 182:494-498.
 11. de Haan JB, Bladier C, Griffiths P, Kelner M, O'Shea RD, Cheung NS, Bronson RT, Silvestro MJ, Wild S, Zheng SS, Beart PM, Hertzog PJ, Kola I (1998) Mice with a homozygous null mutation for the most abundant glutathione peroxidase, Gpx1, show increased susceptibility to the oxidative stress-inducing agents paraquat and hydrogen peroxide. *J.Biol.Chem.*, 273:22528-22536.
 12. Forsberg L, de FU, Marklund SL, Andersson PM, Stegmayr B, Morgenstern R (2000) Phenotype determination of a common Pro-Leu polymorphism in human glutathione peroxidase 1. *Blood Cells Mol.Dis.*, 26:423-426.
 13. Hoffer A, Osmond H, Smythies J (1954) Schizophrenia; a new approach. II. Result of a year's research. *J.Ment.Sci.*, 100:29-45.
 14. Hu YJ, Diamond AM (2003) Role of glutathione peroxidase 1 in breast cancer: loss of heterozygosity and allelic differences in the response to selenium. *Cancer Res.*, 63:3347-3351.
 15. Kane J, Honigfeld G, Singer J, Meltzer H (1988) Clozapine for the treatment-resistant schizophrenic. A double-blind comparison with chlorpromazine. *Arch.Gen.Psychiatry*, 45:789-796.
 16. Kiss C, Li J, Szeles A, Gizatullin RZ, Kashuba VI, Lushnikova T, Protopopov AI, Kelve M, Kiss H, Kholodnyuk ID, Imreh S, Klein G, Zabarovsky ER (1997) Assignment of the ARHA and GPX1 genes to human chromosome bands 3p21.3 by in situ hybridization and with somatic cell hybrids. *Cytogenet.Cell Genet.*, 79:228-230.

17. Kuloghi M, Ustundag B, Atmaca M, Canatan H, Tezean AE, Cinkiline N (2002) Lipid peroxidation and antioxidant enzyme levels in patients with schizophrenia and bipolar disorder. *Cell Biochem Funct* 20:171–175
18. Lahiri DK Nurnberger JI, Jr. (1991) A rapid non-enzymatic method for the preparation of HMW DNA from blood for RFLP studies. *Nucleic Acids Res.*, 19:5444
19. Li XM, Chlan-Fourney J, Juorio AV, Bennett VL, Shrikhande S, Keegan DL, Qi J, Boulton AA (1999) Differential effects of olanzapine on the gene expression of superoxide dismutase and the low affinity nerve growth factor receptor. *J.Neurosci.Res.*, 56:72-75.
20. Liu J Mori A (1993) Monoamine metabolism provides an antioxidant defense in the brain against oxidant- and free radical-induced damage. *Arch.Biochem.Biophys.*, 302:118-127.
21. Mahadik SP, Mukherjee S, Correnti I, Sheffer R (1995) Elevated levels of lipid peroxidation products in plasma from drug-naive patients at onset of psychosis. *Schizophr Res* 15:66–70
22. Mahadik SP Scheffer RE (1996) Oxidative injury and potential use of antioxidants in schizophrenia. *Prostaglandins Leukot.Essent.Fatty Acids*, 55:45-54.
23. Malhotra AK, Murphy GM, Jr., Kennedy JL (2004) Pharmacogenetics of psychotropic drug response. *Am.J.Psychiatry*, 161:780-796.
24. Moore JH, Gilbert JC, Tsai CT, Chiang FT, Holden T, Barney N, White BC (2006) A flexible computational framework for detecting, characterizing, and interpreting statistical patterns of epistasis in genetic studies of human disease susceptibility. *J.Theor.Biol.*, 241:252-261.
25. Parikh V, Terry AV, Khan MM, Mahadik SP (2004) Modulation of nerve growth factor and choline acetyltransferase expression in rat hippocampus after chronic exposure to haloperidol, risperidone, and olanzapine. *Psychopharmacology (Berl)*, 172:365-374.
26. Pavlović D, Tamburić V, Stojanović I, Kocić G, Jevtović T, Đorđević V (2002). Oxidative stress as marker of positive symptoms in schizophrenia *Facta Universitatis Medicine and Biology* 9:157-161
27. Pillai A, Parikh V, Terry AV, Jr., Mahadik SP (2007) Long-term antipsychotic treatments and crossover studies in rats: differential effects of typical and atypical agents on the expression of antioxidant enzymes and membrane lipid peroxidation in rat brain. *J.Psychiatr.Res.*, 41:372-386.
28. Polydoro M, Schroder N, Lima MN, Caldana F, Laranja DC, Bromberg E, Roesler R, Quevedo J, Moreira JC, Dal-Pizzol F (2004) Haloperidol- and clozapine-induced oxidative stress in the rat brain. *Pharmacol.Biochem.Behav.*, 78:751-756.

29. Reddy R, Sahebarao MP, Mukherjee S, Murthy JN (1991) Enzymes of the antioxidant defense system in chronic schizophrenic patients. *Biol.Psychiatry*, 30:409-412.
30. Reinke A, Martins MR, Lima MS, Moreira JC, Dal-Pizzol F, Quevedo J (2004) Haloperidol and clozapine, but not olanzapine, induces oxidative stress in rat brain. *Neurosci.Lett.*, 372:157-160.
31. Shimoda-Matsubayashi S, Matsumine H, Kobayashi T, Nakagawa-Hattori Y, Shimizu Y, Mizuno Y (1996) Structural dimorphism in the mitochondrial targeting sequence in the human manganese superoxide dismutase gene. A predictive evidence for conformational change to influence mitochondrial transport and a study of allelic association in Parkinson's disease. *Biochem.Biophys.Res.Commun.*, 226:561-565.
32. Sirota P, Gavrieli R, Wolach B (2003) Overproduction of neutrophil radical oxygen species correlates with negative symptoms in schizophrenic patients: parallel studies on neutrophil chemotaxis, superoxide production and bactericidal activity. *Psychiatry Res.*, 121:123-132.
33. Souza RP, Romano-Silva MA, Lieberman JA, Meltzer HY, Wong AH, Kennedy JL (2008) Association study of GSK3 gene polymorphisms with schizophrenia and clozapine response. *Psychopharmacology (Berl)*,
34. Streck EL, Rezin GT, Barbosa LM, Assis LC, Grandi E, Quevedo J (2007) Effect of antipsychotics on succinate dehydrogenase and cytochrome oxidase activities in rat brain. *Naunyn Schmiedebergs Arch.Pharmacol.*, 376:127-133.
35. Ursini F, Maiorino M, Gregolin C (1985) The selenoenzyme phospholipid hydroperoxide glutathione peroxidase. *Biochim.Biophys.Acta*, 839:62-70.
36. Vaiva G, Thomas P, Leroux JM, Cottencin O, Dutoit D, Erb F, Goudemand M (1994) [Erythrocyte superoxide dismutase (eSOD) determination in positive moments of psychosis]. *Therapie*, 49:343-348.
37. Wang LI, Miller DP, Sai Y, Liu G, Su L, Wain JC, Lynch TJ, Christiani DC (2001) Manganese superoxide dismutase alanine-to-valine polymorphism at codon 16 and lung cancer risk. *J.Natl.Cancer Inst.*, 93:1818-1821.
38. Weisiger RA, Fridovich I (1973) Mitochondrial superoxide simutase. Site of synthesis and intramitochondrial localization. *J.Biol.Chem.*, 248:4793-4796.
39. Weisiger RA, Fridovich I (1973) Superoxide dismutase. Organelle specificity. *J.Biol.Chem.*, 248:3582-3592.
40. Zhang XY, Tan YL, Cao LY, Wu GY, Xu Q, Shen Y, Zhou DF (2006) Antioxidant enzymes and lipid peroxidation in different forms of schizophrenia treated with typical and atypical antipsychotics. *Schizophr.Res.*, 81:291-300.

Figures and tables

Sample (n)	Male/female [n (%)]	Age (mean \pm SD)	Responder/non- responder [n (%)]	Caucasian/ African- American [n(%)]
Meltzer (100)	72/28 (72/28)	33 \pm 9	49/51 (49/51)	74/26 (74/26)
Lieberman (87)	62/25 (71/29)	35 \pm 8	52/35 (60/40)	72/15 (83/17)
Potkin (29)	22/07 (76/23)	35 \pm 7	11/18 (38/62)	25/04 (86/14)
Caucasian (171)	124/47 (73/27)	35 \pm 18	86/85 (50/50)	-
African-American (45)	31/14 (69/31)	34 \pm 10	26/19 (58/42)	-

Table 1: Demographic characteristics and percentage response in each population.

Marker	Allele frequency				Genotype frequency				Association							
	Non-responder		Responder		Non-responder		Responder		Allele		Genotype					
	n	%	n	%	n	%	n	%	Chi-square	p-value	Chi-square	p-value				
GPX1	1	47	27.6	1	48	25.0	11	1	1.2	11	2	2.1	0.33	0.567	1.03	0.595
	2	123	72.4	2	144	75.0	12	45	52.9	12	44	45.8				
							22	39	45.9	22	50	52.1				
MnSOD	1	84	44.7	1	99	48.5	11	20	21.3	11	26	25.5	0.58	0.445	0.57	0.751
	2	104	55.3	2	105	51.5	12	44	46.8	12	47	46.1				
							22	30	31.9	22	29	28.4				

Table 2: Individual SNP non-responder/responder analyses: frequencies and significance levels




Figure 1: Interaction dendrogram

Gene	Parameter	Time point	Geno type	n	Mean	SD	95%CI		F	p-value		
							Lower	Upper				
GPX1	BPRS	basal	11	3	44.67	8.38	23.83	65.50	0.538	0.586		
			12	51	35.94	14.82	31.77	40.11				
			22	52	35.67	14.66	31.59	39.76				
		% 6 months	11	3	-35.90	47.44	-153.7	81.96			0.379	0.686
			12	51	-19.67	29.01	-27.83	-11.51				
			22	51	-20.99	33.03	-30.28	-11.70				
	BPOS	basal	11	3	13.67	5.13	0.92	26.41	0.405	0.668		
			12	51	13.29	6.35	11.51	15.08				
			22	52	12.25	5.94	10.60	13.90				
		% 6 months	11	3	-35.20	56.18	-174.7	104.38			0.206	0.815
			12	47	-20.20	56.53	-36.80	-3.60				
			22	50	-16.09	54.23	-31.51	-0.68				
BNEG	basal	11	3	8.00	5.00	-4.42	20.42	0.479	0.621			
		12	51	6.10	3.71	5.05	7.14					
		22	52	6.79	5.11	5.36	8.21					
	% 6 months	11	3	14.43	84.61	-195.7	224.63			0.734	0.483	
		12	48	11.48	111.3	-20.86	43.82					
		22	50	-9.34	54.85	-24.93	6.25					
MnSOD	BPRS	basal	11	28	39.61	18.26	32.53	46.69	0.406			0.668
			12	48	36.79	13.86	32.77	40.82				
			22	36	36.33	15.24	31.17	41.49				
		% 6 months	11	27	-23.10	30.12	-35.01	-11.18		0.241	0.786	
			12	48	-20.93	29.70	-29.55	-12.30				
			22	36	-17.81	32.15	-28.68	-6.93				
	BPOS	basal	11	28	13.25	6.76	10.63	15.87	0.007			0.993
			12	49	13.14	6.31	11.33	14.96				
			22	36	13.06	6.09	10.99	15.12				
		% 6 months	11	27	-19.10	45.25	-37.00	-1.20		0.180	0.835	
			12	47	-14.93	69.78	-35.42	5.56				
			22	34	-22.50	41.86	-37.11	-7.90				
BNEG	basal	11	28	7.25	5.70	5.04	9.46	0.183	0.833			
		12	49	6.65	4.87	5.25	8.05					
		22	36	7.19	4.36	5.72	8.67					
	% 6 months	11	27	-9.60	60.15	-33.40	14.20			0.519	0.597	
		12	46	11.82	109.7	-20.76	44.41					
		22	35	1.99	69.29	-21.81	25.80					

Table 3: Basal and after 6 months percentage change scores

3.4 - Genetic association study of NALCN polymorphisms with schizophrenia and antipsychotic treatment

Genetic association study of NALCN polymorphisms with schizophrenia and antipsychotic treatment

Renan P Souza^{1,2}, Marco A Romano-Silva¹, Jeffrey A Lieberman⁴, Herbert Y Meltzer⁵, Mei Zhen⁶, Gary Remington², James L Kennedy^{2,3}, Albert HC Wong². ¹Laboratorio de Neurociencia, Dept. Saude Mental, Faculdade de Medicina, Universidade Federal de Minas Gerais, Brazil; ²Neurogenetics Section, CAMH, Toronto, ON, Canada ³Department of Psychiatry, University of Toronto, ON, Canada; ⁴Department of Psychiatry, University of North Carolina, Chapel Hill, NC, USA; ⁵Psychiatric Hospital, Vanderbilt University, Nashville, TN, USA; ⁶Samuel Lunenfeld Research Institute, Mount Sinai Hospital, Toronto, ON, Canada.

Running title: *NALCN* schizophrenia, treatment response and side-effects

Abstract

NALCN (sodium leak channel, non-selective) is a gene located on chromosome 13q in a suggested linkage region for schizophrenia. Mouse *NALCN* mediates some background sodium leak in hippocampal neurons and plays a role in neuronal excitability. Abnormalities in hippocampal activity and neuronal excitability have been implicated in schizophrenia. In this study we analyzed 26 *NALCN* polymorphisms and examined association with four phenotypes: diagnosis of schizophrenia (case-control and family-based analysis), clozapine response, clozapine-induced weight gain and antipsychotic-induced tardive dyskinesia (TD) in schizophrenia patients. We compared allele; genotype and haplotype frequencies in 219 matched case-control subjects, 85 small nuclear families, 150 schizophrenia patients taking clozapine for 6 months, 67 patients with weight gain accessed after 6 weeks of clozapine treatment and 210 patients with antipsychotic-induced TD. In case-control samples we found significant associations with rs9518320 and rs9518331 and haplotypes composed of rs7317836, rs9518320, rs9518331, rs2584531 and rs3916906. We did not find any significantly altered transmission in the schizophrenia family sample. Our results showed rs2152324 to be associated with clozapine response. A haplotype formed by rs10508059-rs7328287-rs496238 showed association with clozapine-induced weight gain. Five individual SNPs (rs9513851, rs9518307, rs9518349, rs10508059 and rs7328287) and haplotypes composed of rs9513851-rs9518307 and rs7328287-rs496238) showed significant associations with TD. Our results suggest that the *NALCN* may affect susceptibility to schizophrenia, antipsychotic response and side-effects.

Keywords: schizophrenia, clozapine response, *NALCN*, ion channels, genetic association, family-based association test.

Introduction

Schizophrenia affects about 1.0% of the population worldwide, with devastating consequences for both patients and their families and is the seventh most costly medical illness (Freedman, 2003). The hallucinations, delusions, thought disorder, and cognitive deficits associated with schizophrenia impact profoundly on the perception, emotion, and judgment of patients. Current treatments are only partially successful, and therefore the development of novel treatments based on an understanding of the etiology and pathogenesis of schizophrenia is imperative. Until recently, progress in schizophrenia research has been limited by a number of factors, including disease heterogeneity and the lack of clear pathological lesions. Evidence increasingly suggests that schizophrenia is a disorder of brain development and plasticity. Genetic studies have recently begun to identify strong candidate risk genes for schizophrenia, and neurobiological studies of the normal and variant forms of these genes are advancing (Owen et al, 2005; Craddock et al, 2005; Chen et al, 2006; Riley and Kendler, 2006; Ross et al, 2006).

Linkage and association studies have implicated several loci in the genome that likely harbor genes conferring risk for schizophrenia. The interpretation of genetic linkage results is controversial and some degree of subjectivity enters into the determination of which regions of the genome should be considered to have truly significant evidence for linkage to schizophrenia. Two meta-analyses have summarized these findings (Badner and Gershon, 2002; Lewis, *et al* 2003). Badner and Gershon (2002) suggested the existence of susceptibility genes on chromosomes 8p, 13q and 22q, however 13q was not supported by Lewis et al (2003). Among other genes, the 13q region contains *G72* (or *DAOA* at 13q33.2). Several individual replication studies and a meta-analysis have supported the association of *G72* with schizophrenia though, as with other loci, the associated alleles and haplotypes are not identical across studies and some polymorphic variants are located outside of the gene (Chumakov et al, 2002; Detera-Wadleigh and McMahon, 2006).

An adjacent region (13q33.1) also contains other genes that have been associated with neuropsychiatric diseases, such as fibroblast growth factor 14 (*FGF14*) (van Swieten et al, 2003; Dalski et al, 2005) and tripeptidyl peptidase II (*TPPII*). (Radu et al, 2006). In this region, 13q33.1, 4.1Mb upstream of *G72* is located *NALCN* (also known as *VGCNLI*). *NALCN* is a highly conserved protein in mammals (99% identity between human and rat). Close homologues are also found in invertebrates. For example, *D. melanogaster* has a single homologous gene named $\alpha 1U$ (for unusual $\alpha 1$ subunit, 57% identity with human *NALCN*) (Littleton and Ganetzky, 2000). Two homologs also exist in *C. elegans* namely *nca-1* and *nca-2*; both with 48% identity to human *NALCN* (Humphrey et al, 2007). Both *nca* and *na* proteins are expressed specifically in the nervous system.

NALCN family proteins display high homology to the alpha-subunit of voltage-gated cation channels. In *Drosophila*, hypomorphic alleles of this family protein, *na*, that result in a reduced protein expression are viable and fertile, but have altered circadian rhythms (Lear et al, 2005; Nash et al, 2002). The *NALCN* mutant flies also have altered sensitivity to volatile anesthetics such as halothane (Krishnan and Nash, 1990; Mir et al., 1997). In *C. elegans*, the *nca* loss of function mutant is also viable and fertile. It has also

been associated with altered sensitivity to halothane (Humphrey et al,2007), as well as altered locomotion patterns and synaptic functions (Jospin et al, 2007; Yeh et al., 2008).

Originally cloned in 1999 from the rat brain, *NALCN* is expressed in many brain regions (Lu et al, 2007) of vertebrates. *NALCN* encodes a voltage-independent, nonselective, non-inactivating cation channel permeable to sodium, potassium and calcium when exogenously expressed in HEK293 cells (Lu et al, 2007). Deletion of *NALCN* in mice results in a severely disrupted respiratory rhythm characterized by periods of apnea and mutant pups die within 24 hours of birth. *In vivo*, the *NALCN* channel appears to be the main source of the background sodium leak in the hippocampal neurons at rest and is important for neuronal excitability (Lu et al, 2007). Both hippocampal activity and neuronal excitability are processes strongly altered in schizophrenia (Saugstad, 1994; Oxley et al, 2004; Eichhammer et al, 2004; Goldman and Mitchell, 2004; Boyer et al, 2007). The effect of this ion channel on the daily rhythms of the fly and mouse are also especially relevant because of the circadian rhythm disturbances observed in schizophrenia and bipolar disorder (Christian et al. 2002). Because the function of *NALCN* is consistent with manifest some schizophrenia symptoms, and its location is within a suggestive chromosomal linkage region for schizophrenia, we hypothesized that *NALCN* may show a genetic association with schizophrenia, or the patient response to schizophrenia-treatment. To test this hypothesis, we performed an association study using both matched case-control and family samples. We also evaluated whether *NALCN* SNPs are associated with the alternate phenotypes clozapine response as well as two important side effects: clozapine-induced weight gain and antipsychotic-induced tardive dyskinesia (TD).

Methods

Clinical sample

All recruitment and clinical assessments were conducted with written informed consent and approval of our institutional ethics review board. Clinical data and DNA samples were obtained from the probands of 85 small nuclear families, as well as 219 patients with a DSM-III-R or DSM-IV diagnosis of schizophrenia. Healthy controls (N=219) were matched for age (± 5 years), ethnicity and gender (146 male and 73 female cases and controls: mean age 36 ± 8). The Structured Clinical Interview for DSM-IV Axis I Disorders (SCID-I) was administered by trained research assistants to each patient and diagnosis was supplemented by a review of medical records. The diagnosis was established via consensus procedures by two of the investigators. Controls were screened for current or past history of major psychiatric disorders or substance misuse, and excluded if either was detected.

For the clozapine response sample, clinical data from 140 patients with a DSM-III-R or DSM-IV diagnosis of schizophrenia, almost all treatment refractory or intolerant of typical antipsychotic therapy (Kane, *et al* 1988), were collected at Case Western Reserve University in Cleveland, OH (n=90) (63 males and 27 females, mean age 36 ± 8) and Hillside Hospital in Glen Oaks, NY (n=50) (33 males and 17 females, mean age 35 ± 8). After informed consent was obtained, recruited patients underwent a washout period of 2 to 4 weeks during which, unless clinically necessary, they received no

medications before starting clozapine. Clozapine treatment was continued for a minimum of 6 months during which patients' response was evaluated prospectively. Clozapine blood levels were monitored during the course of treatment to ascertain compliance. Treatment response was evaluated as the percentage score change on the 18-item Brief Psychiatric Rating Scale (BPRS). Treatment response was expressed as a dichotomous variable in the whole sample at 6 months using criteria based on those of Kane and coinvestigators (Kane, *et al* 1988): a reduction of $\geq 20\%$ on the overall score of the BPRS from the baseline score at enrolment. There were no differences observed between the sites in terms of gender ratio, mean age, mean age of onset or response ratio. The Caucasian and African-American groups were not significantly different in terms of gender ratio, mean age, mean age of onset or response ratio (data not shown) (Hwang, *et al*, 2005).

For the clozapine-induced weight gain sample (n=67) were recruited from Case Western Reserve (38 males and 18 females, mean age 35 ± 8) and from Hillside Hospital (4 males and 7 females, mean age 33 ± 6). Weight gain at 6 weeks was expressed as a dichotomous variable using as criteria a weight increase $\geq 7\%$ from baseline at enrolment. The FDA has established this threshold as producing a clinically meaningful and significant metabolic outcome.

For the TD samples, 210 subjects were recruited from four clinical sites: the Centre for Addiction and Mental Health in Toronto, ON (n=109; 71 males and 38 females, mean age 42 ± 10); Case Western Reserve (n=60; 44 males and 16 females, mean age 33 ± 10) and the Hillside Hospital (n=41; 25 males and 16 females, mean age 35 ± 7). Regarding ethnicity, 16 subjects from Case Western and 38 from the Centre for Addiction and Mental Health were African-American; all other subjects were Caucasian. Subjects were selected with the same criteria used for the case-control sample. All patients had at least 1 year of cumulative treatment with typical antipsychotics. For the Case Western Reserve and Hillside Hospital samples, the presence or absence of TD was evaluated before any atypical antipsychotic administration. Patients in the Toronto TD sample were on various typical and atypical antipsychotics when TD was evaluated. TD was assessed using AIMS or the modified Hillside Simpson Dyskinesia Scale (HSDS) for patients recruited from the Hillside Hospital (Basile *et al.*, 1999). The seven body area items and the overall global score of HSDS match those of AIMS; thus, TD assessment could be compared across all sites.

Genetic analyses

Genomic DNA was extracted using the high salt method as previously described (Lahiri and Nurnberger, 1991). We analyzed 25 SNPs in *NALCN* (rs1289556, rs9554752, rs17677552, rs686141, rs12867417, rs658213, rs614728, rs9513851, rs9518307, rs12584031, rs1452112, rs7317836, rs9518320, rs9518331, rs2584531, rs12869164, rs3916906, rs9518349, rs10508059, rs7328287, rs496238, rs7318529, rs9554772, rs17486808, rs17584161) and 1 SNP before promoter region (rs2152324). Genotyping was performed by GoldenGate assay (Illumina, San Diego, CA, USA) at The Centre for Applied Genomics (TCAG), Hospital for Sick Children in Toronto, ON, Canada.

Statistical analyses

Individual SNP analyses of case (schizophrenia patients; clozapine responders; weight gain greater than 7%; TD) and control (healthy controls; clozapine non-responders; absence of weight gain greater than 7%; no TD) data and Hardy–Weinberg equilibrium assessment were performed using χ^2 tests. The statistical program used was the Statistical Package for the Social Sciences, version 10.0.7 (SPSS 2000) for genotypic association and Haploview 4.0 (Barrett, 2005) for allelic association. We applied the family-based association test (FBAT, version 1.0, Laird, *et al* 2000) under the assumption of an additive model, and PEDSTATS (Wigginton and Abecasis, 2005) for Hardy–Weinberg equilibrium in the family data. Linkage disequilibrium (LD) was assessed using Haploview, version 4.0. Haplotype analyses were performed using UNPHASED 3.0.10 (Dudbridge, 2003), Haploview version 4.0 and FBAT.

Results

Linkage disequilibrium analysis

Pairwise LD between the SNPs is presented for each gene (Figure S1). In this study, we defined a haplotype block as a region over which less than 5% of pairwise comparisons among informative SNPs showed strong evidence of historical recombination (upper confidence bound on D' less than 0.9; Gabriel, *et al* 2002). Based on this definition we found 5 LD blocks in the case-control sample, 7 in the family sample, 4 in the clozapine response sample, 3 in the clozapine-induced weight gain sample and 7 in the TD sample (figure S1).

Genotype data

Case-control sample: Significant deviation from Hardy–Weinberg equilibrium was observed for rs7318529 ($p = 0.001$) in control samples (Figure 1A). The cases and controls were compared for genotype and allele frequencies across the markers (Table 1 and Figure 1B). Significant associations were observed for: rs9518320 (allele $p = 0.005$, $X^2 = 7.67$; genotype $p = 0.022$, $X^2 = 7.64$) and rs9518331 (allele $p = 0.006$, $X^2 = 7.42$; genotype $p = 0.029$, $X^2 = 7.10$).

Family sample: Three SNPs showed significant deviation from Hardy–Weinberg equilibrium: rs9518320 ($p = 0.023$), rs9518331 ($p = 0.022$) and rs3916906 ($p = 0.032$), analyzing 80 unrelated individuals (Figure 1A). No significant associations were observed (Table 1 and Figure 1B).

Clozapine response sample: Significant deviations from Hardy–Weinberg equilibrium were observed for rs1758416 ($p = 0.032$) and rs2152324 ($p = 0.029$) in the non-responder group. The following markers deviated significantly from Hardy–Weinberg equilibrium in the responder group: rs1289556 ($p = 0.010$), rs9513851 ($p = 0.043$); rs9518307 ($p = 0.043$) and rs496238 ($p = 0.044$) (Figure 1A). Responder/non-responder groups were compared for genotype and allele frequencies across the markers (Table 1 and Figure 1B) and rs2152324 was associated with treatment response (allele $p = 0.030$, $X^2 = 6.98$; genotype $p = 0.061$, $X^2 = 3.49$).

Clozapine-induced weight gain sample: Significant deviations from Hardy–Weinberg equilibrium were observed for rs1050805 ($p = 0.009$) and rs2152324 ($p = 0.023$) in the patients with weight gain (Figure 1A). Patients with and without weight gain were compared for genotype and allele frequencies across the markers (see Table 1 and Figure 4) and no significant association was observed (Table 1 and Figure 1B).

Tardive dyskinesia sample: Two SNPs deviated significantly from Hardy–Weinberg equilibrium in patients without TD (rs9513581, $p = 0.001$ and rs10508059, $p = 0.003$), and one in patients with TD (rs9518307, $p = 0.0006$) (Figure 1A). Comparisons of allele and genotype frequencies revealed significant associations with five SNPs: rs9513851 (allele $p = 0.008$, $X^2 = 9.75$; genotype $p = 0.031$, $X^2 = 4.60$), rs9518307 (allele $p = 0.053$, $X^2 = 5.87$; genotype $p = 0.049$, $X^2 = 3.86$), rs9518349 (allele $p = 0.042$, $X^2 = 6.33$; genotype $p = 0.308$, $X^2 = 1.03$), rs10508059 (allele $p = 0.032$, $X^2 = 6.86$; genotype $p = 0.097$, $X^2 = 2.74$) and rs7328287 (allele $p = 0.079$, $X^2 = 5.08$; genotype $p = 0.030$, $X^2 = 4.67$) (Table 1 and Figure 1B).

Haplotype analysis

Case-control sample: Cases and controls were compared for haplotype frequencies. We did not find associations when considering haplotypes in the same LD block. We then performed three-marker sliding-window haplotype analysis across the whole gene in order to better characterize regions associated with our phenotypes. The following haplotypes showed association rs9518307-rs12584031-rs1452112 (global $p = 0.015$, $X^2 = 13.98$); rs7317836-rs9518320-rs9518331 (global $p = 0.022$, $X^2 = 14.69$; C-A-C control frequency = 0.439, case frequency = 0.371, $p = 0.027$, $X^2 = 4.85$; C-G-C control frequency = 0.234, case frequency = 0.320, $p = 0.005$, $X^2 = 7.78$); rs9518320-rs9518331- rs2584531 (global $p = 0.047$, $X^2 = 14.21$) and rs9518331-rs2584531-rs3916906 (global $p = 0.045$, $X^2 = 12.87$; A-G-A control frequency = 0.239, case frequency = 0.321, $p = 0.008$, $X^2 = 6.82$). The region including SNPs rs7317836, rs9518320, rs9518331, rs2584531 and rs3916906 showed significant associations for all three-marker sliding windows, and so we performed haplotype analysis with all these markers. The C-G-A-G-A haplotype showed significant association with schizophrenia (control frequency = 0.216, case frequency = 0.300, $p = 0.004$, $X^2 = 8.01$).

Family sample: There were no associations with haplotypes in the same LD block or three-marker sliding windows in the family-based sample.

Clozapine response sample: Clozapine responders and non-responders were compared for haplotype frequencies. No haplotypes were associated with clozapine response, either in the same LD block or in three-marker sliding windows.

Clozapine-induced weight gain sample: Subjects were compared for haplotype frequencies relative to weight gain (<7% or $\geq 7\%$), and no significant associations were detected with markers within the same LD block. Three-marker sliding window analysis showed one significant association: rs10508059-rs7328287-rs496238 (global $p = 0.034$, $X^2 = 10.40$; A-A-A control frequency = 0.163, case frequency = 0.030, $p = 0.042$, $X^2 = 4.11$).

TD sample: Subjects were compared for haplotype frequencies relative to the presence or absence of TD. Considering haplotypes in the same LD block, within block 3 (rs9513851 and rs9518307) we found association between haplotype C-A (TD absent

frequency = 0.922, TD present frequency = 0.969, $p = 0.049$, $X^2 = 3.86$; after 1,000 permutations $p = 0.589$) and A-G (TD absent frequency = 0.074., TD present frequency =, $p = 0.025$, $X^2 = 4.60$; after 1,000 permutations $p = 0.430$). Inside block 6 (rs7328287 and rs496238), the G-G haplotype showed association (TD absent frequency = 0.535, TD present frequency = 0.642, $p = 0.030$, $X^2 = 4.67$; after 1,000 permutations $p = 0.409$). No associations were found in the three marker sliding window analysis.

Discussion

This exploratory study examined the association of 26 SNPs in *NALCN* and four phenotypes: diagnosis of schizophrenia (using case-control and family-based analysis), clozapine response, clozapine-induced weight gain and antipsychotic-induced TD in schizophrenia patients. In case-control samples we found significant associations with rs9518320 and rs9518331. Furthermore, a 76.7Kb region containing rs7317836, rs9518320, rs9518331, rs2584531 and rs3916906 showed significant associations for all three-marker sliding window haplotypes. This region contains 186 SNPs in the CEPH HapMap population (Utah Residents with Northern and Western European Ancestry). Further analyses in this region are required to strengthen this association hypothesis. We did not find significantly altered transmission patterns with schizophrenia in our family sample.

Variation in individual clinical response to psychotropic drug treatment remains a critical problem in the management of serious mental illness (Basile, *et al* 2002; Malhotra, *et al* 2004). We examined if *NALCN* might play a role in treatment response and its side effects. Our results showed one SNP (rs2152324) with nominal association with clozapine response. One haplotype (rs10508059-rs7328287-rs496238 A-A-A) showed association with clozapine-induced weight gain. Regarding TD, five individual SNPs (rs9513851, rs9518307, rs9518349, rs10508059 and rs7328287) and three haplotypes (two with rs9513851-rs9518307 and one with rs7328287-rs496238) showed significant associations.

Corrections for multiple testing have been a controversial issue (Aickin, 1999; Bender and Lange, 1999; Perneger, 1998) and considering the exploratory nature of this study, without prespecified hypotheses for most of our SNPs, we assume no clear structure in the multiple tests. Therefore, our statistically significant results should properly be regarded as “exploratory”, with confirmatory studies needed (Bender and Lange, 1999). Bonferroni correction for multiple testing on individual SNP associations renders all the associations non-significant (Bonferroni corrected $p < 0.0019$). Nyholt (2004) correction was not performed because the gene size and low LD level observed among analyzed markers.

There were SNPs that deviated from Hardy-Weinberg equilibrium in the clozapine response and antipsychotic-induced TD samples, as well as one SNP (rs7318529 $p = 0.001$) in healthy controls. We applied X^2 to test for genetic association in these SNPs, but for low genotype counts the Fisher exact test, which does not rely on the X^2 null distribution approximation, is more appropriate (Guo and Thompson, 1992; Wigginton *et al.* 2005). With the Fisher exact test to evaluate Hardy-Weinberg equilibrium, we found a number of significant associations (rs1289556 in non-responders $p = 0.011$; rs9513851 in non-responders $p = 0.151$ and TD absent $p = 0.037$; rs9518307 in

non-responders $p = 0.151$ and TD present $p = 0.061$; rs10808059 in weight gain absent $p = 0.019$ and TD absent $p = 0.007$; rs496238 in responders $p = 0.043$; rs7318529 in healthy controls $p = 0.011$; rs17584161 in non-responders $p = 0.032$ and weight gain absent $p = 0.029$; rs2152324 in non-responders $p = 0.192$). Assuming a significance threshold of $p < 0.01$, one SNP remains deviated in the TD control sample (rs10508059). Trikalinos et al (2006) concluded that Hardy-Weinberg equilibrium should be routinely and transparently assessed in gene-disease association studies. Furthermore, discrepant results in these analyses do not necessarily mean that the observed association should be dismissed, but indicate the need for more evidence and validation. It is possible that deviations from HWE reflect disease associations (Nielsen, *et al* 1998; Wittke-Thompson, *et al* 2005; Balding, 2006).

NALCN is a 361.1Kbp gene with 44 exons located at chromosome 13q in a suggested linkage region for schizophrenia (Chumakov et al, 2002; Badner and Gershon, 2002; Christian et al, 2002). It has been shown that *NALCN* mRNA is expressed in the cerebral cortex and hippocampus in all neurons and layers, and in all neurons of the spinal cord (dorsal and ventral horns). *NALCN* mRNA expression was not detected in liver, muscle, lung, kidney, or testis (Lee et al, 1999; Lu et al, 2007). *NALCN* mutant mouse neonates do not display gross abnormalities in embryonic development, righting responses, spontaneous limb movement, and toe/tail pinch responses, but do not survive beyond 24 hours after birth. Thus, *NALCN* is one of the few members of the family of four homologous repeat (domains I–IV), six transmembrane segment (S1–S6) (4x6TM) ion channels that are required for neonatal survival. Unlike any of the other 20 members in the 4x6TM channel family, *NALCN* forms a voltage-independent and non-inactivating cation channel (Lu et al, 2007).

NALCN's functions in establishing levels of neuronal excitability and thus controlling firing rates (Lu et al, 2007). Proteins that affect patterns of neuronal firing might play a part in the pathogenesis of schizophrenia (Miller et al, 2001; Mansvelder et al, 2006), and other examples include the human calcium-activated potassium channel (*SKCa3* also known as *KCNN3* and *SK3*) and cholinergic nicotinic receptors (nAChRs) (Freedman et al, 1995; Leonard et al, 1996; Chandy et al, 1998; Glatt et al, 2003; De Luca et al, 2004). Interesting questions for the future are whether *NALCN* expression or activity is altered by signal transduction events, neuronal plasticity or antipsychotic drugs (Lu et al, 2007). In *C. elegans*, interesting findings with NCA channels have been reported regarding neuronal activity. Loss of NCA channel activity leads to locomotion deficit called fainters, which fail to sustain active locomotion (Humphrey et al, 2007). These “fainters” partially suppress the locomotor, vesicle depletion, and electrophysiological defects of synaptojanin mutants. Suppressor loci include the genes for the NCA ion channels (Humphrey et al, 2007), which are homologs of the vertebrate cation leak channel *NALCN* (Lu et al, 2007). It was suggested that activation of the NCA ion channel in synaptojanin mutants leads to defects in recycling of synaptic vesicles (Jospin et al (2007). The synaptojanin gene (*SYNJI*) has been evaluated in bipolar disorder (Saito et al, 2001; Stopkova et al, 2004), and it resides in a susceptibility region for schizophrenia (21q22) (Murtagh et al, 2005; Mirnics et al, 2000). These findings suggest that alterations in *NALCN* activity may play a role in neuronal plasticity. Together with our results, suggest that the *NALCN* may be involved with the

manifestation of schizophrenia symptoms and associated with antipsychotic-induced TD, but further work is clearly required to confirm this hypothesis.

Disclosure/Conflicts of interest

Mr. Souza, Dr. Romano-Silva, Dr. Zhen and Dr. Wong have nothing to declare. Dr. Lieberman has served as a consultant/ advisor or grantee of Acadia, Astra Zeneca, Bristol-Myers Squibb, Eli Lilly, GlaxoSmithKline, Janssen Pharmaceutica, Lundbeck, Merck, Organon, Pfizer and Wyeth; and holds a patent from Repligen. Dr. Meltzer declares that he is a consultant or grantee of Abbott, Acadia, ARYx, Astra Zeneca, Bristol Myers Squibb, Eli Lilly, Janssen, Memory, Minster, Organon, Pfizer, Solvay, Wyeth, and Vanda. Dr. Kennedy declares that he is a consultant for GlaxoSmithKline.

Acknowledgements

Funding and grants: 1) CNPq – Brazil (#202447/2006-5; #140950/2005-2; #554496/2005-4); 2) National Institutes of Health (NIH); 3) Canadian Institutes of Health Research (CIHR) #940595, 4) Ontario Mental Health Foundation (OMHF) and 5) Fapemig – Brazil. The authors would like to thank Daniela VF Rosa, Mary Smirniw and Nicole King for their help and support throughout the manuscript preparation.

Reference list

Aickin M (1999) Other method for adjustment of multiple testing exists. *BMJ* 318: 127-128

Badner JA, Gershon ES (2002) Meta-analysis of whole-genome linkage scans of bipolar disorder and schizophrenia. *Mol.Psychiatry* 7: 405-411

Balding DJ (2006) A tutorial on statistical methods for population association studies. *Nat.Rev.Genet.* 7: 781-791

Barrett JC, Fry B, Maller J, Daly MJ (2005) Haploview: analysis and visualization of LD and haplotype maps. *Bioinformatics.* 21: 263-265

Basile VS, Masellis M, Badri F, Paterson AD, Meltzer HY, Lieberman JA, Potkin SG, Macciardi F, Kennedy JL (1999) Association of the MscI polymorphism of the dopamine D3 receptor gene with tardive dyskinesia in schizophrenia. *Neuropsychopharmacology* 21: 17-27

Basile VS, Masellis M, Potkin SG, Kennedy JL (2002) Pharmacogenomics in schizophrenia: the quest for individualized therapy. *Hum.Mol.Genet.* 11: 2517-2530

Bender R, Lange S (1999) Multiple test procedures other than Bonferroni's deserve wider use. *BMJ* 318: 600-601

Boyer P, Phillips JL, Rousseau FL, Ilivitsky S (2007) Hippocampal abnormalities and memory deficits: new evidence of a strong pathophysiological link in schizophrenia. *Brain Res.Rev.* 54: 92-112

Chandy KG, Fantino E, Wittekindt O, Kalman K, Tong LL, Ho TH, Gutman GA, Crocq MA, Ganguli R, Nimgaonkar V, Morris-Rosendahl DJ, Gargus JJ (1998) Isolation of a novel potassium channel gene hSKCa3 containing a polymorphic CAG repeat: a candidate for schizophrenia and bipolar disorder? *Mol.Psychiatry* 3: 32-37

Chen J, Lipska BK, Weinberger DR (2006) Genetic mouse models of schizophrenia: from hypothesis-based to susceptibility gene-based models. *Biol.Psychiatry* 59: 1180-1188

Christian SL, McDonough J, Liu Cy CY, Shaikh S, Vlamakis V, Badner JA, Chakravarti A, Gershon ES (2002) An evaluation of the assembly of an approximately 15-Mb region on human chromosome 13q32-q33 linked to bipolar disorder and schizophrenia. *Genomics* 79: 635-656

Chumakov I, Blumenfeld M, Guerassimenko O, Cavarec L, Palicio M, Abderrahim H, Bougueleret L, Barry C, Tanaka H, La RP, Puech A, Tahri N, Cohen-Akenine A, Delabrosse S, Lissarrague S, Picard FP, Maurice K, Essioux L, Millasseau P, Grel P, Debailleul V, Simon AM, Caterina D, Dufaure I, Malekzadeh K, Belova M, Luan JJ, Bouillot M, Sambucy JL, Primas G, Saumier M, Boubkiri N, Martin-Saumier S, Nasroune M, Peixoto H, Delaye A, Pinchot V, Bastucci M, Guillou S, Chevillon M, Sainz-Fuertes R, Meguenni S, urich-Costa J, Cherif D, Gimalac A, Van DC, Gauvreau D, Ouellette G, Fortier I, Raelson J, Sherbatich T, Riazanskaia N, Rogaev E, Raeymaekers P, Aerssens J, Konings F, Luyten W, Macciardi F, Sham PC, Straub RE, Weinberger DR, Cohen N, Cohen D (2002) Genetic and physiological data implicating the new human gene G72 and the gene for D-amino acid oxidase in schizophrenia. *Proc.Natl.Acad.Sci.U.S.A* 99: 13675-13680

Craddock N, O'Donovan MC, Owen MJ (2006) Genes for schizophrenia and bipolar disorder? Implications for psychiatric nosology. *Schizophr.Bull.* 32: 9-16

Dalski A, Atici J, Kreuz FR, Hellenbroich Y, Schwinger E, Zuhlke C (2005) Mutation analysis in the fibroblast growth factor 14 gene: frameshift mutation and polymorphisms in patients with inherited ataxias. *Eur.J.Hum.Genet.* 13: 118-120

De L, V, Wong AH, Muller DJ, Wong GW, Tyndale RF, Kennedy JL (2004) Evidence of association between smoking and alpha7 nicotinic receptor subunit gene in schizophrenia patients. *Neuropsychopharmacology* 29: 1522-1526

Detera-Wadleigh SD, McMahon FJ (2006) G72/G30 in schizophrenia and bipolar disorder: review and meta-analysis. *Biol.Psychiatry* 60: 106-114

Dudbridge F (2003) Pedigree disequilibrium tests for multilocus haplotypes. *Genet.Epidemiol.* 25: 115-121

Eichhammer P, Wiegand R, Kharraz A, Langguth B, Binder H, Hajak G (2004) Cortical excitability in neuroleptic-naive first-episode schizophrenic patients. *Schizophr.Res.* 67: 253-259

Freedman R, Hall M, Adler LE, Leonard S (1995) Evidence in postmortem brain tissue for decreased numbers of hippocampal nicotinic receptors in schizophrenia. *Biol.Psychiatry* 38: 22-33

Freedman R (2003) Schizophrenia. *N.Engl.J.Med.* 349: 1738-1749

Gabriel SB, Schaffner SF, Nguyen H, Moore JM, Roy J, Blumenstiel B, Higgins J, DeFelice M, Lochner A, Faggart M, Liu-Cordero SN, Rotimi C, Adeyemo A, Cooper R, Ward R, Lander ES, Daly MJ, Altshuler D (2002) The structure of haplotype blocks in the human genome. *Science* 296: 2225-2229

Glatt SJ, Faraone SV, Tsuang MT (2003) CAG-repeat length in exon 1 of KCNN3 does not influence risk for schizophrenia or bipolar disorder: a meta-analysis of association studies. *Am.J.Med.Genet.B Neuropsychiatr.Genet.* 121: 14-20

Goldman MB, Mitchell CP (2004) What is the functional significance of hippocampal pathology in schizophrenia? *Schizophr.Bull.* 30: 367-392

Guo SW, Thompson EA (1992) Performing the exact test of Hardy-Weinberg proportion for multiple alleles. *Biometrics* 48: 361-372

Humphrey JA, Hamming KS, Thacker CM, Scott RL, Sedensky MM, Snutch TP, Morgan PG, Nash HA (2007) A putative cation channel and its novel regulator: cross-species conservation of effects on general anesthesia. *Curr.Biol.* 17: 624-629

Hwang R, Shinkai T, De L, V, Muller DJ, Ni X, Macciardi F, Potkin S, Lieberman JA, Meltzer HY, Kennedy JL (2005) Association study of 12 polymorphisms spanning the dopamine D(2) receptor gene and clozapine treatment response in two treatment refractory/intolerant populations. *Psychopharmacology (Berl)* 181: 179-187

Jospin M, Watanabe S, Joshi D, Young S, Hamming K, Thacker C, Snutch TP, Jorgensen EM, Schuske K (2007) UNC-80 and the NCA ion channels contribute to endocytosis defects in synaptojanin mutants. *Curr.Biol.* 17: 1595-1600

Kane J, Honigfeld G, Singer J, Meltzer H (1988) Clozapine for the treatment-resistant schizophrenic. A double-blind comparison with chlorpromazine. *Arch.Gen.Psychiatry* 45: 789-796

Krishnan KS, Nash HA (1990) A genetic study of the anesthetic response: mutants of *Drosophila melanogaster* altered in sensitivity to halothane. *Proc.Natl.Acad.Sci.U.S.A* 87: 8632-8636

Lahiri DK, Nurnberger JI, Jr. (1991) A rapid non-enzymatic method for the preparation of HMW DNA from blood for RFLP studies. *Nucleic Acids Res.* 19: 5444

Laird NM, Horvath S, Xu X (2000) Implementing a unified approach to family-based tests of association. *Genet.Epidemiol.* 19 Suppl 1: S36-S42

Lear BC, Lin JM, Keath JR, McGill JJ, Raman IM, Allada R (2005) The ion channel narrow abdomen is critical for neural output of the *Drosophila* circadian pacemaker. *Neuron* 48: 965-976

Lee JH, Cribbs LL, Perez-Reyes E (1999) Cloning of a novel four repeat protein related to voltage-gated sodium and calcium channels. *FEBS Lett.* 445: 231-236

Leonard S, Adams C, Breese CR, Adler LE, Bickford P, Byerley W, Coon H, Griffith JM, Miller C, Myles-Worsley M, Nagamoto HT, Rollins Y, Stevens KE, Waldo M, Freedman R (1996) Nicotinic receptor function in schizophrenia. *Schizophr.Bull.* 22: 431-445

Lewis CM, Levinson DF, Wise LH, DeLisi LE, Straub RE, Hovatta I, Williams NM, Schwab SG, Pulver AE, Faraone SV, Brzustowicz LM, Kaufmann CA, Garver DL, Gurling HM, Lindholm E, Coon H, Moises HW, Byerley W, Shaw SH, Mesen A, Sherrington R, O'Neill FA, Walsh D, Kendler KS, Ekelund J, Paunio T, Lonnqvist J, Peltonen L, O'Donovan MC, Owen MJ, Wildenauer DB, Maier W, Nestadt G, Blouin JL, Antonarakis SE, Mowry BJ, Silverman JM, Crowe RR, Cloninger CR, Tsuang MT, Malaspina D, Harkavy-Friedman JM, Svrakic DM, Bassett AS, Holcomb J, Kalsi G, McQuillin A, Brynjolfson J, Sigmundsson T, Petursson H, Jazin E, Zoega T, Helgason T (2003) Genome scan meta-analysis of schizophrenia and bipolar disorder, part II: Schizophrenia. *Am.J.Hum.Genet.* 73: 34-48

Littleton JT, Ganetzky B (2000) Ion channels and synaptic organization: analysis of the *Drosophila* genome. *Neuron* 26: 35-43

Lu B, Su Y, Das S, Liu J, Xia J, Ren D (2007) The neuronal channel NALCN contributes resting sodium permeability and is required for normal respiratory rhythm. *Cell* 129: 371-383

Malhotra AK, Murphy GM, Jr., Kennedy JL (2004) Pharmacogenetics of psychotropic drug response. *Am.J.Psychiatry* 161: 780-796

Mansvelder HD, van Aerde KI, Couey JJ, Brussaard AB (2006) Nicotinic modulation of neuronal networks: from receptors to cognition. *Psychopharmacology (Berl)* 184: 292-305

Miller MJ, Rauer H, Tomita H, Rauer H, Gargus JJ, Gutman GA, Cahalan MD, Chandy KG (2001) Nuclear localization and dominant-negative suppression by a mutant SKCa3 N-terminal channel fragment identified in a patient with schizophrenia. *J.Biol.Chem.* 276: 27753-27756

Mir B, Iyer S, Ramaswami M, Krishnan KS (1997) A genetic and mosaic analysis of a locus involved in the anesthesia response of *Drosophila melanogaster*. *Genetics* 147: 701-712

Mirnics K, Middleton FA, Marquez A, Lewis DA, Levitt P (2000) Molecular characterization of schizophrenia viewed by microarray analysis of gene expression in prefrontal cortex. *Neuron* 28: 53-67

Murtagh A, McTigue O, Ramsay L, Hegarty AM, Green AJ, Stallings RL, Corvin A (2005) Interstitial deletion of chromosome 21q and schizophrenia susceptibility. *Schizophr.Res.* 78: 353-356

Nash HA, Scott RL, Lear BC, Allada R (2002) An unusual cation channel mediates photic control of locomotion in *Drosophila*. *Curr.Biol.* 12: 2152-2158

Nielsen DM, Ehm MG, Weir BS (1998) Detecting marker-disease association by testing for Hardy-Weinberg disequilibrium at a marker locus. *Am.J.Hum.Genet.* 63: 1531-1540

Nyholt DR (2004) A simple correction for multiple testing for single-nucleotide polymorphisms in linkage disequilibrium with each other. *Am.J.Hum.Genet.* 74: 765-769

Owen MJ, Craddock N, O'Donovan MC (2005) Schizophrenia: genes at last? *Trends Genet.* 21: 518-525

Oxley T, Fitzgerald PB, Brown TL, de CA, Daskalakis ZJ, Kulkarni J (2004) Repetitive transcranial magnetic stimulation reveals abnormal plastic response to premotor cortex stimulation in schizophrenia. *Biol.Psychiatry* 56: 628-633

Perneger TV (1998) What's wrong with Bonferroni adjustments. *BMJ* 316: 1236-1238

Radu D, Tomkinson B, Zachrisson O, Weber G, de BJ, Hirsch S, Lindfors N (2006) Overlapping regional distribution of CCK and TPPII mRNAs in Cynomolgus monkey brain and correlated levels in human cerebral cortex (BA 10). *Brain Res.* 1104: 175-182

Riley B, Kendler KS (2006) Molecular genetic studies of schizophrenia. *Eur.J.Hum.Genet.* 14: 669-680

Ross CA, Margolis RL, Reading SA, Pletnikov M, Coyle JT (2006) Neurobiology of schizophrenia. *Neuron* 52: 139-153

Saito T, Guan F, Papolos DF, Lau S, Klein M, Fann CS, Lachman HM (2001) Mutation analysis of SYNJ1: a possible candidate gene for chromosome 21q22-linked bipolar disorder. *Mol.Psychiatry* 6: 387-395

Saugstad LF (1994) Deviation in cerebral excitability: possible clinical implications. *Int.J.Psychophysiol.* 18: 205-212

Stopkova P, Vevera J, Paclt I, Zukov I, Lachman HM (2004) Analysis of SYNJ1, a candidate gene for 21q22 linked bipolar disorder: a replication study. *Psychiatry Res.* 127: 157-161

Trikalinos TA, Salanti G, Khoury MJ, Ioannidis JP (2006) Impact of violations and deviations in Hardy-Weinberg equilibrium on postulated gene-disease associations. *Am.J.Epidemiol.* 163: 300-309

van Swieten JC, Brusse E, de Graaf BM, Krieger E, van de GR, de K, I, Maat-Kievit A, Leegwater P, Dooijes D, Oostra BA, Heutink P (2003) A mutation in the fibroblast growth factor 14 gene is associated with autosomal dominant cerebellar ataxia [corrected]. *Am.J.Hum.Genet.* 72: 191-199

Wigginton JE, Abecasis GR (2005) PEDSTATS: descriptive statistics, graphics and quality assessment for gene mapping data. *Bioinformatics.* 21: 3445-3447

Wigginton JE, Cutler DJ, Abecasis GR (2005) A note on exact tests of Hardy-Weinberg equilibrium. *Am.J.Hum.Genet.* 76: 887-893

Wittke-Thompson JK, Pluzhnikov A, Cox NJ (2005) Rational inferences about departures from Hardy-Weinberg equilibrium. *Am.J.Hum.Genet.* 76: 967-986

Yeh E, Ng S, Zhang M, Bouhours M, Wang Y, Wang M, Hung W, Aoyagi K, Melnik-Martinez K, Li M, Liu F, Schafer WR, Zhen M (2008) A Putative cation channel, NCA-1, and a novel protein, UNC-80, transmit neuronal activity in *C. elegans*. *PloS Biol.* 6: 552-567.

Table 1

		Case-control				Clozapine response				Weight gain				Tardive dyskinesia			
		Control		Case		Non-responder		Responder		<7%		≥ 7%		Absent		Present	
		n	%	n	%	n	%	n	%	n	%	n	%	n	%	n	%
1289556	A	270	61.6	257	60.9	73	50.0	77	57.5	44	57.9	27	46.6	145	56.2	100	61.7
	G	168	38.4	165	39.1	73	50.0	57	42.5	32	42.1	31	53.4	113	43.8	62	38.3
	AA	81	37.0	81	38.4	18	24.7	27	40.3	14	36.8	08	27.6	43	33.3	33	40.7
	AG	108	49.3	95	45.0	37	50.7	23	34.3	16	42.1	11	37.9	59	45.7	34	42.0
	GG	30	13.7	35	16.6	18	24.7	17	25.4	08	21.1	10	34.5	27	20.9	14	17.3
9554752	A	161	36.8	155	36.7	46	31.5	51	38.1	26	34.2	19	32.8	92	35.7	59	36.4
	G	277	63.2	267	63.3	100	68.5	83	61.9	50	65.8	39	67.2	166	63.3	103	63.6
	AA	33	15.1	28	13.3	06	08.2	12	17.9	04	10.5	04	13.8	19	14.7	12	14.8
	AG	95	43.4	99	46.9	34	46.6	27	40.3	18	47.4	11	37.9	54	41.9	35	43.2
	GG	91	41.6	84	39.8	33	45.2	28	41.8	16	42.1	14	48.3	56	43.4	34	42.0
1767752	A	140	32.1	147	34.8	41	28.1	47	35.1	21	27.6	18	31.0	85	32.9	59	34.0
	G	296	67.9	275	65.2	105	71.9	87	64.9	55	72.4	40	69.0	173	67.1	103	66.0
	AA	24	11.0	25	11.8	05	06.8	10	14.9	02	05.3	03	10.3	18	14.0	10	12.3
	AG	92	42.0	97	46.0	31	42.5	27	40.3	17	44.7	12	41.4	49	38.0	35	43.2
	GG	102	46.6	89	42.2	37	50.7	30	44.8	19	50.0	14	48.3	62	48.1	36	44.4
686141	A	140	32.1	127	30.1	47	32.2	30	22.4	25	32.9	13	22.4	72	27.9	49	30.2
	G	296	67.9	295	69.9	99	67.8	104	77.6	51	67.1	45	77.6	186	72.2	113	69.8
	AA	15	06.8	15	07.1	07	09.6	04	06.0	04	10.5	02	06.9	12	09.3	05	06.2
	AG	91	41.6	81	38.4	33	45.2	22	32.8	17	44.7	09	31.0	48	37.2	39	48.1
	GG	113	51.6	115	54.5	33	45.2	41	61.2	17	44.7	18	62.1	69	53.5	37	45.7
12867417	A	91	20.8	78	18.5	37	25.3	26	19.4	22	28.9	10	17.2	58	22.5	46	28.4
	T	347	79.2	344	81.5	109	74.7	108	80.6	54	71.1	48	82.8	200	77.5	116	71.6
	AA	12	05.5	06	02.8	04	05.5	03	04.5	03	07.9	02	06.9	09	07.0	04	04.9
	AT	67	30.6	66	31.3	29	39.7	20	29.9	16	42.1	06	20.7	40	31.0	38	46.9
	TT	140	63.9	139	65.9	40	54.8	44	65.7	19	20.7	21	72.4	80	62.0	39	48.1
658213	A	119	27.3	103	24.5	30	20.5	32	23.9	14	18.4	13	22.4	55	21.3	35	21.6
	C	317	72.7	317	75.5	116	79.5	102	76.1	62	81.6	45	77.6	203	78.7	127	78.4
	AA	14	06.4	14	06.6	03	04.1	05	07.5	00	0.00	01	03.4	09	07.0	04	04.9
	AC	91	41.6	75	35.5	24	32.9	22	32.8	14	36.8	11	37.9	37	28.7	27	33.3
	CC	113	51.6	121	57.3	46	63.0	40	59.7	24	63.2	17	58.6	83	64.3	50	61.7
614728	A	140	32.1	127	30.1	115	78.8	106	79.1	57	75.0	50	86.2	203	21.3	120	25.9
	C	296	67.9	295	69.9	31	21.2	28	20.9	19	25.0	08	13.8	55	78.7	42	74.1
	AA	22	10.0	21	10.0	45	61.6	43	64.2	21	55.3	22	75.9	83	64.3	42	51.9
	AC	96	43.8	85	40.3	25	34.2	20	29.9	15	39.5	06	20.7	37	28.7	36	44.4
	CC	100	45.7	105	49.8	03	04.1	04	06.0	02	05.3	01	03.4	09	07.0	03	03.7
9513851	A	25	05.7	32	07.6	10	06.8	07	05.2	05	06.6	05	08.6	19	07.4	05	02.5
	C	411	94.3	390	92.4	63	93.2	127	94.8	71	93.4	53	91.4	239	92.6	158	97.5
	AA	00	00.0	00	00.0	00	00.0	01	01.5	00	00.0	00	00.0	00	00.0	01	01.2
	AC	25	11.4	32	15.2	10	13.7	05	07.5	05	13.2	05	17.2	19	14.7	02	02.5
	CC	193	88.1	179	84.8	63	86.3	61	91.0	33	86.8	24	82.8	110	85.3	78	96.3
9518307	A	409	93.4	389	92.2	135	92.5	127	94.8	71	93.4	52	89.7	236	92.2	157	96.9
	C	29	6.6	33	7.8	11	07.5	07	05.2	05	06.6	06	10.3	20	07.8	05	03.1
	AA	190	86.8	179	84.8	63	86.3	61	91.0	33	86.8	24	82.8	110	85.3	77	95.1
	AC	29	13.2	31	14.7	09	12.3	05	07.5	05	13.2	04	13.8	18	14.0	03	03.7
	CC	00	00.0	01	00.5	01	01.4	01	01.5	00	00.0	01	03.4	01	00.8	01	01.2

		Case-control				Clozapine response				Weight gain				Tardive dyskinesia			
		Control		Case		Non-responder		Responder		<7%		≥ 7%		Absent		Present	
		n	%	n	%	n	%	n	%	n	%	n	%	n	%	n	%
12584031	A	358	81.7	338	80.1	119	81.5	111	82.8	64	84.2	47	81.0	216	83.7	135	83.3
	G	80	18.3	84	19.9	27	18.5	23	17.2	12	15.8	11	19.0	42	16.3	27	16.7
	AA	146	66.7	133	63.0	47	64.4	48	71.6	27	71.1	20	69.0	92	71.3	56	69.1
	AG	66	30.1	72	34.1	25	34.2	15	22.4	10	26.3	07	24.1	32	24.8	23	28.4
	GG	07	03.2	06	02.8	01	01.4	04	06.0	01	02.6	02	06.9	05	03.9	02	02.5
1452112	A	102	23.3	112	26.5	38	26.0	45	33.6	23	30.3	19	32.8	69	26.7	49	30.2
	G	336	76.7	310	73.5	108	74.0	89	66.4	53	69.7	39	67.2	189	73.3	113	69.8
	AA	07	03.2	13	06.2	04	05.5	06	09.0	03	07.9	02	06.9	07	05.4	06	07.4
	AG	88	40.2	86	40.8	30	41.1	33	49.3	17	44.7	15	51.7	55	42.6	37	45.7
	GG	124	56.6	112	53.1	39	53.4	28	41.8	18	47.4	12	41.4	67	51.9	38	46.9
7317836	A	130	29.7	127	30.1	43	29.5	36	26.9	17	22.4	12	20.7	72	27.9	39	24.1
	G	308	70.3	295	69.9	103	70.5	98	73.1	59	77.6	46	79.3	186	72.1	123	75.9
	AA	18	08.2	19	09.0	04	05.5	05	07.5	00	00.0	00	00.0	09	07.0	06	07.4
	AG	94	42.9	89	42.2	35	47.9	26	38.8	17	44.7	12	41.4	54	41.9	27	33.3
	GG	107	48.9	103	48.8	34	46.6	36	53.7	21	55.3	17	58.6	66	51.2	48	59.3
9518320	A	288	65.8	314	74.4	101	69.2	95	70.9	53	69.7	43	74.1	176	68.2	117	72.2
	G	150	34.2	108	25.6	45	30.8	39	29.1	23	30.3	15	25.9	82	31.8	45	27.8
	AA	96	43.8	120	56.9	38	52.1	35	52.2	20	52.6	17	58.6	62	48.1	42	51.9
	AG	96	43.8	74	35.1	25	34.2	25	37.3	13	34.2	09	31.0	52	40.3	33	40.7
	GG	27	12.3	17	08.1	10	13.7	07	10.4	05	13.2	03	10.3	15	11.6	06	07.4
9518331	A	150	34.2	108	25.7	46	31.5	39	29.1	23	30.3	18	31.0	88	34.1	44	37.2
	T	288	65.8	312	74.3	100	68.5	95	70.9	53	69.7	40	69.0	170	65.9	118	72.8
	AA	28	12.8	15	07.1	09	12.3	06	09.0	04	10.5	03	10.3	17	13.2	05	06.2
	AT	94	42.9	78	37.0	28	38.4	27	40.3	15	39.5	12	41.4	54	41.9	34	42.0
	TT	97	44.3	117	55.5	36	49.3	34	50.7	19	50.0	14	48.3	58	45.0	42	51.9
2584531	A	267	61.0	281	66.6	83	56.8	83	61.9	44	57.9	35	60.3	155	60.1	104	64.2
	C	171	39.0	141	33.4	63	43.2	51	38.1	32	42.1	23	39.7	103	39.9	58	33.8
	AA	82	37.4	97	46.0	25	34.2	29	43.3	15	39.5	12	41.4	50	38.8	37	45.7
	AC	103	47.0	87	41.2	33	45.2	25	37.3	14	36.8	11	37.9	55	42.6	30	37.0
	CC	34	15.5	27	12.8	15	20.5	13	19.4	09	23.7	06	20.7	24	18.6	14	17.3
12869164	A	00	00.0	00	00.0	00	00.0	00	00.0	00	00.0	00	00.0	00	00.0	00	00.0
	C	436	100	422	100	146	100	134	100	76	100	58	100	258	100	162	100
	AA	00	00.0	00	00.0	00	00.0	00	00.0	00	00.0	00	00.0	00	00.0	00	00.0
	AC	00	00.0	00	00.0	00	00.0	00	00.0	00	00.0	00	00.0	00	00.0	00	00.0
	CC	218	100	211	100	73	100	67	100	38	100	29	100	129	100.0	81	100.0
3916906	A	158	36.1	126	29.9	52	35.6	45	33.6	25	32.9	20	34.5	94	36.4	46	28.4
	C	280	63.9	296	70.1	94	64.4	89	66.4	51	67.1	38	65.5	167	63.6	116	71.6
	AA	29	13.2	22	10.4	12	16.4	09	13.4	06	15.8	04	13.8	20	15.5	07	08.6
	AC	100	45.7	82	38.9	28	38.4	27	40.3	13	34.2	12	41.4	54	41.9	32	39.5
	CC	90	41.1	107	50.7	33	45.2	31	46.3	19	50.0	13	44.8	55	42.6	42	51.9
9518349	A	262	59.8	232	55.0	95	65.1	81	60.4	53	69.7	39	67.2	161	62.4	93	57.4
	C	176	40.2	190	45.0	51	34.9	53	39.6	23	30.3	19	32.8	97	37.6	69	42.6
	AA	79	36.1	69	32.7	31	42.5	26	38.8	18	47.4	13	44.8	47	36.4	31	38.3
	AC	104	47.5	94	44.5	33	45.2	29	43.3	17	44.7	13	44.8	67	51.9	31	38.3
	CC	36	16.4	48	22.7	09	12.3	12	17.9	03	07.9	03	10.3	15	11.6	19	23.5

		Case-control				Clozapine response				Weight gain				Tardive dyskinesia			
		Control		Case		Non-responder		Responder		<7%		≥7%		Absent		Present	
		n	%	n	%	n	%	n	%	n	%	n	%	n	%	n	%
10508059	A	83	18.9	82	19.4	27	18.8	22	16.4	15	19.7	11	19.0	38	14.7	34	21.0
	G	355	81.1	340	85.0	119	81.5	112	83.6	61	80.3	47	81.0	220	85.3	128	79.0
	AA	07	03.2	08	03.8	03	04.1	04	06.0	04	10.5	01	03.4	07	05.4	03	03.7
	AG	69	31.5	66	31.3	21	28.8	14	20.9	07	18.4	09	31.0	24	18.6	28	34.6
	GG	143	65.3	137	64.9	49	67.1	49	73.1	27	71.1	19	65.5	98	76.0	50	61.7
7328287	A	180	41.1	170	40.5	65	44.5	60	44.8	35	46.1	28	48.3	120	46.5	58	35.8
	G	258	58.9	250	59.5	81	55.5	74	55.2	41	53.9	30	51.7	138	53.5	104	64.2
	AA	39	17.8	31	14.7	14	19.2	16	23.9	09	23.7	06	23.7	29	22.5	09	11.1
	AG	102	46.6	108	51.2	37	50.7	28	41.8	17	44.7	16	44.7	62	48.1	40	49.4
	GG	78	35.6	71	33.6	22	30.1	23	34.3	12	31.6	07	31.6	38	29.5	32	39.5
496238	A	153	34.9	141	33.4	54	37.0	52	38.8	29	38.2	26	44.8	102	39.5	53	32.7
	G	285	65.1	281	66.6	92	63.0	82	61.2	47	61.8	32	55.2	156	60.5	109	67.3
	AA	29	13.2	26	12.3	09	12.3	14	20.9	07	18.4	06	20.7	21	16.3	08	09.9
	AG	95	43.4	89	42.2	36	49.3	24	35.8	15	39.5	14	48.3	60	46.5	37	45.7
	GG	95	43.4	96	45.5	28	38.4	29	43.3	16	48.3	09	31.0	48	37.2	36	44.4
7318529	A	37	8.4	40	9.5	16	11.0	13	09.7	09	11.8	07	12.1	25	09.7	09	05.6
	G	401	91.6	382	90.5	130	89.0	121	90.3	67	88.2	51	87.9	233	90.3	153	94.4
	AA	05	02.3	03	01.4	01	01.4	01	01.5	01	02.6	01	03.4	02	01.6	01	01.2
	AG	27	12.3	34	16.1	14	19.2	11	16.4	07	18.4	05	17.2	21	16.3	07	08.6
	GG	187	85.4	174	82.5	58	79.5	55	82.1	30	78.9	23	79.3	106	82.2	73	90.1
9554772	A	189	43.2	188	44.8	65	44.5	62	46.3	36	47.7	24	41.4	109	22.6	63	38.9
	T	249	56.8	232	55.2	81	55.5	72	53.7	40	52.6	34	58.6	147	57.4	99	61.1
	AA	40	18.3	41	19.4	12	16.4	17	25.4	09	23.7	06	20.7	21	16.4	15	18.5
	AT	109	49.8	106	50.2	41	56.2	28	41.8	18	47.4	12	41.4	67	52.3	33	40.7
	TT	70	32.0	63	29.9	20	27.4	22	32.8	11	28.9	11	37.9	40	31.3	33	40.7
17486808	A	93	21.2	95	22.5	30	20.5	34	25.4	17	22.4	11	19.0	59	22.9	31	19.1
	C	345	78.8	327	77.5	116	79.5	100	74.6	76	77.6	47	81.0	199	77.1	131	80.9
	AA	11	05.0	14	06.6	04	05.5	05	07.5	02	05.3	02	06.9	05	03.9	05	06.2
	AC	71	32.4	67	31.8	22	30.1	24	35.8	13	34.2	07	24.1	49	38.0	21	25.9
	CC	137	62.6	130	61.6	47	64.4	38	56.7	23	60.5	20	69.0	75	58.1	55	67.9
17584161	A	109	24.9	113	26.8	32	21.9	39	29.1	19	25.0	13	22.4	69	26.7	42	25.9
	C	329	75.1	309	73.2	114	78.1	95	70.9	57	75.0	45	77.6	189	73.3	120	74.1
	AA	17	07.8	20	09.5	07	09.6	07	10.4	05	13.2	03	10.3	09	07.0	06	07.4
	AC	75	34.2	73	34.6	18	24.7	25	37.3	09	23.7	07	24.1	51	39.5	30	37.0
	CC	127	58.0	118	55.9	48	65.8	35	52.2	24	63.2	19	65.5	69	53.5	45	55.6
2152324	A	66	15.1	65	15.4	24	16.4	12	09.0	07	09.2	10	17.2	24	09.3	16	09.9
	C	372	84.9	357	84.6	122	83.6	122	91.0	69	90.8	48	82.8	234	90.7	146	90.1
	AA	05	02.3	03	01.4	00	00.0	01	01.8	00	00.0	01	03.4	02	01.6	01	01.2
	AC	56	25.6	59	28.0	24	32.9	10	14.9	07	18.4	08	27.6	20	15.5	14	17.3
	CC	158	72.1	149	70.6	49	67.1	56	83.6	31	81.6	20	69.0	107	82.9	66	81.5

Table 2

SNP	Allele	Frequency	Family	S	E(S)	Var(S)	Z	P
rs1289556	A	0.644	36	40	42.150	14.344	-0.568	0.570
rs9554752	A	0.413	36	30	31.233	10.434	-0.382	0.702
rs17677552	A	0.384	37	30	31.067	11.462	-0.315	0.752
rs614728	A	0.287	30	20	20.667	09.722	-0.214	0.830
rs686141	A	0.277	28	21	20.417	09.160	0.193	0.847
rs12869164	A	0.220	27	21	18.267	08.962	0.913	0.361
rs658213	A	0.744	29	41	40.083	09.410	0.299	0.765
rs9513851	A	0.087	10	05	04.833	02.472	0.106	0.915
rs9518307	A	0.917	10	15	15.167	02.472	-0.106	0.915
rs12584031	A	0.791	26	42	39.217	07.320	1.029	0.303
rs1452112	A	0.289	28	25	19.517	08.650	1.864	0.062
rs7317836	A	0.320	29	22	22.967	09.132	-0.320	0.749
rs9518320	A	0.680	29	41	41.417	09.024	-0.139	0.889
rs9518331	A	0.324	28	20	19.250	08.688	0.254	0.799
rs2584531	A	0.637	29	42	41.583	09.160	0.138	0.890
rs12867417	A	0.000	-	-	-	-	-	-
rs3916906	A	0.348	30	19	19.917	09.410	-0.299	0.765
rs9518349	A	0.517	33	28	32.250	10.465	-1.314	0.188
rs10508059	A	0.195	24	13	17.500	08.528	-1.541	0.123
rs7328287	A	0.435	33	36	31.667	12.614	1.220	0.222
rs496238	A	0.377	32	30	27.500	12.586	0.705	0.481
rs7318529	A	0.088	14	10	07.667	03.722	1.209	0.226
rs9554772	A	0.421	32	34	33.167	12.364	0.237	0.812
rs17486808	A	0.230	24	19	17.250	08.188	0.612	0.540
rs17584161	A	0.269	27	18	19.417	08.660	-0.481	0.630
rs2152324	A	0.179	23	17	15.917	06.660	0.420	0.674

Figure 1





Figure S1



Legends

Table 1: Allelic and genotypic frequencies

Table 2: Family-based association test results. S represents the test statistic for observed number of alleles, E represents the expected value of S under null hypothesis and $\text{Var}(S)$ represents variance between the observed and expected transmission.

Figure 1: P-values for Hardy-Weinberg equilibrium p-values (A) and association results (B). The solid line represents $p = 0.05$

Figure S1: LD plot for the analyzed markers in case-control, family, response, weight gain and TD samples (A, B, C, D and E); respectively. Values presented are the D' .

3.5 - Association of antipsychotic induced weight gain and body mass index with GNB3 gene: a meta-analysis

(Progress in Neuro-psychopharmacology & Biological Psychiatry; aceito em 22/08/08)

Association of antipsychotic induced weight gain and body mass index with GNB3 gene:
a meta-analysis

Renan P. Souza^{1,2}, Vincenzo De Luca^{1,3*}, Giovanni Muscettola³, Daniela VF Rosa²,
Andrea de Bartolomeis³, Marco Romano Silva² and James L. Kennedy¹

1 Neurogenetics Section, Centre for Addiction and Mental Health, University of Toronto, Canada
2 Departamento de Saude Mental, Universidade Federal de Minas Gerais, Brazil
3 Department of Neuroscience, Section of Psychiatry, University of Naples 'Federico II'

Abstract

It has been reported that C825T variant in the gene encoding the G-protein subunit $\beta 3$ (GNB3) is associated with antipsychotic-induced weight gain and obesity. We investigated the association of the *GNB3* and antipsychotic-induced weight gain as well as body mass index (BMI) using meta-analytical techniques. Our analysis of 402 schizophrenia subjects showed a trend ($p = 0.072$) only under a fixed-model. As it was observed heterogeneity among the studies ($p = 0.007$), we re-analyzed using a random-effects framework and no significance was found ($p = 0.339$). No evidence for bias publication was reported ($p = 0.868$). Our analysis of 18,903 subjects showed a trend ($p = 0.053$) associating CC and lower BMI under a fixed model. Although no significant association was found, the same pattern (CC and lower antipsychotic-induced weight gain) was observed. Our meta-analysis indicates that firmly establishing the role of pharmacogenetics in clinical psychiatry requires much larger sample sizes that have been reported.

Key words: obesity; BMI; antipsychotic-induced weight gain; GNB-3; meta-analysis

Introduction

Antipsychotic medications are an important component in the medical management of many psychotic conditions. Although they have many notable benefits compared with their earlier counterparts, their use has been associated with reports of dramatic weight gain, diabetes (even acute metabolic decompensation, e.g., diabetic ketoacidosis), and an atherogenic lipid profile (increased LDL cholesterol and triglyceride levels and decreased HDL cholesterol). Because of the close associations between obesity, diabetes, and dyslipidemia and cardiovascular disease, there is heightened interest in the relationship between the antipsychotic drugs and the development of these major cardiovascular disease risk factors (Henderson, 2005).

Search for predictors of drug-related morbidity is becoming increasingly important in persons with major mental illness. Weight gain, glucose and lipid abnormalities are observed more frequently in some novel antipsychotics (Newcomer and Haupt, 2006). The relevance of this side-effect clearly arises from the following considerations: (1) a significant increase in weight gain may affect the compliance to pharmacotherapy and be indirectly responsible for psychosis relapses; (2) weight gain may add to schizophrenia stigma the stigma of obesity and this in turn may lead to poor adherence to the therapy; (3) weight gain may increase the risk for diabetes type II; (4) weight gain can be associated with the metabolic syndrome (Haddad, 2004).

Twenty-nine percent of persons with schizophrenia gain at least 7% of their baseline body weight after being treated with olanzapine in short-term studies (Leucht *et al.* 1999). The FDA has established that weight gain $\geq 7\%$ from baseline constitutes a clinically meaningful and significant metabolic outcome. Clinical significance of prolonged therapy is now becoming realized as patients develop Type II diabetes mellitus and other obesity-related problems as a consequence of antipsychotic use (American Diabetes Association *et al.* 2004). When looking for possible ethnic differences, clinical studies suggest that Afro-American patients treated with antipsychotics are at higher risk of weight gain (Blin and Micallef, 2001).

G proteins relay signals from each of more than 1000 receptors to many different effectors, including enzymes and ion channels. G proteins are composed of α -subunit that is loosely bound to a tightly associated structure made up of a β subunit and a γ subunit. The activity of the trimeric G protein is regulated by the binding and hydrolysis of guanosine triphosphate by the $G\alpha$ subunit. The α -subunit to which guanosine diphosphate is bound is inactive and associates with the $\beta\gamma$ dimer (Neves *et al.* 2002). In 1998, Siffert *et al.* described a C825T polymorphism of the *GNB3* gene. C825T polymorphism is located 1700 bp upstream of the alternative splice site, indicating that affect of *GNB3* 825T on the splice process is a complex mechanism. Nevertheless, there are examples that single distant nucleotide exchanges, not related to conserved splice branch, donor, and acceptor sites, can cause such alternative splicing (Stallings-Mann *et al.* 1996; Liu *et al.* 1997).

Recent advances in research on the genetic contributions to obesity and its related phenotypes are providing novel tools and targets for the study of mechanisms and risk factors for antipsychotic-induced weight gain. Candidate gene selection should rely on current knowledge on the molecular pathways to weight gain, antipsychotic pharmacokinetics and pharmacodynamics, as well as possible disease-related genetic

links to the side effects under study (Correll and Malhotra, 2004; Muller *et al.* 2004) 825T allele is associated with G β 3 splice variant, which, despite a deletion of 41 amino acids, is functionally active in reconstituted systems. (Siffert *et al.* 1998). There are few reports linking C825T polymorphism and antipsychotic induced weight gain, that showed no association with either clozapine or olanzapine induced weight gain (Tsai *et al.* 2004; Bishop *et al.* 2006). This variant was chosen as it was previously described associated with obesity in several ethnic groups and weight gain during pregnancy (Bishop *et al.* 2006).

Currently available data suggest a worldwide continuous increase in obesity prevalence, which is recently also being observed in developing countries. This prompts some authors to predict an “obesity epidemic” with an increased prevalence of hypertension, stroke, coronary artery disease, and type 2 diabetes mellitus, for which obesity is a major risk factor. Several studies also have investigated whether the 825T allele increases the risk for obesity (Hegele *et al.* 1999; Siffert *et al.* 1999a; Siffert *et al.* 1999b, Stefan *et al.* 2004), these studies have demonstrated that this allele is associated with obesity or increased BMI. Other similar studies, however, have failed to demonstrate this association (Benjafield *et al.* 2001; Hinney *et al.* 2001; Ohshiro *et al.* 2001; Snapir *et al.* 2001; Poston *et al.* 2002; Suwazono *et al.* 2004; Hayakawa *et al.* 2007). Furthermore, it has been evaluated prediction of successful weight reduction under sibutramine therapy and C825T polymorphism but no association was found with non-pharmacologic weight loss strategies (Hauner *et al.* 2003; Potoczna *et al.* 2004). Because of these varying findings, we considered that the apparent association between GNB3 gene variant and obesity or raised BMI had not been demonstrated conclusively. From an epidemiological point of view, we believe that in order to determine the influence of genetic polymorphisms in the occurrence of a specific disease, it is necessary to undertake large-scale studies in the general population. To elucidate better this question we performed this meta-analysis reaching a more significant number of antipsychotic-induced weight gain and BMI related to GNB3 C825T variant.

Methods

Inclusion criteria

Genetic association studies examining the association between C825T and antipsychotic-induced weight gain among patients with schizophrenia that compared the homozygote genotypes were included. Furthermore, in order to analyze the variant influence in the BMI, genetic association studies examining the association between C825T and BMI among adults that compared the homozygote genotypes were also included.

Search strategy

We searched the National Library of Medicine’s PubMed online using the search strategy: ‘weight gain’, ‘GNB3’, ‘obesity’ and BMI. This database was searched up to December 2007 looking for terms.

Data extraction

For each study, the following data were extracted using standard forms: author, year of publication, sample ethnicity, case and control sample size, allele frequency, mean age, sex ratio. Ethnicity was coded as European, Asian and African-American.

Statistics

Standardized mean differences (SMDs) and their standard error (S.E.) for individual studies were calculated from 2x2 tables in a case control format. Pooled SMDs were calculated using fixed-effects and random-effects approaches (Der-Simonian and Laird, 1986), and the significance of the pooled SMDs determined using a Z test. The assumption that the effect of allele frequency is constant across studies and between-studies variation is due to random variation was checked using a χ^2 test for heterogeneity of SMDs. In absence of significant heterogeneity, data were initially analyzed within a fixed-effects framework; otherwise a random-effects framework was employed using Der-Simonian and Laird methods (Der-Simonian and Laird, 1986). This assumes that between-study variation is due to both random variation and an individual study effect. Random-effects models are more conservative and generate a wider confidence interval. Publication bias was assessed by means of a funnel plot of individual study log SMD against S.E. log SMD, and formally by the method of Egger (Egger et al. 1997), which is based on a weighted linear regression of standard normal deviation of the SMD (standardized effect) on the inverse of the standard error of the SMD (precision). Data were analyzed using the STATA version 8.0 statistical software package (Stata Corporation, College Station, TX, USA).

Results

Antipsychotic-induced weight gain

A total of five studies published between 2004 and 2007, comprising five independent samples, were identified by the search strategy, met the inclusion criteria and contributed to the meta-analysis. Two studies (De Luca et al. 2004; Souza et al. 2007) included in the meta-analysis are from our group and they were published in the conferences abstract book. Each sample was included independently in the analysis. The main outcome investigated in this meta-analysis was the weight percentage change in kilograms after the treatment. To dissect the genotype effect only the CC and TT genotypes were included in the analysis. The data entered in the meta-analysis table were mean, standard deviation and number of subjects for CC and TT genotype respectively. Two were performed with Caucasian subjects, two with Asian (Chinese) and one with mixed population (Caucasian and African-American in Souza *et al.* 2007) (Table 1). When all samples were included there was a trend for CC association with lower antipsychotic-induced weight gain under a fixed model ($z=1.80$, $p=0.072$, SMD 0.27, 95% CI $-0.025 - 0.584$); however, there was a significant heterogeneity between the studies ($X^2= 14.25$, $d.f.=4$, $p=0.007$). As it was found heterogeneity in this sample, this data set has been analyzed using random effects model as it is the most appropriate model. When the analysis was re-run within a random-effects framework no significance was found ($z=0.96$, $p=0.339$, SMD 0.28, 95% CI $-0.03 - 0.58$) (Figure 1A). As there are strong evidences that ethnic background is an important confounding factor in antipsychotic-

induced weight gain genetics, stratified analysis by ethnicity were performed. Asians and Caucasians subpopulations (as there is just one study that analyzed African-American population) were created and their results did not show significant associations (data not shown). A Begg's funnel plot with 95% confidence limits is presented in Figure 1B. Egger's test did not report evidence of publication bias (intercept= 0.784, $t=0.18$, $p=0.868$, 95% CI -12.973 – 14.542) (Figure 1C).

BMI association

A total of 39 studies published between 1999 and 2007, comprising 34 independent samples, were identified by the search strategy, met the inclusion criteria and contributed to the meta-analysis. Five studies that reported the BMI value for one of the homozygous genotypes grouped with the heterozygous were excluded. 16 studies did not report BMI for each genotype and another one that reported it for each allele were also excluded. From the 18 studies that reported BMI for CC and TT groups, 12 were performed with Caucasian subjects, three with Asian (Chinese or Japanese), one with African-American and another two with mixed populations (Caucasian and Asian in Siffert *et al.* 1999b; Caucasian and African-American in Danoviz *et al.* 2006) (Table 2). When all samples were included there was a trend for association of CC and lower BMI under a fixed model ($z=1.93$, $p=0.053$, SMD 0.05, 95% CI 0.00 - 0.09), and there was also a trend of significant heterogeneity between the studies ($X^2= 31.70$, $d.f.=21$, $p=0.063$) (Figure 2A). A Begg's funnel plot with 95% confidence limits is presented in Figure 2B. This illustrates a certain asymmetry with predominance of small and positive studies over small and negative studies. Egger's test did not indicate evidence of publication bias (intercept= 0.897, $t=1.52$, $p=0.143$, 95% CI -0.331 – 2.125), although it is possible to notice that small samples with low precision having a large standardized effect and large samples with high precision having small-standardized effect (Figure 2C).

Discussion

Weight gain is probably the most actual side effect in antipsychotic treatment due to the wide use of new antipsychotics, on the other hand the pharmacogenetic of antipsychotics has focused more on side effects like tardive dyskinesia (Lerer *et al.* 2005), and therefore we were able to find only a few studies exploring weight gain. Our analysis of schizophrenia subjects showed a trend ($p = 0.072$) only under a fixed-model probably because we included only the homozygous CC and TT in the analysis however the overall sample was composed by 406 subjects. As it was observed heterogeneity among the studies ($p = 0.007$), we re-analyzed using a random-effects framework and no significance was found ($p = 0.339$). No evidence for bias publication was reported ($p = 0.868$). However, there are few published studies in the area and a much larger sample size would be needed to test this hypothesis definitively.

A considerable part of obesity is due to environmental factors and lifestyle, but between 40-70% of the variation of body mass index (BMI) is estimated to be heritable (Comuzzie *et al.* 1988). Our analysis of 18,903 subjects showed a trend ($p = 0.053$) associating CC and lower BMI under a fixed model. Even in the BMI analysis we selected only the CC and TT subjects, this can be an advantage since we reduced the

genetic heterogeneity, however this reduces the statistical power of the sample. There may be a trend towards an association but the size of the effect is small and much larger studies are needed to demonstrate this association. Although no significant association was found, the same pattern (CC and lower antipsychotic-induced weight gain) was observed.

The 825T allele has already been analyzed in various contexts that could increase risk for phenotypes metabolic syndrome-related. Total cholesterol is significantly higher in subjects with the T allele among Japanese (Ishikawa *et al.* 2000); and this same allele was associated with end-stage renal disease in type 2 diabetes mellitus (Gumprecht *et al.* 2001). The C825T role in hyperlipidemia, diabetes, and diabetic complications have been controversial (Fogarty *et al.* 1998; Siffert *et al.* 1999; Rosskopf *et al.* 2000; Beige *et al.* 2000; Zychma *et al.* 2000; Gumprecht *et al.* 2001; Shcherbak *et al.* 2001; Hanon *et al.* 2002; Dzida *et al.* 2002; Von Beckerath *et al.* 2003; Brand *et al.* 2003; Yamamoto *et al.* 2004; Andersen *et al.* 2006; Hayakawa *et al.* 2007). At the same way, no conclusive data have been published in studies that evaluated genetic association of C825T with antipsychotic-induced weight gain (De Luca *et al.* 2004; Tsai *et al.* 2004; Wang *et al.* 2005; Bishop *et al.* 2006; Souza *et al.* 2007).

Metabolic imbalance has been found more common in drug-naïve schizophrenics rather than general population (Ryan and Thakore, 2002) thus this strategy to combine studies that have focussed on obesity and the side-effect of schizophrenia treatment analyzing common genetic targets seems very intriguing because it can help to uncover the common genetic background. This meta-analysis pointed out that the number of pharmacogenetic studies of antipsychotic-induced weight gain is very small and sometimes the sample size is not adequate. Furthermore, for some studies, position and dispersion measures are not always specified for both genotype groups and it is very important to specify whether SD or SEM has been described for including the study in future meta-analysis.

On the other hand the phenotype BMI that is closely related to weight gain has been widely investigated in regards of the C825T. This discrepancy is due to the fact that baseline BMI can be measured as a cross-sectional assessment instead weight change requires follow-up with the risk to lose subjects. Therefore, we suggest some points for the pharmacogenetics studies in order to enhance the power of antipsychotic-induced weight gain meta-analysis studies: to select phenotype such as the presence of metabolic syndrome that is measurable in a cross-sectional assessment; use of larger sample size; controlling for ethnicity and previous antipsychotic exposure.

Conclusion

In this paper we suggest a new way how to apply the meta-analytic technique to genetic association studies for dissecting the genetic influence in related phenotypes to show possible bias in the published studies and suggesting different methodological approaches to improve the overall quality of pharmacogenetic studies of antipsychotic-induced weight gain.

Bibliography

American Diabetes Association, American Psychiatric Association, American Association of Clinical Endocrinologists, North American Association for the Study of Obesity. Consensus development conference on antipsychotic drugs and obesity and diabetes. *J Clin Psychiatry* 2004; 65: 267–72

Andersen G, Overgaard J, Albrechtsen A, Glumer C, Borch-Johnsen K, Jorgensen T, et al . Studies of the association of the GNB3 825C>T polymorphism with components of the metabolic syndrome in white Danes. *Diabetologia* 2006; 49: 75-82.

Beige J, Ringel J, Distler A, Sharma AM . G-protein beta(3)-subunit C825T genotype and nephropathy in diabetes mellitus. *Nephrol Dial Transplant* 2000; 15: 1384-7.

Benjafield AV, Lin RC, Dalziel B, Gosby AK, Caterson ID, Morris BJ . G-protein beta3 subunit gene splice variant in obesity and overweight. *Int J Obes Relat Metab Disord* 2001; 25: 777-80.

Bishop JR, Ellingrod VL, Moline J, Miller D . Pilot study of the G-protein beta3 subunit gene (C825T) polymorphism and clinical response to olanzapine or olanzapine-related weight gain in persons with schizophrenia. *Med Sci Monit* 2006; 12: BR47-BR50.

Blin O, Micallef J . Antipsychotic-associated weight gain and clinical outcome parameters. *J Clin Psychiatry* 2001; 62: 11-21.

Brand E, Wang JG, Herrmann SM, Staessen JA . An epidemiological study of blood pressure and metabolic phenotypes in relation to the G beta3 C825T polymorphism. *J Hypertens* 2003; 21: 729-37.

Correll CU, Malhotra AK. Pharmacogenetics of antipsychotic-induced weight gain. *Psychopharmacology (Berl)* 2004;174:477-89.

Danoviz ME, Pereira AC, Mill JG, Krieger JE . Hypertension, obesity and GNB 3 gene variants. *Clin Exp Pharmacol Physiol* 2006; 33: 248-52.

De Luca V; Hwang R, Mueller DJ, Shinkai T, Basile VS, Masellis M, Meltzer HY, Lieberman JA, Kennedy JL. Genetics of clozapine-induced weight gain: beta 3 adrenergic receptor and G-protein beta 3 subunit/protein kinase G transduction pathway polymorphisms. *World Psychiatry Association International Congress Florence 2004*

Der Simonian R, Laird N . Meta-analysis in clinical trials. *Control Clin Trials* 1986; 7: 177-88.

Dzida G, Golon-Siekierska P, Puzniak A, Sobstyl J, Bilan A, Mosiewicz J, et al . G-protein beta3 subunit gene C825T polymorphism is associated with arterial hypertension in Polish patients with type 2 diabetes mellitus. *Med Sci Monit* 2002; 8: CR597-CR602.

Egger M, Smith GD, Phillips AN . Meta-analysis: principles and procedures. *BMJ* 1997; 315: 1533-7.

Fogarty DG, Zychma MJ, Scott LJ, Warram JH, Krolewski AS . The C825T polymorphism in the human G-protein beta3 subunit gene is not associated with diabetic nephropathy in Type I diabetes mellitus. *Diabetologia* 1998; 41: 1304-8.

Gumprecht J, Zychma MJ, Grzeszczak W, Zukowska-Szczechowska E . Transmission of G-protein beta3 subunit C825T alleles to offspring affected with end-stage renal disease. *Am J Nephrol* 2001; 21: 368-72.

Haddad PM . Antipsychotics and diabetes: review of non-prospective data. *Br J Psychiatry Suppl* 2004; 47: S80-S86.

Hanon O, Luong V, Mourad JJ, Bortolotto LA, Safar M, Girerd X . Association between the G protein beta3 subunit 825t allele and radial artery hypertrophy. *J Vasc Res* 2002; 39: 497-503.

Hauer H, Meier M, Jockel KH, Frey UH, Siffert W . Prediction of successful weight reduction under sibutramine therapy through genotyping of the G-protein beta3 subunit gene (GNB3) C825T polymorphism. *Pharmacogenetics* 2003; 13: 453-9.

Hayakawa T, Takamura T, Abe T, Kaneko S . Association of the C825T polymorphism of the G-protein beta3 subunit gene with hypertension, obesity, hyperlipidemia, insulin resistance, diabetes, diabetic complications, and diabetic therapies among Japanese. *Metabolism* 2007; 56: 44-8.

Hegele RA, Anderson C, Young TK, Connelly PW . G-protein beta3 subunit gene splice variant and body fat distribution in Nunavut Inuit. *Genome Res* 1999; 9: 972-7.

Henderson DC . Schizophrenia and comorbid metabolic disorders. *J Clin Psychiatry* 2005; 66 Suppl 6: 11-20.

Hengstenberg C, Schunkert H, Mayer B, Doring A, Lowel H, Hense HW, et al . Association between a polymorphism in the G protein beta3 subunit gene (GNB3) with arterial hypertension but not with myocardial infarction. *Cardiovasc Res* 2001; 49: 820-7.

Hinney A, Geller F, Neupert T, Sommerlad C, Gerber G, Gorg T, et al . No evidence for involvement of alleles of the 825-C/T polymorphism of the G-protein subunit beta 3 in body weight regulation. *Exp Clin Endocrinol Diabetes* 2001; 109: 402-5.

Huang X, Ju Z, Song Y, Zhang H, Sun K, Lu H, et al . Lack of association between the G protein beta3 subunit gene and essential hypertension in Chinese: a case-control and a family-based study. *J Mol Med* 2003; 81: 729-35.

Ishikawa K, Imai Y, Katsuya T, Ohkubo T, Tsuji I, Nagai K, et al . Human G-protein beta3 subunit variant is associated with serum potassium and total cholesterol levels but not with blood pressure. *Am J Hypertens* 2000; 13: 140-5.

Lerer B, Segman RH, Tan EC, Basile VS, Cavallaro R, Aschauer HN, et al . Combined analysis of 635 patients confirms an age-related association of the serotonin 2A receptor gene with tardive dyskinesia and specificity for the non-orofacial subtype. *Int J Neuropsychopharmacol* 2005; 8: 411-25.

Leucht S, Pitschel-Walz G, Abraham D, Kissling W . Efficacy and extrapyramidal side-effects of the new antipsychotics olanzapine, quetiapine, risperidone, and sertindole compared to conventional antipsychotics and placebo. A meta-analysis of randomized controlled trials. *Schizophr Res* 1999; 35: 51-68.

Martin DN, Andreu EP, Ramirez LR, Garcia-Junco PS, Vallejo M, I, Santos RA, et al . G-protein beta-3 subunit gene C825 T polymorphism: influence on plasma sodium and potassium concentrations in essential hypertensive patients. *Life Sci* 2005; 77: 2879-86.

Müller DJ, Muglia P, Fortune T, Kennedy JL. Pharmacogenetics of antipsychotic-induced weight gain. *Pharmacol Res.* 2004; 49:309-29.

Neves SR, Ram PT, Iyengar R . G protein pathways. *Science* 2002; 296: 1636-9.

Newcomer JW,Haupt DW . The metabolic effects of antipsychotic medications. *Can J Psychiatry* 2006; 51: 480-91.

Ohshiro Y, Ueda K, Wakasaki H, Takasu N, Nanjo K . Analysis of 825C/T polymorphism of G proteinbeta3 subunit in obese/diabetic Japanese. *Biochem Biophys Res Commun* 2001; 286: 678-80.

Poston WS, Haddock CK, Spertus J, Catanese DM, Pavlik VN, Hyman DJ, et al . Physical activity does not mitigate G-protein-related genetic risk for obesity in individuals of African descent. *Eat Weight Disord* 2002; 7: 68-71.

Potoczna N, Wertli M, Steffen R, Ricklin T, Lentjes KU, Horber FF . G protein polymorphisms do not predict weight loss and improvement of hypertension in severely obese patients. *J Gastrointest Surg* 2004; 8: 862-8.

Roskopf D, Frey U, Eckhardt S, Schmidt S, Ritz E, Hofmann S, et al . Interaction of the G protein beta 3 subunit T825 allele and the IRS-1 Arg972 variant in type 2 diabetes. *Eur J Med Res* 2000; 5: 484-90.

Ryan MC,Thakore JH . Physical consequences of schizophrenia and its treatment: the metabolic syndrome. *Life Sci* 2002; 71: 239-57.

Sartori M, Semplicini A, Siffert W, Mormino P, Mazzer A, Pegoraro F, et al . G-protein beta3-subunit gene 825T allele and hypertension: a longitudinal study in young grade I hypertensives. *Hypertension* 2003; 42: 909-14.

Shcherbak NS,Schwartz EI . The C825T polymorphism in the G-protein beta3 subunit gene and diabetic complications in IDDM patients. *J Hum Genet* 2001; 46: 188-91.

Siffert W, Roskopf D, Siffert G, Busch S, Moritz A, Erbel R, et al . Association of a human G-protein beta3 subunit variant with hypertension. *Nat Genet* 1998; 18: 45-8.

Siffert W, Forster P, Jockel KH, Mvere DA, Brinkmann B, Naber C, et al . Worldwide ethnic distribution of the G protein beta3 subunit 825T allele and its association with obesity in Caucasian, Chinese, and Black African individuals. *J Am Soc Nephrol* 1999; 10: 1921-30.

Siffert W, Naber C, Walla M, Ritz E . G protein beta3 subunit 825T allele and its potential association with obesity in hypertensive individuals. *J Hypertens* 1999; 17: 1095-8.

Souza RP; De Luca V; Romano-Silva MA; Wong AHC; Volavka J; Lieberman JA; Kennedy JL. Analysis in G-protein-beta3 subunit gene (C825T) polymorphism as a candidate gene to antipsychotic-related weight gain. 1st Canadian Association for Neuroscience Meeting Toronto 2007.

Snapir A, Heinonen P, Tuomainen TP, Lakka TA, Kauhanen J, Salonen JT, et al . G-protein beta3 subunit C825T polymorphism: no association with risk for hypertension and obesity. *J Hypertens* 2001; 19: 2149-55.

Stefan N, Stumvoll M, Machicao F, Koch M, Haring HU, Fritsche A . C825T polymorphism of the G protein beta3 subunit is associated with obesity but not with insulin sensitivity. *Obes Res* 2004; 12: 679-683.

Suwazono Y, Okubo Y, Kobayashi E, Miura K, Morikawa Y, Ishizaki M, et al . Lack of association between human G-protein beta3 subunit variant and overweight in Japanese workers. *Obes Res* 2004; 12: 4-8.

Tsai SJ, Yu YW, Lin CH, Wang YC, Chen JY, Hong CJ . Association study of adrenergic beta3 receptor (Trp64Arg) and G-protein beta3 subunit gene (C825T) polymorphisms and weight change during clozapine treatment. *Neuropsychobiology* 2004; 50: 37-40.

von Beckerath N, Kastrati A, Koch W, Bottiger C, Mehilli J, Seyfarth M, et al . G protein beta3 subunit polymorphism and risk of thrombosis and restenosis following coronary stent placement. *Atherosclerosis* 2000; 149: 151-5.

Wang YC, Bai YM, Chen JY, Lin CC, Lai IC, Liou YJ . C825T polymorphism in the human G protein beta3 subunit gene is associated with long-term clozapine treatment-induced body weight change in the Chinese population. *Pharmacogenet Genomics* 2005; 15: 743-8.

Yamamoto M, Abe M, Jin JJ, Wu Z, Tabara Y, Mogi M, et al . Association of GNB3 gene with pulse pressure and clustering of risk factors for cardiovascular disease in Japanese. *Biochem Biophys Res Commun* 2004; 316: 744-8.

Zychma MJ, Zukowska-Szczehowska E, Ossowska-Szymkowicz I, Trautsolt W, Grzeszczak W . G-Protein beta(3) subunit C825T variant, nephropathy and hypertension in patients with type 2 (Non-insulin-dependent) diabetes mellitus. Am J Nephrol 2000; 20: 305-10.

Table 1

Study	Year	n	Ancestry	Subjects
De Luca <i>et al.</i>	2004	80	Caucasian	Schizophrenia patients weight-gain after 6 weeks taking clozapine
Tsai <i>et al.</i>	2004	87	Asian	Schizophrenia patients weight-gain after 4 months taking clozapine
Wang <i>et al.</i>	2005	134	Asian	Schizophrenia patients weight-gain after 13 months taking clozapine
Bishop <i>et al.</i>	2006	42	Caucasian	Schizophrenia patients weight-gain after 6 weeks taking olanzapine
Souza <i>et al.</i>	2007	59	Caucasian/ African-American	Schizophrenia patients weight-gain after 14 weeks taking mixed antipsychotics

Table 2

Study	Year	n	Ancestry	Subjects
Siffert	1999	277/960	Caucasian/Asian	Healthy male
Hegele	1999	213	Caucasian	Randomly selected individuals
Siffert	1999	197	Caucasian	Hypertensive individuals
Hengstenberg	2001	2052/606	Caucasian	Randomly selected/ Myocardial infarction
Snapiir	2001	903	Caucasian	Randomly selected males
Poston	2001	175	African-American	Randomly selected individuals
Hanon	2002	306	Caucasian	Atherosclerosis Prevention Clinic subjects
Brand	2003	1512	Caucasian	Randomly selected
Huang	2003	1165	Asian	Case-control hypertension study
Beckerath	2003	1338	Caucasian	Case-control coronary artery disease study
Sartori	2003	461	Caucasian	Hypertensive individuals
Stefan	2004	774	Caucasian	Case-control impaired glucose tolerant study
Potoczna	2004	304	Caucasian	Severely obese individuals
Yamamoto	2004	806	Asian	Subject who participated in a medical check-up
Martin	2005	144	Caucasian	Case-control hypertension study
Danoviz	2006	1568	Caucasian/African-American	Randomly selected individuals
Andersen	2006	4387	Caucasian	Glucose tolerant individuals
Hayakawa	2007	755	Asian	Case-control diabetes study

Figure 1





Figure 2



Legends

Table 1: Characteristics of included sample in the antipsychotic-induced weight gain analysis

Table 2: Characteristics of included sample in the BMI analysis

Figure 1: Meta-analysis results of GNB3 C825T association with antipsychotic-induced weight gain. A) Forest plot of studies assessing the effect of CC variant for lower weight gain: overall genotype effect for dichotomous outcomes. Standardized mean difference (SMD) >1 in favour CC effect in lower weight gain; SMD <1 against CC effect in lower weight gain. B) Begg's funnel plot of publication bias with pseudo 95% confidence limits. C) Egger's publication bias plot.

Figure 1: Meta-analysis results of GNB3 C825T association with BMI. A) Forest plot of studies assessing the effect of CC variant for lower BMI: overall genotype effect for dichotomous outcomes. Standardized mean difference (SMD) >1 in favour CC effect in lower BMI; SMD <1 against CC effect in lower BMI. B) Begg's funnel plot of publication bias with pseudo 95% confidence limits. C) Egger's publication bias plot.

4. Conclusões

Após a análise das 90 variantes mapeados nos genes GSK-3 β , GFR α 1-4, NALCN e GNB-3, uma análise individual destes marcadores sugere que:

- GSK-3 β : três variantes (rs7624540, rs4072520 e rs6779828) mostraram genótipos associados com esquizofrenia, sendo que o rs4072520 permanece significativo após correção por 1.000 permutações. Não fora encontrada associação com a esquizofrenia em nossa amostra de famílias e com a resposta ao tratamento com clozapina.

- GFR α 1-4: GFR α 1 rs11197557 foi associado com a esquizofrenia na amostra de caso-controle pareado e rs730357, bem como alguns haplótipos, mostraram um padrão de transmissão alterado. Embora nenhum dos marcadores em GFR α 1 esteve associado com a resposta ao tratamento de forma individual, dois haplótipos (rs11197612-rs3781514 e rs12413585-rs730057-rs1197612) o estiveram. No gene GFR α 2, três variantes (rs1128397, rs13250096 e rs4567028) e alguns haplótipos mostram-se associados com a resposta ao tratamento. No gene GFR α 3, o rs11242417 e o haplótipo contendo todos os quatro marcadores analisados foram associados com a susceptibilidade à esquizofrenia. Nenhuma associação fora encontrada no gene GFR α 4.

- GPX e MnSOD: não foram encontradas associações com a resposta ao tratamento com clozapina nem com a gravidade de sintomas.

- NALCN: na amostra caso-controle foram encontradas associações nos marcadores rs9518320 e rs9518331 e nos haplótipos compostos por rs7317836, rs9518320, rs9518331, rs2584531 e rs3916906. A amostra de famílias não apresentou nenhuma associação significativa. Uma variantes, rs2152324, esteve associada com a resposta ao tratamento, enquanto um haplótipo formado por rs10508059-rs7328287-rs496238 for a associado ao ganho de peso induzido por clozapina. Cinco variantes foram associadas de maneira individual com a TD (rs9513851, rs9518307, rs9518349, rs10508059 e rs7328287) e dois haplótipos (rs9513851-rs9518307 e rs7328287-rs496238).

- GNB-3: a meta-análise mostrou que a variante CC está associada com um menor índice de massa corpórea, bem como com menor ganho de peso, embora não se alcance significância estatística nessas observações ($p = 0,339$ e $p = 0,053$).

5. Referências bibliográficas

1. Abi-Dargham,A., Martinez,D., Mawlawi,O., Simpson,N., Hwang,D.R., Slifstein,M., Anjilvel,S., Pidcock,J., Guo,N.N., Lombardo,I., Mann,J.J., Van,H.R., Foged,C., Halldin,C. e Laruelle,M. 2000. Measurement of striatal and extrastriatal dopamine D1 receptor binding potential with [11C]NNC 112 in humans: validation and reproducibility. *J.Cereb.Blood Flow Metab*, 20 225-243.
2. Akbarian,S., Kim,J.J., Potkin,S.G., Hetrick,W.P., Bunney,W.E., Jr. e Jones,E.G. 1996. Maldistribution of interstitial neurons in prefrontal white matter of the brains of schizophrenic patients. *Arch.Gen.Psychiatry*, 53 425-436.
3. Allison,D.B., Mentore,J.L., Heo,M., Chandler,L.P., Cappelleri,J.C., Infante,M.C. e Weiden,P.J. 1999. Antipsychotic-induced weight gain: a comprehensive research synthesis. *Am.J.Psychiatry*, 156 1686-1696.
4. Allison,D.B. e Casey,D.E. 2001. Antipsychotic-induced weight gain: a review of the literature. *J.Clin.Psychiatry*, 62 Suppl 7 22-31.
5. American Psychiatric Association. 2000. Diagnostic and statistical manual of mental disorders (4th text revision ed.). Washington, DC.
6. Andreasen,N.C. 1982. Negative symptoms in schizophrenia. Definition and reliability. *Arch.Gen.Psychiatry*, 39 784-788.
7. Anttila,S., Illi,A., Kampman,O., Mattila,K.M., Lehtimaki,T. e Leinonen,E. 2004. Interaction between NOTCH4 and catechol-O-methyltransferase genotypes in schizophrenia patients with poor response to typical neuroleptics. *Pharmacogenetics*, 14 303-307.
8. Arndt,S., Tyrrell,G., Flaum,M. e Andreasen,N.C. 1992. Comorbidity of substance abuse and schizophrenia: the role of pre-morbid adjustment. *Psychol.Med.*, 22 379-388.
9. Arnold,S.E., Talbot,K. e Hahn,C.G. 2005. Neurodevelopment, neuroplasticity, and new genes for schizophrenia. *Prog.Brain Res.*, 147 319-345.
10. Arranz,M., Collier,D., Sodhi,M., Ball,D., Roberts,G., Price,J., Sham,P. e Kerwin,R. 1995. Association between clozapine response and allelic variation in 5-HT_{2A} receptor gene. *Lancet*, 346 281-282.
11. Arranz,M.J., Munro,J., Owen,M.J., Spurlock,G., Sham,P.C., Zhao,J., Kirov,G., Collier,D.A. e Kerwin,R.W. 1998. Evidence for association between polymorphisms in the promoter and coding regions of the 5-HT_{2A} receptor gene and response to clozapine. *Mol.Psychiatry*, 3 61-66.
12. Arranz,M.J., Li,T., Munro,J., Liu,X., Murray,R., Collier,D.A. e Kerwin,R.W. 1998. Lack of association between a polymorphism in the promoter region of the dopamine-2 receptor gene and clozapine response. *Pharmacogenetics*, 8 481-484.

13. Arranz,M.J., Bolonna,A.A., Munro,J., Curtis,C.J., Collier,D.A. e Kerwin,R.W. 2000. The serotonin transporter and clozapine response. *Mol.Psychiatry*, 5 124-125.
14. Arranz,M.J., Munro,J., Birkett,J., Bolonna,A., Mancama,D., Sodhi,M., Lesch,K.P., Meyer,J.F., Sham,P., Collier,D.A., Murray,R.M. e Kerwin,R.W. 2000. Pharmacogenetic prediction of clozapine response. *Lancet*, 355 1615-1616.
15. Arranz,M.J. e de,L.J. 2007. Pharmacogenetics and pharmacogenomics of schizophrenia: a review of last decade of research. *Mol.Psychiatry*, 12 707-747.
16. Arthur,H., Dahl,M.L., Siwers,B. e Sjoqvist,F. 1995. Polymorphic drug metabolism in schizophrenic patients with tardive dyskinesia. *J.Clin.Psychopharmacol.*, 15 211-216.
17. Austin,J. 2005. Schizophrenia: an update and review. *J.Genet.Couns.*, 14 329-340.
18. Ban, T.A., 2001. Pharmacotherapy of mental illness. A historical analysis. *Prog. Neuro-Psychopharmacol. Biol. Psychiatry* 25, 709–727.
19. Ban, T.A., 2002. Neuropsychopharmacology: the interface between genes and psychiatric nosology. In: Lerer, B. (Ed.), *Pharmacogenetics of Psychotropic Drugs*. Cambridge Univ. Press, Cambridge, pp. 36– 56.
20. Basile,V.S., Ozdemir,V., Masellis,M., Walker,M.L., Meltzer,H.Y., Lieberman,J.A., Potkin,S.G., Alva,G., Kalow,W., Macciardi,F.M. e Kennedy,J.L. 2000. A functional polymorphism of the cytochrome P450 1A2 (CYP1A2) gene: association with tardive dyskinesia in schizophrenia. *Mol.Psychiatry*, 5 410-417.
21. Basile,V.S., Masellis,M., McIntyre,R.S., Meltzer,H.Y., Lieberman,J.A. e Kennedy,J.L. 2001. Genetic dissection of atypical antipsychotic-induced weight gain: novel preliminary data on the pharmacogenetic puzzle. *J.Clin.Psychiatry*, 62 Suppl 23 45-66.
22. Becker,W., 1921. Stuporlo"sung durch Kokain. *Psychiatr. Neurol. Wschr.* 22, 19–20.
23. Bernstein,J.G. 1987. Induction of obesity by psychotropic drugs. *Ann.N.Y.Acad.Sci.*, 499 203-215.
24. Bersani,G., Orlandi,V., Kotzalidis,G.D. e Pancheri,P. 2002. Cannabis and schizophrenia: impact on onset, course, psychopathology and outcomes. *Eur.Arch.Psychiatry Clin.Neurosci.*, 252 86-92.
25. Birchwood,M. 1999. Commentary on Garety & Freeman. I: 'Cognitive approaches to delusions--a critical review of theories and evidence'. *Br.J.Clin.Psychol.*, 38 (Pt 3) 315-318.

26. Birkett,J.T., Arranz,M.J., Munro,J., Osbourn,S., Kerwin,R.W. e Collier,D.A. 2000. Association analysis of the 5-HT5A gene in depression, psychosis and antipsychotic response. *Neuroreport*, 11 2017-2020.
27. Bolonna,A.A., Arranz,M.J., Munro,J., Osborne,S., Petouni,M., Martinez,M. e Kerwin,R.W. 2000. No influence of adrenergic receptor polymorphisms on schizophrenia and antipsychotic response. *Neurosci.Lett.*, 280 65-68.
28. Bottlender,R., Strauss,A. e Moller,H.J. 2000. Impact of duration of symptoms prior to first hospitalization on acute outcome in 998 schizophrenic patients. *Schizophr.Res.*, 44 145-150.
29. Bottlender,R., Sato,T., Jager,M., Groll,C., Strauss,A. e Moller,H.J. 2002. The impact of duration of untreated psychosis and premorbid functioning on outcome of first inpatient treatment in schizophrenic and schizoaffective patients. *Eur.Arch.Psychiatry Clin.Neurosci.*, 252 226-231.
30. Bottlender,R., Sato,T. e Moller,H.J. 2003. Summer birth and deficit schizophrenia. *Am.J.Psychiatry*, 160 594-595.
31. Brown,A.S., Schaefer,C.A., Wyatt,R.J., Goetz,R., Begg,M.D., Gorman,J.M. e Susser,E.S. 2000. Maternal exposure to respiratory infections and adult schizophrenia spectrum disorders: a prospective birth cohort study. *Schizophr.Bull.*, 26 287-295.
32. Brown,A.S. e Susser,E.S. 2002. In utero infection and adult schizophrenia. *Ment.Retard.Dev.Disabil.Res.Rev.*, 8 51-57.
33. Brzustowicz,L.M., Hodgkinson,K.A., Chow,E.W., Honer,W.G. e Bassett,A.S. 2000. Location of a major susceptibility locus for familial schizophrenia on chromosome 1q21-q22. *Science*, 288 678-682.
34. Burnet,P.W. e Harrison,P.J. 1995. Genetic variation of the 5-HT2A receptor and response to clozapine. *Lancet*, 346 909
35. Cadet,J.L. e Lohr,J.B. 1989. Possible involvement of free radicals in neuroleptic-induced movement disorders. Evidence from treatment of tardive dyskinesia with vitamin E. *Ann.N.Y.Acad.Sci.*, 570 176-185.
36. Cannon,M., Huttunen,M.O., Tanskanen,A.J., Arseneault,L., Jones,P.B. e Murray,R.M. 2002. Perinatal and childhood risk factors for later criminality and violence in schizophrenia. Longitudinal, population-based study. *Br.J.Psychiatry*, 180 496-501.
37. Chakos,M., Lieberman,J., Hoffman,E., Bradford,D. e Sheitman,B. 2001. Effectiveness of second-generation antipsychotics in patients with treatment-resistant schizophrenia: a review and meta-analysis of randomized trials. *Am.J.Psychiatry*, 158 518-526.

38. Chakravarti,A. 1999. Population genetics--making sense out of sequence. *Nat.Genet.*, 21 56-60.
39. Chakravarti,A. e Little,P. 2003. Nature, nurture and human disease. *Nature*, 421 412-414.
40. Charpentier, P., Gailliot, P., Jacob, R., Gaudechon, J. e Buisson, P., 1952. Recherches sur les dimethylaminopropyl-N-phenothiazines substitue`es. *C. R. Acad. Sci. (Paris)* 235, 59– 60.
41. Chen,C.H., Wei,F.C., Koong,F.J. e Hsiao,K.J. 1997. Association of TaqI A polymorphism of dopamine D2 receptor gene and tardive dyskinesia in schizophrenia. *Biol.Psychiatry*, 41 827-829.
42. Chong,S.A., Tan,E.C., Tan,C.H., Mahendren,R., Tay,A.H. e Chua,H.C. 2000. Tardive dyskinesia is not associated with the serotonin gene polymorphism (5-HTTLPR) in Chinese. *Am.J.Med.Genet.*, 96 712-715.
43. Chowdari,K.V., Mirnics,K., Semwal,P., Wood,J., Lawrence,E., Bhatia,T., Deshpande,S.N., B K T, Ferrell,R.E., Middleton,F.A., Devlin,B., Levitt,P., Lewis,D.A. e Nimgaonkar,V.L. 2002. Association and linkage analyses of RGS4 polymorphisms in schizophrenia. *Hum.Mol.Genet.*, 11 1373-1380.
44. Chumakov,I., Blumenfeld,M., Guerassimenko,O., Cavarec,L., Palicio,M., Abderrahim,H., Bougueleret,L., Barry,C., Tanaka,H., La,R.P., Puech,A., Tahri,N., Cohen-Akenine,A., Delabrosse,S., Lissarrague,S., Picard,F.P., Maurice,K., Essioux,L., Millasseau,P., Grel,P., Debailleul,V., Simon,A.M., Caterina,D., Dufaure,I., Malekzadeh,K., Belova,M., Luan,J.J., Bouillot,M., Sambucy,J.L., Primas,G., Saumier,M., Boubkiri,N., Martin-Saumier,S., Nasroune,M., Peixoto,H., Delaye,A., Pinchot,V., Bastucci,M., Guillou,S., Chevillon,M., Sainz-Fuertes,R., Meguenni,S., urich-Costa,J., Cherif,D., Gimalac,A., Van,D.C., Gauvreau,D., Ouellette,G., Fortier,I., Raelson,J., Sherbatich,T., Riazanskaia,N., Rogaev,E., Raeymaekers,P., Aerssens,J., Konings,F., Luyten,W., Macciardi,F., Sham,P.C., Straub,R.E., Weinberger,D.R., Cohen,N. e Cohen,D. 2002. Genetic and physiological data implicating the new human gene G72 and the gene for D-amino acid oxidase in schizophrenia. *Proc.Natl.Acad.Sci.U.S.A.*, 99 13675-13680.
45. Conneally,P.M. 2003. The complexity of complex diseases. *Am.J.Hum.Genet.*, 72 229-232.
46. Coyle,J.T. 2006. Glutamate and schizophrenia: beyond the dopamine hypothesis. *Cell Mol.Neurobiol.*, 26 365-384.
47. Craddock,N., O'Donovan,M.C. e Owen,M.J. 2006. Genes for schizophrenia and bipolar disorder? Implications for psychiatric nosology. *Schizophr.Bull.*, 32 9-16.
48. Croce, G., 1932. Pyretotherapy of schizophrenia in initial stages by injection of sulphur. *Am. J. Psychiatry* 132, 1237–1245.

49. Crow,T.J. 1985. The two-syndrome concept: origins and current status. *Schizophr.Bull.*, 11 471-486.
50. Davidson,L. e McGlashan,T.H. 1997. The varied outcomes of schizophrenia. *Can.J.Psychiatry*, 42 34-43.
51. Davis,K.L., Stewart,D.G., Friedman,J.I., Buchsbaum,M., Harvey,P.D., Hof,P.R., Buxbaum,J. e Haroutunian,V. 2003. White matter changes in schizophrenia: evidence for myelin-related dysfunction. *Arch.Gen.Psychiatry*, 60 443-456.
52. Davis,R. e Faulds,D. 1996. Dexfenfluramine. An updated review of its therapeutic use in the management of obesity. *Drugs*, 52 696-724.
53. Dawson,M.E. e Nuechterlein,K.H. 1984. Psychophysiological dysfunctions in the developmental course of schizophrenic disorders. *Schizophr.Bull.*, 10 204-232.
54. Delay, J., Deniker, P. e Harl, J., 1952. Traitement des etats d'excitation et d'agitation par une methode medicamenteuse derive de l'hibernotherapie. *Ann. Medicopsychol. (Paris)* 119, 267– 273.
55. Delay, J., Deniker, P., Tardieu, Y. e Lemperiere, T., 1954. Le R1625, nouvelle therapeutique psychiatrique de la reserpina, alcaloide nouveau de la Rauwolfia serpentina. *C. R. Congr. Alien Neurol. France* 52, 836– 841.
56. de Leon, J., Susce,M.T., Pan,R.M., Koch,W.H. e Wedlund,P.J. 2005. Polymorphic variations in GSTM1, GSTT1, PgP, CYP2D6, CYP3A5, and dopamine D2 and D3 receptors and their association with tardive dyskinesia in severe mental illness. *J.Clin.Psychopharmacol.*, 25 448-456.
57. de Leon, J., Susce,M.T. e Murray-Carmichael,E. 2006. The AmpliChip CYP450 genotyping test: Integrating a new clinical tool. *Mol.Diagn.Ther.*, 10 135-151.
58. de Luca, V, Vincent,J.B., Muller,D.J., Hwang,R., Shinkai,T., Volavka,J., Czobor,P., Sheitman,B.B., Lindenmayer,J.P., Citrome,L., McEvoy,J.P., Lieberman,J.A. e Kennedy,J.L. 2005. Identification of a naturally occurring 21 bp deletion in alpha 2c noradrenergic receptor gene and cognitive correlates to antipsychotic treatment. *Pharmacol.Res.*, 51 381-384.
59. de Lisi,L.E., Shaw,S.H., Crow,T.J., Shields,G., Smith,A.B., Larach,V.W., Wellman,N., Loftus,J., Nanthakumar,B., Razi,K., Stewart,J., Comazzi,M., Vita,A., Heffner,T. e Sherrington,R. 2002. A genome-wide scan for linkage to chromosomal regions in 382 sibling pairs with schizophrenia or schizoaffective disorder. *Am.J.Psychiatry*, 159 803-812.
60. Divry, P., Bobon, J. e Collard, J., 1959. Le R1625, nouvelle therapeutique symptomatique de l'agitation psychometrice. *Acta Neurol. Psychiatr. Belg.* 58, 878–888.

61. Dohrenwend,B.P., Levav,I., ShROUT,P.E., Schwartz,S., Naveh,G., Link,B.G., Skodol,A.E. e Stueve,A. 1992. Socioeconomic status and psychiatric disorders: the causation-selection issue. *Science*, 255 946-952.
62. Ekelund,J., Hennah,W., Hiekkalinna,T., Parker,A., Meyer,J., Lonnqvist,J. e Peltonen,L. 2004. Replication of 1q42 linkage in Finnish schizophrenia pedigrees. *Mol.Psychiatry*, 9 1037-1041.
63. Ellingrod,V.L., Miller,D., Schultz,S.K., Wehring,H. e Arndt,S. 2002. CYP2D6 polymorphisms and atypical antipsychotic weight gain. *Psychiatr.Genet.*, 12 55-58.
64. Falkai,P., Wobrock,T., Lieberman,J., Glenthøj,B., Gattaz,W.F. e Moller,H.J. 2005. World Federation of Societies of Biological Psychiatry (WFSBP) guidelines for biological treatment of schizophrenia, Part 1: acute treatment of schizophrenia. *World J.Biol.Psychiatry*, 6 132-191.
65. Faurvye, A., Rasch,P.J., Petersen,P.B., Brandborg,G. E Pakkenberg,H. 1964. Neurological symptoms in pharmacotherapy of psychoses. *Acta Psychiatr.Scand.*, 40 10-27.
66. Fenton,W.S. e McGlashan,T.H. 1987. Sustained remission in drug-free schizophrenic patients. *Am.J.Psychiatry*, 144 1306-1309.
67. Freedman,R. e Leonard,S. 2001. Genetic linkage to schizophrenia at chromosome 15q14. *Am.J.Med.Genet.*, 105 655-657.
68. Freedman,R. 2003. Schizophrenia. *N.Engl.J.Med.*, 349 1738-1749.
69. Gaebel,W., Frick,U., Kopcke,W., Linden,M., Muller,P., Muller-Spahn,F., Pietzcker,A. e Tegeler,J. 1993. Early neuroleptic intervention in schizophrenia: are prodromal symptoms valid predictors of relapse? *Br.J.Psychiatry Suppl*, 8-12.
70. Glatt,S.J., Faraone,S.V. e Tsuang,M.T. 2003. Association between a functional catechol O-methyltransferase gene polymorphism and schizophrenia: meta-analysis of case-control and family-based studies. *Am.J.Psychiatry*, 160 469-476.
71. Gottesman, II. 1991. *Schizophrenia Genesis: The Origins of Madness* New York, NY W.H. Freeman and Company.
72. Gottesman, II e Bertelsen, A. 1989. Confirming unexpressed genotypes for schizophrenia. *Arch. General Psychiatry* 46 867-872
73. Gottesman,II. e Shields,J. 1967. A polygenic theory of schizophrenia. *Proc.Natl.Acad.Sci.U.S.A*, 58 199-205.
74. Goudie,A.J., Cooper,G.D. e Halford,J.C. 2005. Antipsychotic-induced weight gain. *Diabetes Obes.Metab*, 7 478-487.

75. Greenbaum,L., Strous,R.D., Kanyas,K., Merbl,Y., Horowitz,A., Karni,O., Katz,E., Kotler,M., Olender,T., Deshpande,S.N., Lancet,D., Ben-Asher,E. e Lerer,B. 2007. Association of the RGS2 gene with extrapyramidal symptoms induced by treatment with antipsychotic medication. *Pharmacogenet.Genomics*, 17 519-528.
76. Guest,J.F. e Cookson,R.F. 1999. Cost of schizophrenia to UK Society. An incidence-based cost-of-illness model for the first 5 years following diagnosis. *Pharmacoeconomics.*, 15 597-610.
77. Gurling,H.M., Kalsi,G., Brynjolfson,J., Sigmundsson,T., Sherrington,R., Mankoo,B.S., Read,T., Murphy,P., Blaveri,E., McQuillin,A., Petursson,H. e Curtis,D. 2001. Genomewide genetic linkage analysis confirms the presence of susceptibility loci for schizophrenia, on chromosomes 1q32.2, 5q33.2, and 8p21-22 and provides support for linkage to schizophrenia, on chromosomes 11q23.3-24 and 20q12.1-11.23. *Am.J.Hum.Genet.*, 68 661-673.
78. Hafner,H. 2003. Gender differences in schizophrenia. *Psychoneuroendocrinology*, 28 Suppl 2 17-54.
79. Hamilton, M. (Ed.), 1976. *Fish's Schizophrenia*. John Wright & Sons, Bristol, pp. 2– 4.
80. Hamshere,M.L., Williams,N.M., Norton,N., Williams,H., Cardno,A.G., Zammit,S., Jones,L.A., Murphy,K.C., Sanders,R.D., McCarthy,G., Gray,M.Y., Jones,G., Holmans,P., O'Donovan,M.C., Owen,M.J. e Craddock,N. 2006. Genome wide significant linkage in schizophrenia conditioning on occurrence of depressive episodes. *J.Med.Genet.*, 43 563-567.
81. Harrison,P.J. 1999. The neuropathology of schizophrenia. A critical review of the data and their interpretation. *Brain*, 122 593-624.
82. Hegarty,J.D., Baldessarini,R.J., Tohen,M., Wateraux,C. e Oepen,G. 1994. One hundred years of schizophrenia: a meta-analysis of the outcome literature. *Am.J.Psychiatry*, 151 1409-1416.
83. Henderson,D.C., Cagliero,E., Gray,C., Nasrallah,R.A., Hayden,D.L., Schoenfeld,D.A. e Goff,D.C. 2000. Clozapine, diabetes mellitus, weight gain, and lipid abnormalities: A five-year naturalistic study. *Am.J.Psychiatry*, 157 975-981.
84. Henderson,D.C. 2007. Weight gain with atypical antipsychotics: evidence and insights. *J.Clin.Psychiatry*, 68 Suppl 12 18-26.
85. Heresco-Levy,U., Ermilov,M., Shimoni,J., Shapira,B., Silipo,G. e Javitt,D.C. 2002. Placebo-controlled trial of D-cycloserine added to conventional neuroleptics, olanzapine, or risperidone in schizophrenia. *Am.J.Psychiatry*, 159 480-482.

86. Herken,H., Erdal,M.E., Boke,O. e Savas,H.A. 2003. Tardive dyskinesia is not associated with the polymorphisms of 5-HT_{2A} receptor gene, serotonin transporter gene and catechol-o-methyltransferase gene. *Eur.Psychiatry*, 18 77-81.
87. Hoenberg, K., Goetz, K., 2006. *Antipsychotics: Analysis of Disease Markets and Emerging Agents*. Decision Resources, Inc., Waltham, Massachusetts.
88. Hoff,A.L., Sakuma,M., Wieneke,M., Horon,R., Kushner,M. e DeLisi,L.E. 1999. Longitudinal neuropsychological follow-up study of patients with first-episode schizophrenia. *Am.J.Psychiatry*, 156 1336-1341.
89. Hong,C.J., Yu,Y.W., Lin,C.H., Song,H.L., Lai,H.C., Yang,K.H. e Tsai,S.J. 2000. Association study of apolipoprotein E epsilon4 with clinical phenotype and clozapine response in schizophrenia. *Neuropsychobiology*, 42 172-174.
90. Hong,C.J., Yu,Y.W., Lin,C.H., Cheng,C.Y. e Tsai,S.J. 2001. Association analysis for NMDA receptor subunit 2B (GRIN2B) genetic variants and psychopathology and clozapine response in schizophrenia. *Psychiatr.Genet.*, 11 219-222.
91. Hong,C.J., Lin,C.H., Yu,Y.W., Yang,K.H. e Tsai,S.J. 2001. Genetic variants of the serotonin system and weight change during clozapine treatment. *Pharmacogenetics*, 11 265-268.
92. Hong,C.J., Liu,H.C., Liu,T.Y., Liao,D.L. e Tsai,S.J. 2005. Association studies of the adenosine A_{2a} receptor (1976T > C) genetic polymorphism in Parkinson's disease and schizophrenia. *J.Neural Transm.*, 112 1503-1510.
93. Hori,H., Ohmori,O., Shinkai,T., Kojima,H., Okano,C., Suzuki,T. e Nakamura,J. 2000. Manganese superoxide dismutase gene polymorphism and schizophrenia: relation to tardive dyskinesia. *Neuropsychopharmacology*, 23 170-177.
94. Huerdo-Diaz,P., Arranz,M.J., Munro,J., Osborne,S., Makoff,A., Kerwin,R.W., Austin,J. e O'Donovan,M. 2004. An association study of the neurotensin receptor gene with schizophrenia and clozapine response. *Schizophr.Res.*, 66 193-195.
95. Illi,A., Kampman,O., Anttila,S., Roivas,M., Mattila,K.M., Lehtimaki,T. e Leinonen,E. 2003. Interaction between angiotensin-converting enzyme and catechol-O-methyltransferase genotypes in schizophrenics with poor response to conventional neuroleptics. *Eur.Neuropsychopharmacol.*, 13 147-151.
96. Ingham, S.D., 1930. Favorable results in dementia praecox with the use of castor oil and force feeding. *Tr. Am. Neurol.* 56, 401– 407.
97. Jacquet,H., Raux,G., Thibaut,F., Hecketsweiler,B., Houy,E., Demilly,C., Haouzir,S., Allio,G., Fouldrin,G., Drouin,V., Bou,J., Petit,M., Campion,D. e Frebourg,T. 2002. PRODH mutations and hyperprolinemia in a subset of schizophrenic patients. *Hum.Mol.Genet.*, 11 2243-2249.

98. Janssen, P.A.J., 1996. Haloperidol and the butyrophenones: the early years. In: Ban, T.A., Ray, O.S. (Eds.), *A History of the CINP*. JM Productions, Brentwood, pp. 44–48.
99. Javitt, D.C. e Zukin, S.R. 1991. Recent advances in the phencyclidine model of schizophrenia. *Am.J.Psychiatry*, 148 1301-1308.
100. Javitt, D.C. e Coyle, J.T. 2004. Decoding schizophrenia. *Sci.Am.*, 290 48-55.
101. Kampman, O., Anttila, S., Illi, A., Saarela, M., Rontu, R., Mattila, K.M., Leinonen, E. e Lehtimäki, T. 2004. Neuregulin genotype and medication response in Finnish patients with schizophrenia. *Neuroreport*, 15 2517-2520.
102. Kampman, O., Illi, A., Hanninen, K., Katila, H., Anttila, S., Rontu, R., Mattila, K.M., Leinonen, E. e Lehtimäki, T. 2006. RGS4 genotype is not associated with antipsychotic medication response in schizophrenia. *J.Neural Transm.*, 113 1563-1568.
103. Kane, J.M. 1982. Coping with a drug-induced movement disorder: tardive dyskinesia. *Geriatrics*, 37 83-87.
104. Kane, J.M., Honigfeld, G., Singer, J. e Meltzer, H. 1988. Clozapine in treatment-resistant schizophrenics. *Psychopharmacol.Bull.*, 24 62-67.
105. Kane, J.M. 2001. Extrapyramidal side effects are unacceptable. *Eur.Neuropsychopharmacol.*, 11 Suppl 4 S397-S403
106. Kane, J.M., Krystal, J. e Correll, C.U. 2003. Treatment models and designs for intervention research during the psychotic prodrome. *Schizophr.Bull.*, 29 747-756.
107. Kapur, S. e Remington, G. 2001. Dopamine D(2) receptors and their role in atypical antipsychotic action: still necessary and may even be sufficient. *Biol.Psychiatry*, 50 873-883.
108. Kendler, K.S. e Zerbin-Rudin, E. 1996. Abstract and review of "Zur Erbpathologie der Schizophrenie" (Contribution to the genetics of schizophrenia). 1916. *Am.J.Med.Genet.*, 67 343-346.
109. Kendler, K.S., Karkowski-Shuman, L. e Walsh, D. 1996. Age at onset in schizophrenia and risk of illness in relatives. Results from the Roscommon Family Study. *Br.J.Psychiatry*, 169 213-218.
110. Knapp, M. 1997. Costs of schizophrenia. *Br.J.Psychiatry*, 171 509-518.
111. Kovasznay, B., Fleischer, J., Tanenberg-Karant, M., Jandorf, L., Miller, A.D. e Bromet, E. 1997. Substance use disorder and the early course of illness in schizophrenia and affective psychosis. *Schizophr.Bull.*, 23 195-201.
112. Kraepelin, E., 1913. *Lehrbuch der Psychiatrie*. 8 Aufl Barth, Leipzig.

113. Kyle,U.G. e Pichard,C. 2006. The Dutch Famine of 1944-1945: a pathophysiological model of long-term consequences of wasting disease. *Curr.Opin.Clin.Nutr.Metab Care*, 9 388-394.
114. Lai,I.C., Liao,D.L., Bai,Y.M., Lin,C.C., Yu,S.C., Chen,J.Y. e Wang,Y.C. 2002. Association study of the estrogen receptor polymorphisms with tardive dyskinesia in schizophrenia. *Neuropsychobiology*, 46 173-175.
115. Lane,H.Y., Chang,Y.C., Liu,Y.C., Chiu,C.C. e Tsai,G.E. 2005. Sarcosine or D-serine add-on treatment for acute exacerbation of schizophrenia: a randomized, double-blind, placebo-controlled study. *Arch.Gen.Psychiatry*, 62 1196-1204.
116. Lane,H.Y., Liu,Y.C., Huang,C.L., Chang,Y.C., Wu,P.L., Lu,C.T. e Chang,W.H. 2006. Risperidone-related weight gain: genetic and nongenetic predictors. *J.Clin.Psychopharmacol.*, 26 128-134.
117. Lawrie,S.M., Whalley,H., Kestelman,J.N., Abukmeil,S.S., Byrne,M., Hodges,A., Rimmington,J.E., Best,J.J., Owens,D.G. e Johnstone,E.C. 1999. Magnetic resonance imaging of brain in people at high risk of developing schizophrenia. *Lancet*, 353 30-33.
118. Lee,H.J., Kang,S.G., Paik,J.W., Lee,M.S., Cho,B.H., Park,Y.M., Kim,W., Choi,J.E., Jung,I.K., Kim,L. e Lee,M.S. 2007. No evidence for an association between G protein beta3 subunit gene C825T polymorphism and tardive dyskinesia in schizophrenia. *Hum.Psychopharmacol.*, 22 501-504.
119. Lehmann, H.E., 1993. Before they called it psychopharmacology. *Neuropsychopharmacology*, 8, 291– 303.
120. Lehman,A.F., Lieberman,J.A., Dixon,L.B., McGlashan,T.H., Miller,A.L., Perkins,D.O. e Kreyenbuhl,J. 2004. Practice guideline for the treatment of patients with schizophrenia, second edition. *Am.J.Psychiatry*, 161 1-56.
121. Lerer,B., Segman,R.H., Hamdan,A., Kanyas,K., Karni,O., Kohn,Y., Korner,M., Lanktree,M., Kaadan,M., Turetsky,N., Yakir,A., Kerem,B. e Macciardi,F. 2003. Genome scan of Arab Israeli families maps a schizophrenia susceptibility gene to chromosome 6q23 and supports a locus at chromosome 10q24. *Mol.Psychiatry*, 8 488-498.
122. Lewis,C.M., Levinson,D.F., Wise,L.H., DeLisi,L.E., Straub,R.E., Hovatta,I., Williams,N.M., Schwab,S.G., Pulver,A.E., Faraone,S.V., Brzustowicz,L.M., Kaufmann,C.A., Garver,D.L., Gurling,H.M., Lindholm,E., Coon,H., Moises,H.W., Byerley,W., Shaw,S.H., Mesen,A., Sherrington,R., O'Neill,F.A., Walsh,D., Kendler,K.S., Ekelund,J., Paunio,T., Lonnqvist,J., Peltonen,L., O'Donovan,M.C., Owen,M.J., Wildenauer,D.B., Maier,W., Nestadt,G., Blouin,J.L., Antonarakis,S.E., Mowry,B.J., Silverman,J.M., Crowe,R.R., Cloninger,C.R., Tsuang,M.T., Malaspina,D., Harkavy-Friedman,J.M., Svrakic,D.M., Bassett,A.S., Holcomb,J., Kalsi,G., McQuillin,A., Brynjolfson,J., Sigmundsson,T., Petursson,H., Jazin,E., Zoega,T. e Helgason,T. 2003. Genome scan meta-analysis of schizophrenia and bipolar disorder, part II: Schizophrenia. *Am.J.Hum.Genet.*, 73 34-48.

123. Lewis,D.A., Hashimoto,T. e Volk,D.W. 2005. Cortical inhibitory neurons and schizophrenia. *Nat.Rev.Neurosci.*, 6 312-324.
124. Lewis,S.W. 1987. Causes of schizophrenia. *Lancet*, 1 1216-1217.
125. Lewis,S.W., Barnes,T.R., Davies,L., Murray,R.M., Dunn,G., Hayhurst,K.P., Markwick,A., Lloyd,H. e Jones,P.B. 2006. Randomized controlled trial of effect of prescription of clozapine versus other second-generation antipsychotic drugs in resistant schizophrenia. *Schizophr.Bull.*, 32 715-723.
126. Lieberman,J.A., Phillips,M., Gu,H., Stroup,S., Zhang,P., Kong,L., Ji,Z., Koch,G. e Hamer,R.M. 2003. Atypical and conventional antipsychotic drugs in treatment-naive first-episode schizophrenia: a 52-week randomized trial of clozapine vs chlorpromazine. *Neuropsychopharmacology*, 28 995-1003.
127. Lin,Y.C., Ellingrod,V.L., Bishop,J.R. e Miller,d.D. 2006. The relationship between P-glycoprotein (PGP) polymorphisms and response to olanzapine treatment in schizophrenia. *Ther.Drug Monit.*, 28 668-672.
128. Linszen,D.H., Dingemans,P.M. e Lenior,M.E. 1994. Cannabis abuse and the course of recent-onset schizophrenic disorders. *Arch.Gen.Psychiatry*, 51 273-279.
129. Liou,Y.J., Liao,D.L., Chen,J.Y., Wang,Y.C., Lin,C.C., Bai,Y.M., Yu,S.C., Lin,M.W. e Lai,I.C. 2004. Association analysis of the dopamine D3 receptor gene ser9gly and brain-derived neurotrophic factor gene val66met polymorphisms with antipsychotic-induced persistent tardive dyskinesia and clinical expression in Chinese schizophrenic patients. *Neuromolecular.Med.*, 5 243-251.
130. Liou,Y.J., Wang,Y.C., Chen,J.Y., Bai,Y.M., Lin,C.C., Liao,D.L., Chen,T.T., Chen,M.L., Mo,G.H. e Lai,I.C. 2007. Association analysis of polymorphisms in the N-methyl-D-aspartate (NMDA) receptor subunit 2B (GRIN2B) gene and tardive dyskinesia in schizophrenia. *Psychiatry Res.*, 153 271-275.
131. Loebel,A.D., Lieberman,J.A., Alvir,J.M., Mayerhoff,D.I., Geisler,S.H. e Szymanski,S.R. 1992. Duration of psychosis and outcome in first-episode schizophrenia. *Am.J.Psychiatry*, 149 1183-1188.
132. Malhotra,A.K., Goldman,D., Ozaki,N., Breier,A., Buchanan,R. e Pickar,D. 1996. Lack of association between polymorphisms in the 5-HT2A receptor gene and the antipsychotic response to clozapine. *Am.J.Psychiatry*, 153 1092-1094.
133. Malhotra,A.K., Murphy,G.M., Jr. e Kennedy,J.L. 2004. Pharmacogenetics of psychotropic drug response. *Am.J.Psychiatry*, 161 780-796.
134. Malhotra,A.K., Lencz,T., Correll,C.U. e Kane,J.M. 2007. Genomics and the future of pharmacotherapy in psychiatry. *Int.Rev.Psychiatry*, 19 523-530.

135. Mancama,D., Arranz,M.J., Munro,J., Osborne,S., Makoff,A., Collier,D. e Kerwin,R. 2002. Investigation of promoter variants of the histamine 1 and 2 receptors in schizophrenia and clozapine response. *Neurosci.Lett.*, 333 207-211.
136. Mancama,D., Mata,I., Kerwin,R.W. e Arranz,M.J. 2007. Choline acetyltransferase variants and their influence in schizophrenia and olanzapine response. *Am.J.Med.Genet.B Neuropsychiatr.Genet.*, 144 849-853.
137. Marcelis,M., Takei,N. e van,O.J. 1999. Urbanization and risk for schizophrenia: does the effect operate before or around the time of illness onset? *Psychol.Med.*, 29 1197-1203.
138. Marenco,S. e Weinberger,D.R. 2000. The neurodevelopmental hypothesis of schizophrenia: following a trail of evidence from cradle to grave. *Dev.Psychopathol.*, 12 501-527.
139. Masellis,M., Paterson,A.D., Badri,F., Lieberman,J.A., Meltzer,H.Y., Cavazzoni,P. e Kennedy,J.L. 1995. Genetic variation of 5-HT_{2A} receptor and response to clozapine. *Lancet*, 346 1108
140. Masellis,M., Basile,V.S., Meltzer,H.Y., Lieberman,J.A., Sevy,S., Goldman,D.A., Hamblin,M.W., Macciardi,F.M. e Kennedy,J.L. 2001. Lack of association between the T->C 267 serotonin 5-HT₆ receptor gene (HTR6) polymorphism and prediction of response to clozapine in schizophrenia. *Schizophr.Res.*, 47 49-58.
141. Matsumoto,C., Shinkai,T., Hori,H., Ohmori,O. e Nakamura,J. 2004. Polymorphisms of dopamine degradation enzyme (COMT and MAO) genes and tardive dyskinesia in patients with schizophrenia. *Psychiatry Res.*, 127 1-7.
142. McEvoy,J.P. 2006. An overview of the Clinical Antipsychotic Trials of Intervention Effectiveness (CATIE) study. *CNS.Spectr.*, 11 4-8.
143. McGue, M. e Gottesman, I.I. (1989). Genetic linkage in schizophrenia: Perspectives from genetic epidemiology. *Schizophrenia Bulletin*, 15: 453-464.
144. Meary,A., Brousse,G., Jamain,S., Schmitt,A., Szoke,A., Schurhoff,F., Gavaudan,G., Lancon,C., quin-Mavier,I., Leboyer,M. e Llorca,P.M. 2007. Pharmacogenetic study of atypical antipsychotic drug response: Involvement of the norepinephrine transporter gene. *Am.J.Med.Genet.B Neuropsychiatr.Genet.*,
145. Meltzer,H.Y. 1989. Clinical studies on the mechanism of action of clozapine: the dopamine-serotonin hypothesis of schizophrenia. *Psychopharmacology (Berl)*, 99 Suppl S18-S27
146. Meltzer,H.Y. 2001. Putting metabolic side effects into perspective: risks versus benefits of atypical antipsychotics. *J.Clin.Psychiatry*, 62 Suppl 27 35-39.

147. Meltzer,H.Y., Alphs,L., Green,A.I., Altamura,A.C., Anand,R., Bertoldi,A., Bourgeois,M., Chouinard,G., Islam,M.Z., Kane,J., Krishnan,R., Lindenmayer,J.P. e Potkin,S. 2003. Clozapine treatment for suicidality in schizophrenia: International Suicide Prevention Trial (InterSePT). *Arch.Gen.Psychiatry*, 60 82-91.
148. Meltzer,H.Y. 2004. What's atypical about atypical antipsychotic drugs? *Curr.Opin.Pharmacol.*, 4 53-57.
149. Millar,J.K., Wilson-Annan,J.C., Anderson,S., Christie,S., Taylor,M.S., Semple,C.A., Devon,R.S., Clair,D.M., Muir,W.J., Blackwood,D.H. e Porteous,D.J. 2000. Disruption of two novel genes by a translocation co-segregating with schizophrenia. *Hum.Mol.Genet.*, 9 1415-1423.
150. Miller,A.L., Hall,C.S., Buchanan,R.W., Buckley,P.F., Chiles,J.A., Conley,R.R., Crismon,M.L., Ereshefsky,L., Essock,S.M., Finnerty,M., Marder,S.R., Miller,D.D., McEvoy,J.P., Rush,A.J., Saeed,S.A., Schooler,N.R., Shon,S.P., Stroup,S. e Tarin-Godoy,B. 2004. The Texas Medication Algorithm Project antipsychotic algorithm for schizophrenia: 2003 update. *J.Clin.Psychiatry*, 65 500-508.
151. Mimmack,M.L., Ryan,M., Baba,H., Navarro-Ruiz,J., Iritani,S., Faull,R.L., McKenna,P.J., Jones,P.B., Arai,H., Starkey,M., Emson,P.C. e Bahn,S. 2002. Gene expression analysis in schizophrenia: reproducible up-regulation of several members of the apolipoprotein L family located in a high-susceptibility locus for schizophrenia on chromosome 22. *Proc.Natl.Acad.Sci.U.S.A.*, 99 4680-4685.
152. Mirnics,K., Middleton,F.A., Lewis,D.A. e Levitt,P. 2001. The human genome: gene expression profiling and schizophrenia. *Am.J.Psychiatry*, 158 1384
153. Moller,H.J. 2000. Definition, psychopharmacological basis and clinical evaluation of novel/atypical neuroleptics: methodological issues and clinical consequences. *World J.Biol.Psychiatry*, 1 75-91.
154. Moller,H.J. 2003. Management of the negative symptoms of schizophrenia: new treatment options. *CNS.Drugs*, 17 793-823.
155. Muller,D.J., Schulze,T.G., Knapp,M., Held,T., Krauss,H., Weber,T., Ahle,G., Maroldt,A., Alfter,D., Maier,W., Nothen,M.M. e Rietschel,M. 2001. Familial occurrence of tardive dyskinesia. *Acta Psychiatr.Scand.*, 104 375-379.
156. Muller,D.J., De,L., V, Sicard,T., King,N., Hwang,R., Volavka,J., Czobor,P., Sheitman,B.B., Lindenmayer,J.P., Citrome,L., McEvoy,J.P., Lieberman,J.A., Meltzer,H.Y. e Kennedy,J.L. 2005. Suggestive association between the C825T polymorphism of the G-protein beta3 subunit gene (GNB3) and clinical improvement with antipsychotics in schizophrenia. *Eur.Neuropsychopharmacol.*, 15 525-531.
157. Muller,D.J., Klempan,T.A., De,L., V, Sicard,T., Volavka,J., Czobor,P., Sheitman,B.B., Lindenmayer,J.P., Citrome,L., McEvoy,J.P., Lieberman,J.A., Honer,W.G. e Kennedy,J.L. 2005. The SNAP-25 gene may be associated with clinical

response and weight gain in antipsychotic treatment of schizophrenia. *Neurosci.Lett.*, 379 81-89.

158. Muller,D.J. e Kennedy,J.L. 2006. Genetics of antipsychotic treatment emergent weight gain in schizophrenia. *Pharmacogenomics.*, 7 863-887.

159. Mueller, J.M., Schlittler, E. e Bein, H.J., 1952. Reserpin, der sedative wirkstoff aus *Rauwolfia serpentine* Benth. *Experientia* 8, 338– 339.

160. Murray,R.M. e Lewis,S.W. 1987. Is schizophrenia a neurodevelopmental disorder? *Br.Med.J.(Clin.Res.Ed)*, 295 681-682.

161. Nothen,M.M., Rietschel,M., Erdmann,J., Oberlander,H., Moller,H.J., Nober,D. e Propping,P. 1995. Genetic variation of the 5-HT_{2A} receptor and response to clozapine. *Lancet*, 346 908-909.

162. NIH. 1998. Clinical guidelines on the identification evaluation and treatment of overweight and obesity in adults. The evidence report. *Obes. Res.*, 6 51S-209S.

163. Ohmori,O., Shinkai,T., Hori,H., Kojima,H. e Nakamura,J. 2001. Polymorphisms of mu and delta opioid receptor genes and tardive dyskinesia in patients with schizophrenia. *Schizophr.Res.*, 52 137-138.

164. Ohmori,O., Shinkai,T., Hori,H. e Nakamura,J. 2002. Genetic association analysis of 5-HT₆ receptor gene polymorphism (267C/T) with tardive dyskinesia. *Psychiatry Res.*, 110 97-102.

165. Owen,M.J., Williams,N.M. e O'Donovan,M.C. 2004. The molecular genetics of schizophrenia: new findings promise new insights. *Mol.Psychiatry*, 9 14-27.

166. Owen,M.J., Craddock,N. e O'Donovan,M.C. 2005. Schizophrenia: genes at last? *Trends Genet.*, 21 518-525.

167. Ozdemir,V., Basile,V.S., Masellis,M. e Kennedy,J.L. 2001. Pharmacogenetic assessment of antipsychotic-induced movement disorders: contribution of the dopamine D₃ receptor and cytochrome P450 1A₂ genes. *J.Biochem.Biophys.Methods*, 47 151-157.

168. Pae,C.U., Yu,H.S., Kim,J.J., Lee,C.U., Lee,S.J., Jun,T.Y., Lee,C. e Paik,I.H. 2004. Quinone oxidoreductase (NQO1) gene polymorphism (609C/T) may be associated with tardive dyskinesia, but not with the development of schizophrenia. *Int.J.Neuropsychopharmacol.*, 7 495-500.

169. Pantelis,C., Velakoulis,D., McGorry,P.D., Wood,S.J., Suckling,J., Phillips,L.J., Yung,A.R., Bullmore,E.T., Brewer,W., Soulsby,B., Desmond,P. e McGuire,P.K. 2003. Neuroanatomical abnormalities before and after onset of psychosis: a cross-sectional and longitudinal MRI comparison. *Lancet*, 361 281-288.

170. Potkin,S.G., Basile,V.S., Jin,Y., Masellis,M., Badri,F., Keator,D., Wu,J.C., Alva,G., Carreon,D.T., Bunney,W.E., Jr., Fallon,J.H. e Kennedy,J.L. 2003. D1 receptor alleles predict PET metabolic correlates of clinical response to clozapine. *Mol.Psychiatry*, 8 109-113.
171. Reed, G.E., 1929. The use of manganese chloride in dementia praecox. *CMAJ* 21, 46– 49.
172. U.S. Department of Health e Human Services 1998 Mental Health: A Report of the Surgeon General.
173. Richardson,M.A., Chao,H.M., Read,L.L., Clelland,J.D. e Suckow,R.F. 2006. Investigation of the phenylalanine hydroxylase gene and tardive dyskinesia. *Am.J.Med.Genet.B Neuropsychiatr.Genet.*, 141 195-197.
174. Rietschel,M., Nothen,M.M., Lannfelt,L., Sokoloff,P., Schwartz,J.C., Lanczik,M., Fritze,J., Cichon,S., Fimmers,R., Korner,J. e . 1993. A serine to glycine substitution at position 9 in the extracellular N-terminal part of the dopamine D3 receptor protein: no role in the genetic predisposition to bipolar affective disorder. *Psychiatry Res.*, 46 253-259.
175. Rietschel,M., Naber,D., Oberlander,H., Holzbach,R., Fimmers,R., Eggermann,K., Moller,H.J., Propping,P. e Nothen,M.M. 1996. Efficacy and side-effects of clozapine: testing for association with allelic variation in the dopamine D4 receptor gene. *Neuropsychopharmacology*, 15 491-496.
176. Rietschel,M., Naber,D., Fimmers,R., Moller,H.J., Propping,P. e Nothen,M.M. 1997. Efficacy and side-effects of clozapine not associated with variation in the 5-HT2C receptor. *Neuroreport*, 8 1999-2003.
177. Riley,B. e Kendler,K.S. 2006. Molecular genetic studies of schizophrenia. *Eur.J.Hum.Genet.*, 14 669-680.
178. Robinson,D., Woerner,M.G., Alvir,J.M., Bilder,R., Goldman,R., Geisler,S., Koreen,A., Sheitman,B., Chakos,M., Mayerhoff,D. e Lieberman,J.A. 1999. Predictors of relapse following response from a first episode of schizophrenia or schizoaffective disorder. *Arch.Gen.Psychiatry*, 56 241-247.
179. Ross,C.A., Margolis,R.L., Reading,S.A., Pletnikov,M. e Coyle,J.T. 2006. Neurobiology of schizophrenia. *Neuron*, 52 139-153.
180. Ruano,G., Goethe,J.W., Caley,C., Woolley,S., Holford,T.R., Kocherla,M., Windemuth,A. e de,L.J. 2007. Physiogenomic comparison of weight profiles of olanzapine- and risperidone-treated patients. *Mol.Psychiatry*, 12 474-482.
181. Rupp,A. e Keith,S.J. 1993. The costs of schizophrenia. Assessing the burden. *Psychiatr.Clin.North Am.*, 16 413-423.

182. Rutter,M., Silberg,J., O'Connor,T. e Simonoff,E. 1999. Genetics and child psychiatry: I Advances in quantitative and molecular genetics. *J.Child Psychol.Psychiatry*, 40 3-18.
183. Sedvall,G. e Farde,L. 1995. Chemical brain anatomy in schizophrenia. *Lancet*, 346 743-749.
184. Seeman,P. e Lee,T. 1975. Antipsychotic drugs: direct correlation between clinical potency and presynaptic action on dopamine neurons. *Science*, 188 1217-1219.
185. Segman,R.H., Heresco-Levy,U., Finkel,B., Inbar,R., Neeman,T., Schlafman,M., Dorevitch,A., Yakir,A., Lerner,A., Goltser,T., Shelevoy,A. e Lerer,B. 2000. Association between the serotonin 2C receptor gene and tardive dyskinesia in chronic schizophrenia: additive contribution of 5-HT_{2C}ser and DRD₃gly alleles to susceptibility. *Psychopharmacology (Berl)*, 152 408-413.
186. Segman,R.H., Heresco-Levy,U., Finkel,B., Goltser,T., Shalem,R., Schlafman,M., Dorevitch,A., Yakir,A., Greenberg,D., Lerner,A. e Lerer,B. 2001. Association between the serotonin 2A receptor gene and tardive dyskinesia in chronic schizophrenia. *Mol.Psychiatry*, 6 225-229.
187. Segman,R.H., Shapira,Y., Modai,I., Hamdan,A., Zislin,J., Heresco-Levy,U., Kanyas,K., Hirschmann,S., Karni,O., Finkel,B., Schlafman,M., Lerner,A., Shapira,B., Macciardi,F. e Lerer,B. 2002. Angiotensin converting enzyme gene insertion/deletion polymorphism: case-control association studies in schizophrenia, major affective disorder, and tardive dyskinesia and a family-based association study in schizophrenia. *Am.J.Med.Genet.*, 114 310-314.
188. Segman,R.H., Goltser,T., Heresco-Levy,U., Finkel,B., Shalem,R., Schlafman,M., Yakir,A., Greenberg,D., Strous,R., Lerner,A., Shelevoy,A. e Lerer,B. 2003. Association of dopaminergic and serotonergic genes with tardive dyskinesia in patients with chronic schizophrenia. *Pharmacogenomics.J.*, 3 277-283.
189. Shaikh,S., Collier,D., Kerwin,R.W., Pilowsky,L.S., Gill,M., Xu,W.M. e Thornton,A. 1993. Dopamine D4 receptor subtypes and response to clozapine. *Lancet*, 341 116
190. Shaikh,S., Collier,D.A., Sham,P.C., Ball,D., Aitchison,K., Vallada,H., Smith,I., Gill,M. e Kerwin,R.W. 1996. Allelic association between a Ser-9-Gly polymorphism in the dopamine D3 receptor gene and schizophrenia. *Hum.Genet.*, 97 714-719.
191. Sherrington,R., Brynjolfsson,J., Petursson,H., Potter,M., Dudleston,K., Barraclough,B., Wasmuth,J., Dobbs,M. e Gurling,H. 1988. Localization of a susceptibility locus for schizophrenia on chromosome 5. *Nature*, 336 164-167.
192. Shinkai,T., Ohmori,O., Matsumoto,C., Hori,H., Kennedy,J.L. e Nakamura,J. 2004. Genetic association analysis of neuronal nitric oxide synthase gene polymorphism with tardive dyskinesia. *Neuromolecular.Med.*, 5 163-170.

193. Shinkai,T., Muller,D.J., De,L., V, Shaikh,S., Matsumoto,C., Hwang,R., King,N., Trakalo,J., Potapova,N., Zai,G., Hori,H., Ohmori,O., Meltzer,H.Y., Nakamura,J. e Kennedy,J.L. 2006. Genetic association analysis of the glutathione peroxidase (GPX1) gene polymorphism (Pro197Leu) with tardive dyskinesia. *Psychiatry Res.*, 141 123-128.
194. Shorter, E., 1997. *A History of Psychiatry*. Wiley, New York, pp. 26– 28, 93– 99, 196– 200.
195. Sklar,P., Pato,M.T., Kirby,A., Petryshen,T.L., Medeiros,H., Carvalho,C., Macedo,A., Dourado,A., Coelho,I., Valente,J., Soares,M.J., Ferreira,C.P., Lei,M., Verner,A., Hudson,T.J., Morley,C.P., Kennedy,J.L., Azevedo,M.H., Lander,E., Daly,M.J. e Pato,C.N. 2004. Genome-wide scan in Portuguese Island families identifies 5q31-5q35 as a susceptibility locus for schizophrenia and psychosis. *Mol.Psychiatry*, 9 213-218.
196. Sodhi,M.S., Arranz,M.J., Curtis,D., Ball,D.M., Sham,P., Roberts,G.W., Price,J., Collier,D.A. e Kerwin,R.W. 1995. Association between clozapine response and allelic variation in the 5-HT_{2C} receptor gene. *Neuroreport*, 7 169-172.
197. Souza,R.P., Romano-Silva,M.A., Lieberman,J.A., Meltzer,H.Y., Wong,A.H. e Kennedy,J.L. 2008. Association study of GSK3 gene polymorphisms with schizophrenia and clozapine response. *Psychopharmacology (Berl)*,
198. Soyka,M., Albus,M., Kathmann,N., Finelli,A., Hofstetter,S., Holzbach,R., Immler,B. e Sand,P. 1993. Prevalence of alcohol and drug abuse in schizophrenic inpatients. *Eur.Arch.Psychiatry Clin.Neurosci.*, 242 362-372.
199. Srivastava,V., Varma,P.G., Prasad,S., Semwal,P., Nimgaonkar,V.L., Lerer,B., Deshpande,S.N. e BK,T. 2006. Genetic susceptibility to tardive dyskinesia among schizophrenia subjects: IV. Role of dopaminergic pathway gene polymorphisms. *Pharmacogenet.Genomics*, 16 111-117.
200. St Clair D., Xu,M., Wang,P., Yu,Y., Fang,Y., Zhang,F., Zheng,X., Gu,N., Feng,G., Sham,P. e He,L. 2005. Rates of adult schizophrenia following prenatal exposure to the Chinese famine of 1959-1961. *JAMA*, 294 557-562.
201. Stefansson,H., Sigurdsson,E., Steinthorsdottir,V., Bjornsdottir,S., Sigmundsson,T., Ghosh,S., Brynjolfsson,J., Gunnarsdottir,S., Ivarsson,O., Chou,T.T., Hjaltason,O., Birgisdottir,B., Jonsson,H., Gudnadottir,V.G., Gudmundsdottir,E., Bjornsson,A., Ingvarsson,B., Ingason,A., Sigfusson,S., Hardardottir,H., Harvey,R.P., Lai,D., Zhou,M., Brunner,D., Mutel,V., Gonzalo,A., Lemke,G., Sainz,J., Johannesson,G., Andresson,T., Gudbjartsson,D., Manolescu,A., Frigge,M.L., Gurney,M.E., Kong,A., Gulcher,J.R., Petursson,H. e Stefansson,K. 2002. Neuregulin 1 and susceptibility to schizophrenia. *Am.J.Hum.Genet.*, 71 877-892.
202. Straub,R.E., Jiang,Y., MacLean,C.J., Ma,Y., Webb,B.T., Myakishev,M.V., Harris-Kerr,C., Wormley,B., Sadek,H., Kadambi,B., Cesare,A.J., Gibberman,A., Wang,X., O'Neill,F.A., Walsh,D. e Kendler,K.S. 2002. Genetic variation in the 6p22.3 gene

DTNBP1, the human ortholog of the mouse dysbindin gene, is associated with schizophrenia. *Am.J.Hum.Genet.*, 71 337-348.

203. Strous,R.D., Greenbaum,L., Kanyas,K., Merbl,Y., Horowitz,A., Karni,O., Viglin,D., Olender,T., Deshpande,S.N., Lancet,D., Ben-Asher,E. e Lerer,B. 2007. Association of the dopamine receptor interacting protein gene, NEF3, with early response to antipsychotic medication. *Int.J.Neuropsychopharmacol.*, 10 321-333.

204. Suarez,B.K., Duan,J., Sanders,A.R., Hinrichs,A.L., Jin,C.H., Hou,C., Buccola,N.G., Hale,N., Weilbaecher,A.N., Nertney,D.A., Olincy,A., Green,S., Schaffer,A.W., Smith,C.J., Hannah,D.E., Rice,J.P., Cox,N.J., Martinez,M., Mowry,B.J., Amin,F., Silverman,J.M., Black,D.W., Byerley,W.F., Crowe,R.R., Freedman,R., Cloninger,C.R., Levinson,D.F. e Gejman,P.V. 2006. Genomewide linkage scan of 409 European-ancestry and African American families with schizophrenia: suggestive evidence of linkage at 8p23.3-p21.2 and 11p13.1-q14.1 in the combined sample. *Am.J.Hum.Genet.*, 78 315-333.

205. Susser,E., Neugebauer,R., Hoek,H.W., Brown,A.S., Lin,S., Labovitz,D. e Gorman,J.M. 1996. Schizophrenia after prenatal famine. Further evidence. *Arch.Gen.Psychiatry*, 53 25-31.

206. Szekeres,G., Keri,S., Juhasz,A., Rimanoczy,A., Szendi,I., Czimmer,C. e Janka,Z. 2004. Role of dopamine D3 receptor (DRD3) and dopamine transporter (DAT) polymorphism in cognitive dysfunctions and therapeutic response to atypical antipsychotics in patients with schizophrenia. *Am.J.Med.Genet.B Neuropsychiatr.Genet.*, 124 1-5.

207. Tandon,R. 2007. Antipsychotic treatment of schizophrenia: two steps forward, one step back. *Curr.Psychiatry Rep.*, 9 263-264.

208. Tandon,R., Belmaker,R.H., Gattaz,W.F., Lopez-Ibor,J.J., Jr., Okasha,A., Singh,B., Stein,D.J., Olie,J.P., Fleischhacker,W.W. e Moeller,H.J. 2008. World Psychiatric Association Pharmacopsychiatry Section statement on comparative effectiveness of antipsychotics in the treatment of schizophrenia. *Schizophr.Res.*, 100 20-38.

209. Theisen,F.M., Gebhardt,S., Haberhausen,M., Heinzl-Gutenbrunner,M., Wehmeier,P.M., Krieg,J.C., Kuhnau,W., Schmidtke,J., Remschmidt,H. e Hebebrand,J. 2005. Clozapine-induced weight gain: a study in monozygotic twins and same-sex sib pairs. *Psychiatr.Genet.*, 15 285-289.

210. Thelma,B.K., Tiwari,A.K., Deshpande,S.N., Lerer,B. e Nimgaonkar,V.L. 2007. Genetic susceptibility to Tardive Dyskinesia in chronic schizophrenia subjects: role of oxidative stress pathway genes. *Schizophr.Res.*, 92 278-279.

211. Tiwari,A.K., Deshpande,S.N., Rao,A.R., Bhatia,T., Lerer,B., Nimgaonkar,V.L. e Thelma,B.K. 2005. Genetic susceptibility to tardive dyskinesia in chronic schizophrenia subjects: III. Lack of association of CYP3A4 and CYP2D6 gene polymorphisms. *Schizophr.Res.*, 75 21-26.

212. Torrey,E.F., Bowler,A.E. e Clark,K. 1997. Urban birth and residence as risk factors for psychoses: an analysis of 1880 data. *Schizophr.Res.*, 25 169-176.
213. Tsai,L.H. e Gleeson,J.G. 2005. Nucleokinesis in neuronal migration. *Neuron*, 46 383-388.
214. Tsai,S.J., Hong,C.J., Yu,Y.W., Lin,C.H., Song,H.L., Lai,H.C. e Yang,K.H. 2000. Association study of a functional serotonin transporter gene polymorphism with schizophrenia, psychopathology and clozapine response. *Schizophr.Res.*, 44 177-181.
215. Tsai,S.J., Wang,Y.C., Yu Younger,W.Y., Lin,C.H., Yang,K.H. e Hong,C.J. 2001. Association analysis of polymorphism in the promoter region of the alpha2a-adrenoceptor gene with schizophrenia and clozapine response. *Schizophr.Res.*, 49 53-58.
216. Tsai,S.J., Hong,C.J., Yu,Y.W., Lin,C.H. e Liu,L.L. 2003. No association of tumor necrosis factor alpha gene polymorphisms with schizophrenia or response to clozapine. *Schizophr.Res.*, 65 27-32.
217. Tsai,S.J., Yu,Y.W., Lin,C.H., Wang,Y.C., Chen,J.Y. e Hong,C.J. 2004. Association study of adrenergic beta3 receptor (Trp64Arg) and G-protein beta3 subunit gene (C825T) polymorphisms and weight change during clozapine treatment. *Neuropsychobiology*, 50 37-40.
218. Tsuang, M.T., Vandermey, R., 1980. *Genes and the Mind*. Oxford Univ. Press, Oxford, p. 71.
219. Tuunainen,A., Wahlbeck,K. e Gilbody,S. 2002. Newer atypical antipsychotic medication in comparison to clozapine: a systematic review of randomized trials. *Schizophr.Res.*, 56 1-10.
220. Wehmeier,P.M., Gebhardt,S., Schmidtke,J., Remschmidt,H., Hebebrand,J. e Theisen,F.M. 2005. Clozapine: weight gain in a pair of monozygotic twins concordant for schizophrenia and mild mental retardation. *Psychiatry Res.*, 133 273-276.
221. Weinberger,D.R. 1995. From neuropathology to neurodevelopment. *Lancet*, 346 552-557.
222. Weinberger,D.R., Egan,M.F., Bertolino,A., Callicott,J.H., Mattay,V.S., Lipska,B.K., Berman,K.F. e Goldberg,T.E. 2001. Prefrontal neurons and the genetics of schizophrenia. *Biol.Psychiatry*, 50 825-844.
223. Woerner,M.G., Kane,J.M., Lieberman,J.A., Alvir,J., Bergmann,K.J., Borenstein,M., Schooler,N.R., Mukherjee,S., Rotrosen,J., Rubinstein,M. e . 1991. The prevalence of tardive dyskinesia. *J.Clin.Psychopharmacol.*, 11 34-42.
224. Xu,M.Q., Xing,Q.H., Zheng,Y.L., Li,S., Gao,J.J., He,G., Guo,T.W., Feng,G.Y., Xu,F. e He,L. 2007. Association of AKT1 gene polymorphisms with risk of

schizophrenia and with response to antipsychotics in the Chinese population. *J.Clin.Psychiatry*, 68 1358-1367.

225. Yamanouchi,Y., Iwata,N., Suzuki,T., Kitajima,T., Ikeda,M. e Ozaki,N. 2003. Effect of DRD2, 5-HT2A, and COMT genes on antipsychotic response to risperidone. *Pharmacogenomics.J.*, 3 356-361.

226. Yasui-Furukori,N., Saito,M., Nakagami,T., Kaneda,A., Tateishi,T. e Kaneko,S. 2006. Association between multidrug resistance 1 (MDR1) gene polymorphisms and therapeutic response to bromperidol in schizophrenic patients: a preliminary study. *Prog.Neuropsychopharmacol.Biol.Psychiatry*, 30 286-291.

227. Youssef,H., Lyster,G. e Youssef,F. 1989. Familial psychosis and vulnerability to tardive dyskinesia. *Int.Clin.Psychopharmacol.*, 4 323-328.

228. Yu,Y.W., Tsai,S.J., Lin,C.H., Hsu,C.P., Yang,K.H. e Hong,C.J. 1999. Serotonin-6 receptor variant (C267T) and clinical response to clozapine. *Neuroreport*, 10 1231-1233.

229. Zai,G., Muller,D.J., Volavka,J., Czobor,P., Lieberman,J.A., Meltzer,H.Y. e Kennedy,J.L. 2006. Family and case-control association study of the tumor necrosis factor-alpha (TNF-alpha) gene with schizophrenia and response to antipsychotic medication. *Psychopharmacology (Berl)*, 188 171-182.

230. Zerbin-Rudin, E., 1967. Endogene Psychosen. In: Becker, P.E. (Ed.), *Humangenetik ein Kurzes Handbuch, Band 2*. Thieme, Stuttgart.

231. Zhang,Z.J., Yao,Z.J., Mou,X.D., Chen,J.F., Zhu,R.X., Liu,W., Zhang,X.R., Sun,J. e Hou,G. 2003. [Association of -2548G/A functional polymorphism in the promoter region of leptin gene with antipsychotic agent-induced weight gain]. *Zhonghua Yi.Xue.Za Zhi.*, 83 2119-2123.

Livros Grátis

(<http://www.livrosgratis.com.br>)

Milhares de Livros para Download:

[Baixar livros de Administração](#)

[Baixar livros de Agronomia](#)

[Baixar livros de Arquitetura](#)

[Baixar livros de Artes](#)

[Baixar livros de Astronomia](#)

[Baixar livros de Biologia Geral](#)

[Baixar livros de Ciência da Computação](#)

[Baixar livros de Ciência da Informação](#)

[Baixar livros de Ciência Política](#)

[Baixar livros de Ciências da Saúde](#)

[Baixar livros de Comunicação](#)

[Baixar livros do Conselho Nacional de Educação - CNE](#)

[Baixar livros de Defesa civil](#)

[Baixar livros de Direito](#)

[Baixar livros de Direitos humanos](#)

[Baixar livros de Economia](#)

[Baixar livros de Economia Doméstica](#)

[Baixar livros de Educação](#)

[Baixar livros de Educação - Trânsito](#)

[Baixar livros de Educação Física](#)

[Baixar livros de Engenharia Aeroespacial](#)

[Baixar livros de Farmácia](#)

[Baixar livros de Filosofia](#)

[Baixar livros de Física](#)

[Baixar livros de Geociências](#)

[Baixar livros de Geografia](#)

[Baixar livros de História](#)

[Baixar livros de Línguas](#)

[Baixar livros de Literatura](#)
[Baixar livros de Literatura de Cordel](#)
[Baixar livros de Literatura Infantil](#)
[Baixar livros de Matemática](#)
[Baixar livros de Medicina](#)
[Baixar livros de Medicina Veterinária](#)
[Baixar livros de Meio Ambiente](#)
[Baixar livros de Meteorologia](#)
[Baixar Monografias e TCC](#)
[Baixar livros Multidisciplinar](#)
[Baixar livros de Música](#)
[Baixar livros de Psicologia](#)
[Baixar livros de Química](#)
[Baixar livros de Saúde Coletiva](#)
[Baixar livros de Serviço Social](#)
[Baixar livros de Sociologia](#)
[Baixar livros de Teologia](#)
[Baixar livros de Trabalho](#)
[Baixar livros de Turismo](#)