



Universidade Federal do Rio Grande do Sul  
Instituto de Biociências  
Departamento de Zoologia  
Programa de Pós-Graduação em Biologia Animal

CLAITON MARTINS FERREIRA

ESTRUTURA POPULACIONAL, DIVERSIDADE GENÉTICA, ÁREA DE DISTRIBUIÇÃO E  
CONSERVAÇÃO DO CARDEAL-AMARELO – *GUBERNATRIX CRISTATA* (VIEILLOT, 1817)  
(AVES, PASSERIFORMES, EMBERIZIDAE).

Tese apresentada ao Programa de Pós-Graduação em  
Biologia Animal, Instituto de Biociências da Universidade  
Federal do Rio Grande do Sul, como requisito parcial à  
obtenção do título de Doutor em Biologia Animal.

Área de Concentração: Biologia Comparada

Orientador: Prof. Dr. Thales Renato Ochotorena de Freitas

UNIVERSIDADE FEDERAL DO RIO GRANDE DO SUL  
PORTO ALEGRE  
2010

# **Livros Grátis**

<http://www.livrosgratis.com.br>

Milhares de livros grátis para download.

Tese intitulada “ESTRUTURA POPULACIONAL, DIVERSIDADE GENÉTICA, ÁREA DE DISTRIBUIÇÃO E CONSERVAÇÃO DO CARDEAL-AMARELO - *GUBERNATRIX CRISTATA* (VIEILLOT, 1817) (AVES, PASSERIFORMES, EMBERIZIDAE)” apresentada por **Claiton Martins Ferreira** como parte dos requisitos para obtenção do grau de Doutor em Biologia Animal, Área de Concentração Biologia Comparada, aprovada em 12/08/2010 pela Comissão Examinadora:

Dr. Thales Renato Ochotorena de Freitas  
Orientador

Dra. Vera Lúcia Valente Gaiesty

Dra. Márcio Borges Martins

Dr. Jorge Luiz Berger Albuquerque

Homologada em                      conforme ata N.º                      pela Comissão Coordenadora.

Dra. Helena Piccoli Romanowski  
Coordenadora

## SUMÁRIO

DEDICATÓRIA.....	iii
AGRADECIMENTOS.....	iv
ACKNOWLEDGEMENTS.....	vii
RESUMO.....	x
ABSTRACT .....	xii
CAPÍTULO UM .....	1
1. INTRODUÇÃO .....	1
1.1.Estruturação da Tese e Objetivos.....	2
1.2.O Cardeal-Amarelo .....	3
1.3.As Ameaças .....	5
1.4.Métodos Gerais .....	6
1.5.Resultados Gerais .....	9
1. INTRODUCTION.....	7
1.1.General Introduction and Thesis Aims .....	12
1.2.The bird .....	12
1.3.The Threats .....	14
1.4.General Methods .....	15
1.5.General Results .....	18
CAPÍTULO DOIS.....	22
2. Isolation and characterization of 10 microsatellite loci in the Yellow Cardinal <i>Gubernatrix cristata</i> En.....	23
Abstract.....	23
Introduction .....	23
Material and Methods.....	24
Acknowledgements .....	25
References .....	25
CAPÍTULO TRÊS.....	29
3. Lack of population structure in the Yellow Cardinal <i>Gubernatrix cristata</i> En and implications to its conservation .....	30

Abstract.....	30
Introduction .....	31
Methods.....	32
Results.....	36
Discussion .....	41
Conclusions .....	43
Acknowledgements .....	44
References .....	44
CAPÍTULO QUATRO.....	47
4. Potential Distribution Model to Yellow Cardinal <i>Gubernatrix cristata</i> : how much effort we should put to preserve this Endangered bird? .....	48
Abstract.....	48
Introduction .....	49
Methods.....	50
Results.....	55
Discussion .....	61
Conclusion.....	64
Acknowledgements .....	64
References .....	65
CAPÍTULO CINCO .....	68
5. CONCLUSÕES E RECOMENDAÇÕES PARA A CONSERVAÇÃO DA ESPÉCIE.....	68
5.1.Viabilidade do cardeal-amarelo no RS, Uruguai e Argentina.....	69
5.2.Recomendações para a conservação: um primeiro passo para a construção de um plano internacional para a conservação do cardeal-amarelo no Sul da América do Sul .....	71
5. CONCLUSIONS AND CONSERVATION RECOMMENDATIONS .....	73
5.1. Viability of yellow cardinal in RS, Uruguay and Argentina .....	73
5.2. Conservation Recommendations: A first step in building a regional action plan for the conservation of yellow cardinal in Southern South America .....	75
6. REFERENCIAS .....	76
7. ANEXOS.....	81

## DEDICATÓRIA

*Para todas as aves que necessitam urgente de planos de conservação, especialmente o cardeal-amarelo.*

*Para a minha amada esposa, Regina. Verdadeira companheira e amiga, que me deu muito de seu apoio em todos os momentos dessa caminhada. Te amo.*

## AGRADECIMENTOS

Na realização de um estudo desta dimensão muitas pessoas acabam invariavelmente sendo envolvidas, direta ou indiretamente, do começo ao fim do trabalho. É quase impossível nomear a todos sem cometer injustiças. Tenho inúmeras pessoas a agradecer. Mas algumas delas se destacaram. Seja pela sua presença constante, seja para ajudar no momento crítico, seja dando um ombro amigo nos momentos mais estressantes (que não foram poucos).

Agradeço à Fundação O Boticário de Proteção à Natureza pelo apoio financeiro. Essa concessão tornou esta tese possível.

Ao CNPq, pelo apoio financeiro através de bolsa de doutorado e taxa de bancada, a qual foi realmente valiosa durante o curso deste doutorado.

À CAPES pelo suporte financeiro através da bolsa sanduíche e passagens aéreas POA - TO - POA. Ter feito este estágio no Canadá foi uma das melhores experiências que eu tive durante a minha formação profissional como cientista.

Ao ICMBio e IBAMA pela emissão de todas as licenças e apoio durante a elaboração do Primeiro Plano de Ação do Cardeal-Amarelo.

Agradeço a todos os professores do PPG Biologia Animal, muito bem representados por sua coordenadora, profa. Helena Romanowski, por todo o conhecimento transmitido e saudáveis discussões realizadas ao longo da realização dessa tese. Todo o apoio e vivência que lá tive carregou em minha experiência profissional.

Agradeço aos membros da banca pelo aceite em revisar minha tese em curtíssimo espaço de tempo, especialmente à Dra. Vera Gaiety e ao Dr. Jorge Albuquerque, que há 20 anos acompanham minha trajetória profissional sempre com gentis palavras de incentivo.

Agradeço às professoras do Departamento de Zoologia, Clarice Fialho e Laura Verrastro, por todo o apoio recebido durante minha estada nesse departamento. De amáveis colegas se tornaram queridas amigas.

Agradeço ao Prof. Dr. Eduardo Eizirik da PUCRS pela providencial carta avaliando positivamente meu projeto a ser realizado no Canadá.

Agradeço ao meu orientador Thales de Freitas, por acreditar em mim e me permitir fazer tudo o que eu considere importante para o doutorado.

Agradeço imensamente a todos os colegas do laboratório de citogenética e evolução pela amizade e a parceria sempre presente. Um agradecimento especial a Carla Lopes, Paula Rorato, Gabriela Fernandes, Gislene Lopes e Darlise Santos. Vocês todas são definitivamente tanto o motor quanto o cérebro da área molecular do laboratório.

Agradeço ao meu co-orientador do Royal Ontario Museum, o Prof Dr. Allan J. Baker por me dar a oportunidade de fazer um estágio em seu laboratório de primeira linha, o qual acrescentou muito à minha formação profissional.

Agradeço aos meus colegas do laboratório de biologia molecular de aves do Royal Ontario Museum, especialmente Oliver Haddrath por sua boa vontade e paciência para ensinar alguém, sem uma grande experiência molecular (para não dizer quase nenhuma); a Erica Tavares, Rosemary Gomes, Pasan Dinasayake, Yvonne Verkuil e Jacqueline Muller pelas incontáveis boas dicas, hospitalidade e amizade sempre presentes.

À minha querida amiga Prof Dra. Cristina Myiaki do laboratório de genética molecular de aves da USP, pela calorosa acolhida e oportunidade de treinamento em seu laboratório, com discussões importantes para minha pesquisa de tese.

Aos meus colegas do laboratório de genética molecular de aves da USP, pelas dicas, paciência e amizade durante a minha estadia lá, especialmente o Fernando, que me ajudou muito.

À Profa. Anete Pereira do CBMEG/UNICAMP pelo aprendizado na criação da biblioteca de microssatélites do cardeal-amarelo e a Dra. Melissa O. S. Garcia pela genotipagem dos clones.

Ao Sr. Fernando Leon Lucas e sua esposa, incansáveis e amorosas pessoas, que não mediram esforços para deixar-nos à vontade em todas as vezes que fomos à cidade de Pinheiro Machado e permanecemos em sua casa.

Ao meu amigo Mestre Glayson Bencke da FZB pela sua palestra inspiradora lá em 2004. Aquela conversa me incentivou a trabalhar com esta espécie e tentar contribuir para sua preservação.

Ao Rodrigo Damiani por sua parceria durante o trabalho de campo e boa



vontade nas viagens sozinha ao Espinilho durante o projeto de monitoramento do cardeal-amarelo. Ele acreditou que esta ave não era um fantasma e que um dia iríamos encontrá-la na natureza. O que de fato acabou acontecendo.

Agradeço a Elmo Cardoso, secretário do PPGGBM que, embora não estando diretamente vinculado ao PPGBAN, nunca se furtou de me esclarecer alguma questão acadêmica, burocrática, prática ou simplesmente para bater um papo.

Agradeço ao meu querido amigo Prof. Dr. Márcio Efe, parceiro de ONG, de projetos, de discussões e um grande incentivador de minha carreira profissional como ornitólogo.

Agradeço a todos os meus amigos mais próximos, principalmente o Adriano e o Enio por continuarem acreditando em mim e sendo leais amigos, mesmo durante as freqüentes ausências durante a realização do doutorado.

Agradeço ao meu filho Cesar Schunemann por continuar fazendo parte da minha vida e sempre compreender as minhas freqüentes ausências nos últimos tempos. É um filho que conheci já adulto e que não pude ensinar tudo o que eu gostaria de ter ensinado durante seu desenvolvimento. Mas ele teve uma excelente mãe. E o resultado é que acabei aprendendo muito com ele. E, principalmente, aprendi a admirá-lo.

Agradeço a todas as outras pessoas que eu não estou colocando aqui seus nomes. Como afirmei antes, há muitas pessoas que eu tenho que agradecer e não dá para nominar todo mundo. Sou profundamente grato pela sua ajuda.

Ao cardeal-amarelo o qual generosamente doou algumas penas e gotas de sangue, embora não completamente de acordo. Sem estas preciosas amostras essa tese não existiria.

E agradeço à minha esposa Regina. Ela é a melhor mulher que eu conheci durante minha vida inteira. Ela é paciente (sendo minha mulher diz muito sobre sua paciência), querida, delicada, generosa, apoiadora e sempre uma agradável presença em minha vida. Querida, não há palavras para dizer o quanto sou grato a você. Tudo o que você fez para mim durante esta cansativa experiência eu gostaria de poder lhe dar de volta em dobro. Mas você é aquele tipo de pessoa que é rara hoje em dia. Uma mulher verdadeiramente nobre e elegante, em todos os sentidos. Obrigado por todo o apoio antes, durante e após essa tese. Obrigado por compartilhar sua vida comigo.

## ACKNOWLEDGEMENTS

In conducting a study of this size many people invariably end up being involved, directly or indirectly, from beginning to the ending. It is almost impossible to name everybody without committing injustice. I have so many people to thank for. But some of them stood out. Whether by their constant presence, whether for help at the critical moment, either by giving a friendly shoulder at the most stressful times (which were not few).

I thank my advisor Thales de Freitas for believing me and let me do whatever I deemed important for the doctorate.

I greatly appreciate all the colleagues of the laboratory of cytogenetics and evolution for the friendship and partnership always present. Special thanks to Carla Lopes, Paula Rorato, Gabriela Fernandez, Gislene Lopes, and Darlise Santos. You all are definitely both the engine and brain of the laboratory molecular area.

I thank my co-advisor from Royal Ontario Museum, Prof. Dr. Allan J. Baker for giving me the opportunity to do an internship at his first class lab, which has added a lot to my professional training.

I thank my colleagues of avian molecular biology laboratory in the Royal Ontario Museum, especially Oliver Haddrath for his goodwill and patience in teaching someone with not such a great molecular experience; to Erica Tavares, Rosemary Gibson, Pasan Dinasayake, Yvonne Verkuil, and Jacqueline Muller for the countless good advices, hospitality and friendship always present.

To my dear friend Prof. Dr. Cristina Myiaki of avian molecular genetics laboratory, USP, for the warm welcome and training opportunity at her lab, with valuable discussions to my thesis research.

To my colleagues of avian molecular genetics laboratory, USP, for the good advices, patience and friendship during my stay there, especially to Fernando.

To Prof. Anete Pereira from CBMEG / UNICAMP for the opportunity of attending an excellent course conducted by her which has resulted in the creation of Yellow Cardinal microsatellite library.

To Dr. Melissa O. S. Garcia who kindly offered herself to genotype my clones. This assistance yielded a publication in Molecular Ecology Resources.

To Mr. Fernando Leon Lucas and his wife, tireless and loving people, who did not measure efforts to make us comfortable at all times we went to Pinheiro Machado City and stayed at their home.

To my friend M.Sc. Glayson Bencke from FZB by his inspirational lecture back in 2004. That talk encouraged me to work with this species and seek to contribute to its preservation.

To Rodrigo Damiani by his partnership during fieldwork and good will in the lonely trips to Espinilho monitoring the Yellow Cardinal. He had believed that this bird was not a ghost and we would find it in nature someday. What eventually happened indeed.

Thanks to Elmo Cardoso, Secretary of PPGGBM, while not directly tied to PPGBAN, have never refused to clarify any academic questions, bureaucratic, practice, or simply to chat.

Thanks to my dear friend Prof. Dr. Márcio Efe, NGO fellow, partner of different projects, and someone who always gave a great encouragement to my career as an ornithologist.

To Fundação O Boticário de Proteção à Natureza for its financial support. That grant made this thesis possible.

To CNPq, for financial support through doctoral scholarship and fee bench, which was really valuable during the course of this doctorate.

To CAPES for financial support through the sandwich scholarship and airline tickets POA – TO – POA. Having done this internship in Canada was one of the best experiences I had made to improve my professional career as scientist.

To ICMBIO and IBAMA for licenses and support provided during the preparation of the first Yellow Cardinal action plan.

To everybody else whom I am not putting their names here. As I stated before, there are too many people I have to thank. I am deeply grateful for their help.

To Yellow Cardinal which generously gave away some feathers and blood drops, even though not quite agreed. Without those precious samples this thesis wouldn't exist at all.

And to my wife Regina. She is the best woman I ever met during my whole life. She is patient (being my wife tells a lot about her patience), kind, supportive, delicate,

generous, and always a good presence in my life. Dear, there isn't words to tell you how much grateful I am to you. Everything what you have done to me during this exhilarated experience I wish to be able give you back at double. But you are that sort of person which is rare nowadays. A truly noble and elegant woman, in all senses. Thanks for all the support before, during and after this thesis. Thank you for sharing your life with me.

## RESUMO

### **Estrutura populacional, diversidade genética, área de distribuição e conservação do cardeal-amarelo En – *Gubernatrix cristata* (Vieillot, 1817) (Aves, Passeriformes)(Emberizidae)**

*Gubernatrix cristata*, o cardeal-amarelo, é uma espécie de ave rara do Pampa. Ele tem uma distribuição geográfica restrita ao sul da América do Sul (Uruguai, Argentina e sul do Brasil) e é exclusivo deste bioma. Quatorze viagens de campo foram feitas de Abril de 2006 a março de 2009, em uma tentativa de encontrar indivíduos e coletar amostras para análise genética. Desenvolvemos dez marcadores microsatélites isolados a partir desta espécie e caracterizamos a sua variabilidade alélica. O número de alelos observados em cada locus variou de 4 a 14, com uma média de 7,5 alelos por locus. Os microsatélites mostraram-se úteis em revelar os níveis de diversidade de *G. cristata* e, portanto, podem ser usados para explorar a estrutura genética das populações dispersas ao longo de sua distribuição geográfica atual. Um total de 72 amostras de cardeal-amarelo foi usado neste estudo, 59 foram de amostras contemporâneas e 13 amostras de espécimes de museu. Nós acessamos a diversidade genética da espécie através dos dez loci polimórficos de microsatélites nucleares do *G. cristata* desenvolvidos nesse estudo e através de um fragmento do gene ND2 do DNA mitocondrial. Encontramos apenas três haplótipos com diversidade nucleotídica e haplotípica respectivamente igual a  $\pi = 0,00277$  e  $Hd = 0,6219$ . O  $F_{st}$  (0,00340) e  $Nm$  (73,18) mostraram uma fraca estruturação com pouca diferenciação genética. A estatística  $F_s$  de  $F_u$  foi significativamente diferente de zero (2,248) e  $D$  de Tajima foi positivo (2,17506). Ambos os testes mostraram um indicativo de evidência provável de uma diminuição no tamanho da população e/ou seleção equilibradora. As análises usando o "no admixture model" e um grande burn-in (500000) não resultaram em clustering dos indivíduos. Devido a esse resultado nenhuma atribuição individual a uma área geográfica foi possível. O presente estudo auxilia na compreensão das necessidades de conservação do cardeal-amarelo, fornecendo informações sobre a diversidade genética e estruturação populacional ao longo de sua área de distribuição. Embora não tenhamos encontrado estruturação dentro da nossa área de estudo, é preciso mais estudos, analisando a diversidade genética e estruturação populacional em toda a área de distribuição da espécie, e adicionando mais amostras de animais silvestres da população de La Pampa e de Corrientes. O mapa da área de distribuição do cardeal-amarelo parece estar ultrapassado devido à situação cada vez mais crítica que a espécie está enfrentando na natureza. Dada a recente falta de conhecimento sobre sua distribuição e seu estado de conservação em seu habitat natural, decidimos modelar a distribuição potencial da espécie. A principal idéia por trás dessa informação era propor estratégias para sua conservação. Para

melhor alcançar esse objetivo, focamos nossa pesquisa especificamente em a) localizar as populações silvestres desta espécie no Rio Grande do Sul, Uruguai e Argentina, e b) identificar as áreas de ocorrência do cardeal-amarelo que poderiam ser transformadas em áreas protegidas. Todos os registros georeferenciados de *G. cristata* foram extraídos de diferentes fontes: através de trabalho de campo, peles de museu, literatura e bases de dados da Internet. Para modelar a distribuição da espécie, foram selecionadas 20 variáveis ambientais e o aplicativo de modelagem Maxent v. 3.3.2. Mapas de distribuição histórica e atual de *G. cristata* foram criados usando DIVA GIS. O modelo que melhor prediz o índice relativo de adequação ambiental para a espécie é o modelo onde a AUC resultante sobre dados de testes foi 0,868. O resultado do teste de jackknife acerca da importância das variáveis mostrou que a variável ambiental com maior ganho quando usada isoladamente é bio1\_sa\_30s\_cut (temperatura média anual) que, portanto, parece ter as informações mais úteis por si só. A variável ambiental que mais diminui o ganho quando é omitida é alt\_sa\_30s\_cut (altitude) que, portanto, parece ter o máximo de informação que não está presente nas outras variáveis. Se o objetivo de pesquisas futuras do cardeal-amarelo em sua área de distribuição for detectar outras populações, então as áreas preditas em ter elevada adequabilidade relativa no modelo espacial são um bom ponto de partida para a pesquisa mais direcionada. A pressão da caça sobre a espécie é tão grande que o cardeal-amarelo hoje em dia parece ser encontrado apenas em áreas de difícil acesso. Muitas das áreas adequadas para a espécie mostrada pelo modelo podem ainda conter populações e também podem ser usadas como áreas-chave para futuras reintroduções dentro do programa de conservação do cardeal-amarelo.

## ABSTRACT

### **Population Structure, genetic diversity, distribution range and conservation of yellow cardinal En – *Gubernatrix cristata* (Vieillot, 1817) (Aves, Passeriformes)(Emberizidae)**

*Gubernatrix cristata*, the yellow cardinal, is a rare bird species from the Pampas grassland. It has a restricted geographical distribution in southern South America and is unique to this Biome (Uruguay, Argentina, and Southern Brazil). Fourteen field trips were made from April 2006 to March 2009 in an attempt to find individuals and collect samples for genetic analysis. We developed ten microsatellite markers isolated from this species and the characterization of their allele variability. The number of alleles observed for each locus ranged from 4 to 14, with an average of 7.5 alleles per locus. The microsatellites proved to be useful in revealing levels of diversity in *G. cristata*, and thus can be used to explore the genetic structure of scattered populations across its present geographic range. A total of 72 yellow cardinal samples was taken in this study, 59 were from contemporary specimens and 13 samples were from museum specimens. We accessed the species genetic diversity through ten polymorphic nuclear microsatellite loci of *G. cristata* and a ND2 mtDNA fragment. We found only three haplotypes with nucleotide and haplotype diversity equals to  $\pi = 0.00277$  and  $Hd = 0.6219$ . The  $F_{st}$  (0,00340) and  $Nm$  (73,18) shown a weak structuring with little genetic differentiation. Fu's  $F_s$  statistic was significantly different from zero (2,248) and Tajima's  $D$  was positive (2,17506). Both tests showed likely evidence indicating a decrease in population size and/or balancing selection. Analyses using the "no admixture model" and a larger burn-in (500000) yielded no clustering of individuals. Due to this result no individual assignment to any geographical area was possible. The present study assists in understanding the conservation needs for yellow cardinal by providing information on the genetic diversity and population structuring along its distribution range. Although we found no structuring within our study area, further study is needed, examining the genetic diversity and population structuring throughout the species' range, adding more samples from wild animals from La Pampa population and Corrientes. The yellow cardinal distribution range map seems to be outdated because of the increasingly critical situation the species is facing in nature. Given the recent lack of knowledge regarding its distribution and its state of preservation in its natural habitat we decided to model the species potential distribution. The main idea behind this information was to propose strategies for their conservation. To better achieve that target we specifically focused our research on a) locating wild populations of this species in Rio Grande do Sul, Uruguay and Argentina, and b) identifying areas of occurrence of yellow cardinal that could be turned into protected areas. We extracted all georeferenced *G.*

*cristata* records from different sources: field working, museum skins, literature, and internet datasets. To model the species distributions, 20 environmental predictors were selected. To model the species distributions we selected the modeling application Maxent v. 3.3.2. Historical and current distribution maps of *G. cristata* were created using DIVA GIS. The model that best predicts the relative index of environmental suitability for the species is the model where resulting AUC on test data was 0.868. The results of the jackknife test of variable importance showed that the environmental variable with highest gain when used in isolation is bio1\_sa\_30s\_cut (Annual Mean Temperature), which therefore appears to have the most useful information by itself. The environmental variable that decreases the gain the most when it is omitted is alt\_sa\_30s\_cut (Altitude), which therefore appears to have the most information that isn't present in the other variables. If the objective of future surveys of the yellow cardinal in its distribution range is to detect other populations, then areas predicted to have high relative suitability in the spatial model is a good starting point for further targeted survey. The hunting pressure on the species is so great that the yellow cardinal nowadays seems to be only found in areas of difficult access. Many of the areas suitable for the species shown by the model may still contain populations and can also be used as key areas for future reintroductions into the yellow cardinal conservation program.



# CAPÍTULO UM

## Introdução



## 1.1. Estruturação da Tese e Objetivos

A tese está estruturada na forma de artigos. Para facilitar a consulta, foi dividida em cinco capítulos.

Capítulo Um – Introdução: Esse capítulo fornece informações detalhadas sobre a espécie e área de estudo, bem como métodos gerais utilizados nesta pesquisa. A primeira parte do capítulo consiste em uma descrição detalhada do cardeal-amarelo (*Gubernatrix cristata*), fornecendo informações sobre aspectos da taxonomia, morfologia, distribuição geográfica, ecologia, comportamento e uma visão geral do estado de conservação da espécie e as ameaças que enfrenta em seu ambiente natural. A segunda parte fornece informações sobre as ameaças enfrentadas pelo cardeal-amarelo. A terceira parte apresenta os métodos gerais utilizados na pesquisa e a quarta parte apresenta uma síntese dos resultados encontrados.

Capítulo Dois – Primeiro artigo: Esse capítulo apresenta o artigo apresentado durante o exame de qualificação do doutorado e que foi publicado esse ano na revista *Molecular Ecology Resources*.

Capítulo Três – Segundo artigo: Esse capítulo apresenta o artigo que aborda a estrutura populacional e diversidade genética de *G. cristata*. Após a defesa o mesmo será submetido à revista *Animal Conservation*.

Capítulo Quatro – Terceiro artigo: Esse capítulo apresenta o artigo sobre a distribuição potencial modelada para a espécie. Após a defesa o mesmo será submetido à revista *Diversity and Distribution*.

Capítulo Cinco – Conclusão: Esse capítulo sintetiza as conclusões gerais resultantes da tese e recomenda ações para a conservação da espécie.

Referencias – As referencias bibliográficas seguem o estilo da APA (*American Psychological Association*), como sugerido nas orientações ao autor de ambas as revistas - *Animal Conservation* e *Diversity and Distribution*.

Além da introdução e conclusão em Português foi também incluída uma versão em Inglês desses dois capítulos bem como dos agradecimentos. Esse cuidado foi tomado em virtude dos colaboradores canadenses dessa tese.

Nosso principal objetivo foi determinar o status das populações de *Gubernatrix cristata* dentro de sua área de distribuição, especialmente aquelas encontradas no Rio Grande do Sul. Nos focamos em aspectos da biologia reprodutiva, estrutura populacional e fluxo gênico. A principal idéia por trás dessa informação foi propor estratégias para sua conservação. Para melhor atingir

esse alvo, especificamente, buscamos centrar nossa pesquisa em a) localizar as populações selvagens desta espécie no Rio Grande do Sul, Uruguai e Argentina, b) comparar as diferentes populações em termos de variabilidade genética e para verificar o fluxo gênico entre elas, utilizando marcadores moleculares tanto nucleares (microsatélites) quanto mitocondriais (ND2), c) identificação das populações mais importante em termos de variabilidade genética, onde os esforços de conservação deveriam ser concentrados, e d) identificação de áreas de ocorrência de cardeal-amarelo que poderiam ser transformadas em áreas protegidas.

## 1.2. O cardeal-amarelo

*Gubernatrix cristata* é classificado como uma ave da ordem Passeriformes Linnaeus, 1758, Subordem Passeri Linnaeus, 1758, Parvordem Passerida Linnaeus, 1758, Família Emberizidae Vigors, 1825, e a única espécie do gênero *Gubernatrix* (Bencke, 2001; CBRO, 2009). Sinonímia - *Coccothraustes cristata* (Wetmore, 1926), *Gubernatrix cristatella* (Hudson, 1920).

O macho tem peito e costas amarelo brilhante e crista, linha dos olhos e garganta pretos (Fig. 1.1). Ele tem a sobrancelha e faixa malar amarela enquanto que na fêmea são brancas, com um peito cinzento (Belton, 1994; Sick, 1997). O macho mede entre 19,2-20,1 cm e pesa cerca de 46g.

Cristi (1958) descreveu a infestação de um macho adulto por filarídeos (Nematoda, Filarioidea) subfamília Aproctinae e Muniz-Pereira e colaboradores (2009) registraram um filarídeo *Pelecitus* sp. (Nematoda, Filarioidea) da Família Onchocercidae infestando um indivíduo da espécie.

Durante a estação reprodutiva o macho defende seu território atacando de forma agressiva outros machos. Usando um macho de cativeiro como chamariz, esse comportamento territorialista passou a ser usado pelos caçadores como uma forma de capturá-los (Fontana et al. 2003). O acasalamento ocorre na primavera austral, de novembro a fevereiro, com ninhos contendo 2 a 3 ovos azuis esverdeados e manchados de preto (Fig. 1.2) com um período de incubação de cerca de 13 dias (Belton, 1994). Em um trabalho de campo realizado em janeiro de 2007 em Lavalleya, Uruguai, encontramos um ninho (Fig. 1.3) em uma árvore de cerca de 4m de altura conhecida como Viraró, *Ruprechtia salicifolia* da família Polygonaceae.

A alimentação consiste principalmente de grãos, embora quando filhotes são alimentados basicamente de insetos.

O cardeal-amarelo é uma espécie de ave rara do Pampa. Ele tem uma distribuição

geográfica restrita ao sul da América do Sul e é exclusivo desse bioma (Uruguai, Argentina e sul do Brasil) (Azpiroz, 2003). Esta espécie tem sofrido um dramático declínio populacional ao longo de sua área de distribuição e seu estado de conservação é definido como ameaçado de extinção (En) (*Endangered* da sigla em inglês) (BirdLife International, 2000; Fontana et al., 2003).

Na Argentina é agora raro exceto muito localmente (Fraga, 1997). Pode ser encontrada em San Luis, Buenos Aires, La Pampa e Río Negro (especialmente entre General Conesa, San Antonio Oeste e Viedma). Há populações importantes em Corrientes (Pay Urbe e Estancia San Antonio), Entre Ríos (recentemente na área de Montiel, Ceibas e Estancia La Choza) e Córdoba [recentemente em Chancaní (Chebez et al., 1998)] (Di Giacomo, 2005). Ao mesmo tempo, Córdoba apresenta uma redução enorme da população de cardeal-amarelo, apesar de alguns novos registros (Kopta, 1999). Há registros anteriores a 1975 de Salta, Tucumán, Santiago del Estero, Santa Fe e San Juan, e foi registrado sem detalhes para Formosa, Chaco, La Rioja e (erroneamente) em Misiones (Chebez, 1996). Matteucci e colaboradores (2007) afirmaram que a zona dos *bajos submeridionales* é a área mais importante em Santa Fé uma que é a zona onde se concentram a maior densidade de cardeal-amarelo no Chaco argentino. Os autores, entretanto não dizem o quão grande é a população nem se está baixando.

No Uruguai, ele foi historicamente conhecido em 13 departamentos, mas recentemente apenas de Paysandú, Río Negro, Florida, Rocha, Lavalleja e Artigas (Azpiroz, 2003; BirdLife International, 2000; Lapitz, 2010; Martins-Ferreira, 2007). Em 1999, a população uruguaia de cardeal-amarelo foi estimada em menos de 300 indivíduos, concentrados principalmente na Bacia do Rio Uruguai (Azpiroz em litt. 1999).

Dois indivíduos coletados no Paraguai em 1905 eram provavelmente aves de gaiolas que escaparam do cativeiro (Di Giacomo, 2005).

No Sul do Brasil (Rio Grande do Sul) já era considerado raro na década de 1970 e 1980 (Belton, 1985). Desde então, ele sofreu uma drástica queda e agora se acredita que esteja limitado a locais restritos da Serra do Sudeste, próximo a fronteira com o Uruguai e na ponta Oeste do Rio Grande do Sul (Fontana et al., 2003). Há um registro de um macho de *G. cristata* em um parque urbano de Porto Alegre (Parque Mascarenhas de Moraes) em outubro/novembro de 2005 (Scherer et al., 2010). Este registro não representa qualquer extensão de sua distribuição real, uma vez que é muito provavelmente uma fuga do cativeiro como concluíram os próprios autores do trabalho.

*Gubernatrix cristata* é encontrado em bosques (incluindo bosques de *Prosopis*), planícies, matagais e estepes arbustivas, com cerca de 700 m acima do nível do mar (Di Giacomo, 2005). Há uma população reprodutiva no Rio Grande do Sul, Brasil, no Parque Estadual do

Espinilho (Damiani et al., 2009). Este parque é uma extensão da província do Espinal, unidade fitogeográfica que descreve um grande arco irregular em território argentino, a partir do centro da cidade de Corrientes e Entre Ríos, na costa atlântica, na província do sul de Buenos Aires, passando pela região central de Santa Fé, Córdoba, San Luís e La Pampa (Marchiori, 2004). Membro do domínio Chaqueño o Espinal se separa da Província Chaqueña pela ausência de espécies do gênero *Schinopsis* (Anacardiaceae) (Marchiori 2004). Dos seus três distritos fitogeográficos, o do "Nandubay", situado mais ao norte (ou nordeste do arco), inclui o Parque do Espinilho e distingue-se pela associação de *Prosopis affinis* e *P. nigra*. Na Mesopotâmia argentina, o "Espinal" também pode incluir: o chañar (*Geoffroea decorticans*), sombra-de-touro (*Jodina rhombifolia*) e até mesmo duas palmas - a jataí (*Butia yatay*) e carandá (*Trithrinax campestris*) - além das espécies anteriormente citadas (Cabrera & Willink, 1973).

### 1.3. As Ameaças

Devido a falta de avistamentos do cardeal-amarelo nos últimos anos e sua situação ruim em áreas adjacentes no Uruguai, a população da serra do sudeste parece estar isolada ou mesmo extinta (Martins-Ferreira, 2007). A única população reprodutiva selvagem conhecida no Brasil de *G. cristata* parece estar limitada ao Parque Estadual do Espinilho e arredores (Damiani et al., 2009).

A exploração constante e crônica como ave para o mercado de aves de gaiola continua a ser a ameaça mais significativa. A destruição e a fragmentação de seu habitat foram enumerados como possíveis ameaças adicionais para a espécie, bem como a extrema pressão de captura para abastecer o tráfico de animais silvestres (Chebez, 1994; BirdLife International 2000). A principal razão por trás do tráfico é que é um pássaro de bonito colorido e canto melodioso. O crescente registro na Argentina de híbridos entre diuca (*Diuca diuca*) e cardeal-amarelo parece ser uma resposta à escassez de indivíduos do sexo masculino na população da espécie (Bertonatti & Guerra, 1997, 2001). No Rio Grande do Sul, o cardeal-amarelo é capturado principalmente nos municípios de Piratini e Pinheiro Machado. A descaracterização e destruição do habitat foram listados como potenciais ameaças adicionais à espécie (Chebez, 1994; BirdLife International, 2000). No Rio Grande do Sul, a maior parte da paisagem original do sul e do oeste foi convertida em plantações de arroz e pastagens para o gado. No entanto, desconhece-se até que ponto essas atividades afetam a espécie. Presumivelmente, o seu habitat sofreu com a conversão para pastagem de gado, extração de madeira para lenha e mobiliário (Chebez, 1994) e, especialmente, pela arborização rápida com plantios de eucalipto.

Hudson (1920) já atestava que suas qualidades como belo pássaro com voz alta e musical fizeram dele um pássaro de gaiola preferido. Loydi (2008) encontrou oito indivíduos de cardeal-amarelo em pet shops da cidade de Bahía Blanca, Província de Buenos Aires, em 2004 e mais 3 em 2007. A comercialização das espécies ainda continua. Pereira e Brito (2005) encontraram *G. cristata* sendo vendido na região metropolitana do Recife em feiras livres. Pessino e Titarelli (2006) disseram que é uma ave ainda muito valorizada como pássaro de estimação e sugeriram uma ação conjunta entre as três províncias do limite sul da sua distribuição (Buenos Aires, Río Negro e La Pampa) como forma de diminuir a pressão sofrida pelas populações da espécie. Dois indivíduos foram avistados em dezembro de 1996 em Río Negro, onde era uma espécie comum que foi diminuindo devido à caça (Seewald e Perez, 2009). Uma ameaça sofrida pelas aves em geral em comunidades locais pobres, incluindo espécies ameaçadas de extinção, é o uso delas como meio de subsistência através da venda em autoestradas (Pautasso, 2003).

#### **1.4. Métodos gerais**

Atualmente muitas espécies enfrentam os problemas da redução severa de suas áreas de distribuição causadas pela fragmentação e, em casos mais extremos, a perda total dos seus habitats devido ao crescimento da atividade humana nas últimas décadas. Tanto a perda do habitat quanto o isolamento dos fragmentos remanescentes, pode reduzir o tamanho populacional a níveis tão baixos que a extinção local pode ser atingida (Gaines *et al.*, 1997; Lacy, 1997).. Dois processos têm uma grande importância no aumento do risco de extinção em pequenas populações. Em primeiro lugar, processos não genéticos, tanto no nível demográfico quanto ambiental, podem afetar parâmetros tais como a reprodução, mortalidade, distribuição por classes de idade e proporção sexual seguidas de flutuações no tamanho populacional (Goodman, 1987; Shaffer, 1987). Segundo, quanto menor é o tamanho populacional, mais suscetível ficam as populações a processos genéticos, tais como, deriva genética e endocruzamento, seguidos de perda da variabilidade genética populacional ou aumento da homozigose (Shields, 1993; Hartl & Clark, 1997; Broders *et al.*, 1999) e do efeito de depressão por endocruzamento, ou seja, diminuição do *fitness* da descendência por efeito do cruzamento entre aparentados. Esses processos genéticos conhecidos como erosão genética, podem causar degradação de importantes características da espécie como a sobrevivência, crescimento e reprodução (Shields, 1993; Hedrick *et al.*, 1996).

Apesar de que a discussão a respeito do real impacto da depressão por endocruzamento em populações naturais ainda existe, e que os padrões observados em animais de cativeiro podem ser bem diferentes daqueles achados na natureza (Hoelzel *et al.*, 1993; Craig, 1994) a maioria dos planos de conservação atuais monitoram, tanto o tamanho, quanto a estrutura populacional e a variabilidade genética das espécies (Lande, 1995; Gaines *et al.* 1997).

A ferramenta molecular mais usada neste tipo de abordagem são as análises de DNA mitocondrial (mtDNA) que permitem o seguimento de rastros genealógicos além dos limites genéticos entre as populações, espécies e grupos taxonômicos mais elevados.

Outra vantagem no uso das informações obtidas a partir da análise da diversidade genética são: redução do risco de extinção reduzindo o endocruzamento e perda da diversidade genética; identificação de populações importantes, do ponto de vista de variabilidade genética; resolução da estrutura populacional; definição de unidades de manejo dentro das espécies; detecção de hibridizações; permite amostragem não-intrusiva para análise genética; possibilita a definição de sítios para reintrodução e escolha da melhor população para reintrodução; permite uma melhor compreensão da biologia da espécie (Awise *et al.*, 1987a; Gaines *et al.*, 1997; Lacy, 1997; Warning, 1994, 2000; Frankham *et al.*, 2002; Sutherland *et al.*, 2004).

Isto levou a melhor compreensão de biogeografias regionais e áreas de endemismo, bem como à elaboração de propostas prioritárias para a conservação das biodiversidades taxonômicas e locais (Rojas, 1995; Moritz & Bermingham, 1998; da Silva & Patton, 1998; Firestone *et al.*, 1999; Jerusalinsky, 2001).

A maioria das espécies distribuem-se em populações geograficamente estruturadas, muitas das quais podem experimentar pouco ou nenhum contato por longos períodos de tempo. Outras espécies, entretanto, podem estar caracterizadas por expansões recentes, o que determina estreitas conexões genealógicas entre as populações resultantes (Ibrahim *et al.*, 1996).

Características demográficas históricas e contemporâneas podem afetar a estrutura espacial de populações co-específicas e com ela influenciar nas genealogias matrilineares em vários sentidos. Um importante desafio dos estudos filogeográficos é decifrar os fatores demográficos passados e presentes que produzem um determinado padrão de relações genealógicas (Awise, 2000).

Duas metas fundamentais da genética da conservação são a prevenção de problemas associados ao endocruzamento e à deriva genética em pequenas populações e a descrição da estrutura populacional com a finalidade de identificar unidades de conservação. A genética

molecular tem proporcionado numerosas técnicas de acesso à variabilidade genética dentro e entre as populações, tanto através de polimorfismos de proteínas quanto de DNA (Awise, 1994). Por serem altamente polimórficos e possuírem altas taxas de mutação (dentre outras características), os marcadores moleculares mais utilizados em estudos populacionais e filogeográficos são o DNA mitocondrial e os *loci* de microssatélites.

O genoma mitocondrial de animais é haplóide e está formado por uma dupla fita circular entre 15000 e 17000 pares de bases de comprimento estando presente em centenas a milhares de cópias por célula (Meyer, 1993; Li & Graur, 2000).

Tipicamente cada genoma de DNA mitocondrial (mtDNA) consiste de 37 genes funcionais sem longos espaços intergênicos. Estes *loci* codificam para 22 RNAs de transferência, dois RNAs ribossômicos e 13 RNAs mensageiros para a síntese de subunidades de proteínas específicas envolvidas na fosforilação oxidativa e o transporte de elétrons (Awise, 2000).

O mtDNA é simples em estrutura e econômico em tamanho sendo que o único grande fragmento não codificante (de aproximadamente 1000 pares de bases) é a região controladora (CR - *Control Region*) da molécula que tem função regulatória na dinâmica da molécula e é predominantemente rica em bases AT (Brown, 1985). Nos vertebrados esta região é referida como *D-loop* (*Displacement-loop*) devido à formação de uma estrutura de fita tripla que gera deslocamentos dinâmicos no começo da replicação do mtDNA (Brown et al., 1986).

Devido à rápida evolução o mtDNA torna-se um bom marcador para o estudo de diferenciações genéticas recentes por acumular substituições de base, inserções e deleções com uma taxa média de 5 a 10 vezes mais rápida que o DNA nuclear cópia simples (Brown et al., 1979). Várias hipóteses têm sido levantadas na tentativa de explicar a rápida evolução do DNA mitocondrial: alta taxa de mutação devido à ineficiência dos mecanismos de reparo de DNA, alta exposição aos radicais livres (mutagênicos) produzidos no interior da mitocôndria ou ao rápido *turnover* dentro das linhagens celulares. Também o fato do DNA não estar complexado com proteínas histonas, que são evolutivamente conservadas e poderiam limitar a taxa evolutiva do DNA nuclear (Awise, 2000). Outra vantagem do mtDNA, como marcador para o estudo de diferenciações genéticas recentes, é a frequência extremamente baixa de transposições, inversões, rearranjos e recombinação (Brown, 1985). Isto permite a caracterização de linhagens filogenéticas sem a ambigüidade causada pela recombinação meiótica que ocorre nos genes nucleares.

Foram feitas 14 viagens de campo de Abril de 2006 a março de 2009 ao Rio Grande do Sul e Uruguai, em uma tentativa de encontrar indivíduos e coletar amostras para análise genética.



Todos os exemplares foram anilhados com anilhas fornecidas pelo CEMAVE/ICMBio (Centro Nacional de Pesquisa para a Conservação de Aves Silvestres). Na Argentina, uma visita foi feita ao Museo de Ciencias Naturales Bernardino Rivadavia onde, a partir de oito peles escolhidas foram coletadas amostras de tecido, uma de cada província onde a espécie ocorreu, e uma do Uruguai. Um total de 97 amostras de cardeal-amarelo foram usadas neste estudo (Apêndice A.1) e apenas 72 funcionaram adequadamente.

Uma biblioteca de DNA genômico enriquecido em microssatélites de *Gubernatrix cristata* foi construído e dez loci de microssatélites polimórficos nucleares (com repetições de 2 bp) foram selecionados com base nos níveis de polimorfismo e tamanho dos produtos de amplificação. Os Primers desenhados e as condições de PCR são dadas no capítulo 2.

Nós extraímos DNA de amostras de pena utilizando o protocolo Alkali para extração de DNA e tentamos sequenciar vários loci de DNA mitocondrial (mtDNA) utilizando primers de aves degenerados de Sorenson (2003) para determinar se variabilidade genética suficiente estava presente para análise. A região controle, ATPase8 e ND2 foram escolhidos, uma vez que estas regiões têm se mostrado úteis em estudos de genética de populações de aves (Burg & Croxall, 2001; Oyler-McCance et al., 2005; Sorenson, 2003). Nós amplificamos com sucesso porções do ND2, mas não fomos capazes de amplificar a região controle e ATPase8. Dada a baixa qualidade do DNA extraído de várias amostras decidimos amplificar o gene em dois pedaços e correr quatro reações de seqüenciamento (ambas as fitas de cada produto de PCR). Os primers usados e as condições de PCR estão descritos no Capítulo 3.

Todos os registros georeferenciados de *Gubernatrix cristata* foram extraídos de diferentes fontes: registros em GPS durante trabalhos de campo, peles de museu, literatura e bases de dados da Internet. No total este conjunto foi composto por 172 registros georeferenciados pertencentes a indivíduos, casais ou um pequeno número de aves vistas na zona onde os registros foram tomados.

Para modelar a distribuição das espécies, foram selecionados 20 indicadores ambientais. Nós fizemos o download de 19 variáveis bioclimáticas (1950 - 2000) e altitude (em m) do conjunto de dados do WORLDCLIM (<http://www.worldclim.org>) para a América do Sul na resolução de 30 arco-segundo (~ 1km) (Hijmans et al., 2005a).

BIOCLIM (usando a plataforma DIVA-GIS) e MaxEnt foram escolhidos para nossos dois modelos preditivos biogeoclimáticos separados. DIVA-GIS é um plataforma baseada em SIG do modelo BIOCLIM (Hijmans et al., 2005b) que se encaixa num envelope de habitat mínimo da espécie em um espaço climático multidimensional. MaxEnt é um programa desenvolvido para a

modelagem de entropia máxima de distribuição geográfica das espécies (Phillips et al., 2006) que expressa a adequação de cada célula da grade em função das variáveis ambientais nessa quadrícula. Alto valor de funções indica condições adequadas previstas para a espécie. Estes modelos foram selecionados por acomodarem os dados de presença apenas e abrangem uma variedade de plataformas de modelagem (Elith et al., 2006).

### 1.5. Resultados gerais

Em populações sofrendo um estrangulamento genético, a diversidade genética observada excede a diversidade genética esperada em equilíbrio sob a suposição de equilíbrio mutação-deriva. A hipótese nula testada para o excesso de heteroziguidade usando o teste Wilcoxon de rank de sinal fornecido ( $P < 0,05273$ ), ( $P > 0,09668$ ) e ( $P < 0,42285$ ) probabilidades no âmbito do Modelo de Alelos Infinitos (IAM), Mutação de Duas Fases (TPM) e Mutação Step-wise (SMM), respectivamente. A hipótese nula é aceita no modelo SMM apenas, o que implica que *Gubernatrix cristata* não sofreu qualquer estrangulamento genético recente. Os valores estimados do excesso de heteroziguidade e suas probabilidades de teste do sinal foram 5,89 ( $P < 0,15033$ ) para o IAM, 5,90 ( $P < 0,15148$ ) para o TPM e 5,91 ( $P < 0,38936$ ) para o SMM. A hipótese de equilíbrio mutação-deriva também não foi aceita com base no teste de sinal sob os modelos de mutação.

Somente 67 amostras foram seqüenciadas com sucesso para o fragmento ND2 alvo (seqüências foram depositadas no GenBank sob números de acesso tal e tal). Um total de 349 pb foram alinhados em amostras de *G. cristata* contendo sítios informativos parcimoniosos e variáveis e um total de três haplótipos (H1 - H3). Entre os haplótipos H1 e H2 75 passos mutacionais foram encontrados, enquanto 34 entre H2 e H3. Medidas de variabilidade genética tais como diversidade haplotípica ( $Hd$ ), variação nucleotídica estimada com base no número de sítios segregantes ( $\pi$ ) e o número médio de diferenças em pares de populações amostradas ( $k$ ) são mostrados no Capítulo 3. Estimativas de diversidade nucleotídica e haplotípica na amostra foram relativamente elevadas, com  $\pi = 0,00277$  e  $Hd = 0,6219$ . A análise de network não revelou padrão de estrutura algum. O  $F_{st}$  (0,00340) e  $Nm$  (73,18) mostraram uma estruturação fraca com pouca diferenciação genética. Falta de apoio para a hipótese de expansão demográfica foi também obtida através de diferentes testes realizados para testar a teoria neutra da evolução molecular: teste estatístico  $D^*$  de Fu e Li (0,71916) não apresentaram significância estatística ( $P > 0,10$ ), bem como não foi significativo o teste estatístico  $F^*$  de Fu e Li (1,34814) ( $P > 0,10$ ). Estatística  $F_s$  de Fu foi

2,248 o qual foi significativamente diferente de zero e é uma evidência provável indicando uma diminuição no tamanho da população e/ou seleção equilibradora.  $D$  de Tajima foi positivo (2,17506), o que significa baixos níveis de polimorfismos de frequência alta e baixa, indicando uma diminuição no tamanho da população e/ou seleção equilibradora.

Uma vez que a origem geográfica exata de muitos dos indivíduos amostrados (45%) cardeal-amarelo é desconhecida ou incerto (muitos dos indivíduos que foram amostrados em zoológicos ou de particulares, ou recebido de apreensões policiais não foram associados a informações confiáveis sobre a sua localização exata de captura), a análise do Structure (Falush *et al.*, 2003) seria particularmente relevante porque isso permitiria o agrupamento de indivíduos, sem a necessidade de uma informação geográfica a priori. Cinco independentes corridas para 1 a 6 populações, utilizando o modelo admixture e frequências alélicas afins entre as populações (Falush *et al.* 2003), foram executados no programa Structure. Os resultados de todas as corridas (mostrado no capítulo 3) foram praticamente os mesmos sugerindo que não houve aglomeração de indivíduos. Devido a esse resultado nenhuma atribuição individual a qualquer área geográfica foi possível.

A distribuição das diferenças da variação haplotípica observada foi unimodal revelando uma partida da variação esperada em uma população constante ( $k = 0,965$ , variação de  $k = 0,7205$ ,  $r = 0,1319$  estatística de irregularidade).

Os mapas com uma discussão detalhada de nossos resultados sobre os modelos de distribuição da espécie são encontrados no Capítulo 4.

A partir da execução desse projeto avançamos o conhecimento sobre *Gubernatrix cristata*, principalmente no que se refere ao seu status populacional dentro de sua área de distribuição. Conseguimos determinar claramente onde a espécie ainda sobrevive no Brasil e quais são, de fato, os melhores locais para se investir na sua conservação. A partir desse projeto, definiu-se ações para o manejo da espécie em cativeiro. Foram estabelecidas parcerias e novos contatos que possibilitaram a centralização de todos os exemplares oriundos de apreensão no RS em um local: o zôo de Gramado. Esse foi o primeiro passo para a criação de um programa de reprodução em cativeiro da espécie, bem como um protocolo sanitário nacional e o engajamento de instituições públicas, privadas e ONGs para a conservação de *G. cristata*. Um dos impactos mais significativos do projeto foi o estímulo à criação do Plano de Ação do Cardeal-Amarelo, uma vez que enquanto não há um projeto efetivo de conservação para uma espécie ameaçada não há como definir ações por parte do ICMBio. Algumas dessas ações, embora ainda não tenha sido publicado o plano, já foram executadas e outras estão sendo realizadas. O Projeto Cardeal-Amarelo iniciou

em 2005 como um projeto-piloto na região de Pinheiro Machado e Pedras Altas. Ele tomou forma em 2006 e 2007 com o apoio da Fundação O Boticário de Proteção à Natureza. E seguirá ainda por muitos anos à frente, pois muitas questões ainda precisam ser respondidas.

### **1.1. General Introduction and Thesis Aims**

This chapter provides detailed information about the study species and the study area, as well as general methods used in this research study. The first part of the chapter consists of a detailed description of Yellow Cardinal (*Gubernatrix cristata*), providing information on aspects of taxonomy, morphology, geographic distribution, ecology, behavior, and an overview of the conservation status of the species and the threats it faces in the wild. The second part provides information about the threats faced by Yellow Cardinal. The third part gives the general methods used in this research study and the fourth part provides a synthesis of found results.

Our main objective was to determine the status of *Gubernatrix cristata* populations within its distribution range, especially those found in Rio Grande do Sul. We focused on aspects of reproductive biology, population structure and gene flow. The main idea behind this information was to propose strategies for their conservation. To better achieve that target we specifically focus our research on a) locating wild populations of this species in Rio Grande do Sul, Uruguay and Argentina, b) comparing the different populations in terms of genetic variability and to check for gene flow between them using both nuclear (microsatellites) and mitochondrial (ND2) molecular markers, c) identifying the most important populations in terms of genetic variability, where conservation efforts should be concentrated, and d) identifying areas of occurrence of yellow cardinal that could be turned into protected areas.

### **1.2. The bird**

*Gubernatrix cristata* is classified as a bird in the Order Passeriformes Linnaeus, 1758, Suborder Passeri Linnaeus, 1758, Parvorder Passerida Linnaeus, 1758, Family Emberizidae Vigors, 1825, and the only species of genus *Gubernatrix* (Bencke, 2001; CBRO, 2009). Synonymy – *Coccothraustes cristata* (Wetmore, 1926), *Gubernatrix cristatella* (Hudson, 1920).

The male has a bright yellow chest and back with black crest, eye line and throat (Fig. 1.1). It has the full eyebrow and malar stripe yellow while the female are white with a gray breast (Belton, 1994; Sick, 1997). The male measured 19.2 to 20.1 cm and weighs about 46 g.

Cristi (1958) described the infestation of an adult male by filarids (Nematoda, Filarioidea) subfamily Aproctinae and Muniz-Pereira *et al.* (2009) have recorded a filarid *Pelecitus* sp. (Nematoda, Filarioidea) from Family Onchocercidae infesting the bird.

During the breeding season the male defends its territory aggressively attacking other males. That behavior turns to be a way how poachers capture them using a captive male as decoy (Fontana *et al.* 2003). The breeding season occurs in the austral spring, from November to February, with nests containing two to three greenish blue and black spotted eggs (Fig. 1.2) with an incubation period of around 13 days (Belton, 1994). In a fieldwork in Jan 2007 in Lavalleja Department, Uruguay, we found a nest (Fig. 1.3) in a tree of circa 4m height known as *Viraró*, *Ruprechtia salicifolia* of family *Polygonaceae*.

The food consists mainly of grains, although while nestlings are fed essentially on insects.

The Yellow Cardinal, is a rare bird species from the Pampas grassland. It has a restricted geographical distribution in Southern South America and is unique to this Biome (Uruguay, Argentina, and Southern Brazil) (Azpiroz, 2003). This species has undergone a dramatic population decline across its range and its conservation status is defined as endangered (BirdLife International, 2000; Fontana *et al.*, 2003).

In Argentina, it is now rare except very locally (Fraga, 1997). It is found in San Luis, Buenos Aires, La Pampa and Río Negro (especially among General Conesa, San Antonio Oeste and Viedma). There are important populations in Corrientes (Pay Urbe and Estancia San Antonio), Entre Ríos (recently in the area of Montiel, Ceibas and Estancia La Choza) and Cordoba [recently in Chancaní (Chebez *et al.*, 1998)] (Di Giacomo, 2005). At the same time Cordoba present a huge lowering of Yellow Cardinal population, despite some new records (Kopta, 1999). There are records prior to 1975 from Salta, Tucumán, Santiago del Estero, Santa Fe and San Juan, and it was recorded with no details to Formosa, Chaco, La Rioja and (wrongly) in Misiones (Chebez, 1996). Matteucci *et al.* (2007) say that *bajos submeridionales* zone is the most important area in Santa Fe since is the zone which concentrate the largest density of yellow cardinal in the Argentinean *chaco*. The authors although didn't say how large is that population neither if it is lowering.

In Uruguay, it was historically known from 13 departments, but recently from only Paysandú, Río Negro, Florida, Rocha, Lavalleja and Artigas (BirdLife International, 2000; Azpiroz, 2003; Martins-Ferreira, 2007; Lapitz, 2010). In 1999 the Uruguayan population of yellow cardinal was estimated less than 300 individuals, mostly concentrated in the Uruguay River Basin (Azpiroz *in litt.* 1999).

Two individuals collected in Paraguay in 1905 were probably caged birds escaped from captivity (Di Giacomo, 2005).

In Southern Brazil (Rio Grande do Sul State) it was already considered rare in the 1970s and 1980s (Belton, 1985). Since then it suffered a dramatic decline and now is believed it is confined to restricted spots in the SE hills, near Uruguayan border and at W tip of Rio Grande do Sul (Fontana *et al.*, 2003). There is a record of a male *G. cristata* to an urban park (Mascarenhas de Moraes Park) in Porto Alegre City (Brazil, RS) in Oct/Nov 2005 (Scherer *et al.*, 2010). This record doesn't represent any extension of its actual distribution since it is most probably an escape from captivity, as conclude by the papers authors themselves.

*Gubernatrix cristata* is found in woodlands (including forests of *Prosopis*), plains, bush, and shrub steppes, to about 700 m above sea level (Di Giacomo, 2005). There is a breeding population in Rio Grande do Sul State, Brazil, at State Park of *Espinilho* (*Parque Estadual do Espinilho*) (Damiani *et al.*, 2009). This park is an extension of the Province of *Espinal*, phytogeographical unit that describes a large irregular arc in Argentinean territory, from the center of Corrientes and Entre Ríos, to the Atlantic coast at the Southern Buenos Aires province, passing through the central region of Santa Fe, Cordoba, San Luis and La Pampa (Marchiori, 2004). Member of "*Chaqueño* Domain" the *espinal* branches off the "*Chaco* Province" by the absence of actual species of the genus *Schinopsis* (Anacardiaceae) (Marchiori 2004). Of his three phytogeographic districts, "*Nandubay*" located further north (or northeast of the arch), includes the Park of *Espinilho* and distinguishes itself by the association of *Prosopis affinis* and *P. nigra*. In the Argentinean Mesopotamia, the "*Espinal*" may also include: the *chañar* (*Geoffroea decorticans*), *sombra-de-touro* (*Jodina rhombifolia*) and even two palms - the *jataí* (*Butia yatay*) and *carandá* (*Trithrinax campestris*) – in addition to species previously cited (Cabrera & Willink, 1973).

### 1.3. The Threats

Once the lack of yellow cardinal sightings in the last few years and its bad situation in the adjacent areas in Uruguay, the SE hills population seems to be isolated or even extinct (Martins-Ferreira, 2007). The only breeding wild population known in Brazil of *G. cristata* seems to be limited to State Park of *Espinilho* (*Parque Estadual do Espinilho*) and surroundings (Damiani *et al.*, 2009).

The constant and chronic exploitation as songbird for the cage bird market remains the most significant threat. The destruction and fragmentation of its habitat have been enumerated as

possible additional threats for the species, as well as the extreme pressure of capture to supply the traffic of wild animals (Chebez, 1994; BirdLife International 2000). The main reason behind the traffick is that bird is a colorful and singing passerine. The growing record of hybrids among yellow cardinal and common diuca finch (*Diuca diuca*) in Argentina seems to be a response to the shortage of male individuals in the population of the species (Bertonatti & Guerra, 1997, 2001). In Rio Grande do Sul, Yellow Cardinal is caught mainly in the municipalities of Piratini and Pinheiro Machado. The mischaracterization and destruction of habitat have been listed as potential additional threats to the species (Chebez, 1994; BirdLife International, 2000). In Rio Grande do Sul, most of the original landscape of the south and west has been converted into rice fields and pastures for cattle. However, it is unknown to what degree these activities affected the species. Presumably its habitat suffered with conversion for livestock grazing, timber extraction for firewood and furniture (Chebez, 1994) and especially by the rapid forestation with *Eucalyptus* plantations.

Hudson (1920) already attested that its qualities as beautiful bird with loud and musical voice made it a favorite cage bird. Loydi (2008) had found 8 Yellow Cardinal individuals in pet shops in Bahía Blanca City, Buenos Aires Province, in 2004 and 3 more in 2007. The trading of the species still goes. Pereira and Brito (2005) had found *G. cristata* being sold at Recife metropolitan area on free market. Pessino and Titarelli (2006) said that is still fairly valued as a pet bird and suggested joint action between the three provinces of the Southern limit of its distribution (Buenos Aires, Rio Negro and La Pampa) as a way to slow the pressure suffered by the species populations. Seewald and Perez (2009) reported the sighting of two individuals in Dec 1996 in Río Negro and it was a common species that has been decreasing due to hunting. Pautasso (2003) reports the use of birds in general by local poor communities, including endangered species as a means of livelihood for sale in routes.

#### **1.4. General Methods**

Today many species face the problems of severe reduction of their distribution areas caused by fragmentation, and in more extreme cases, total loss of their habitat due to increased human activity in recent decades. Both loss of habitat and the isolation of fragments, can reduce the population size to levels so low that local extinction can be achieved (Gaines *et al.*, 1997; Lacy, 1997). Two processes are of great importance in increasing the risk of extinction in small populations. First, non-genetic processes at both the demographic and environmental level, can

affect parameters such as reproduction, mortality, distribution by age and sex ratio followed by fluctuations in population size (Goodman, 1987; Shaffer, 1987). Second, the smaller the population size, the more susceptible to genetic processes they are, such as genetic drift and inbreeding. These processes are followed by loss of genetic variability and population increase of homozygosity (Broders *et al.*, 1999; Hartl & Clark, 1997; Shields, 1994;) and the effect of inbreeding depression, i.e., reduced fitness of offspring due to crossing among relatives. These genetic processes known as genetic erosion can cause degradation of important species characteristics such as survival, growth and reproduction (Hedrick *et al.*, 1996; Shields, 1994).

Although the discussion about the actual impact of inbreeding depression in natural populations still exists, and that the patterns seen in captive animals may be quite different from those found in nature (Craig, 1994; Hoelzel *et al.*, 1993) most conservation plans current monitor, both the size and the population structure and genetic variability (Gaines *et al.*, 1997; Lande, 1995). The most widely used molecular tool in this approach is the analysis of mitochondrial DNA (mtDNA) that allow the following of pedigree trace beyond the genetic boundaries between populations, species and higher taxonomic groups. Another advantages in the use of information obtained from the analysis of genetic diversity are: reducing the risk of extinction by reducing inbreeding and loss of genetic diversity; identification of important populations from the standpoint of genetic variability; resolution of population structure; definition of management units within species; detection of hybridization; allows non-intrusive sampling for genetic analysis; enables the definition of sites for reintroduction and choosing the best population for reintroduction; and allows a better understanding of the species biology (Avisé *et al.*, 1987a; Frankham *et al.*, 2002; Gaines *et al.*, 1997; Lacy, 1997; Sutherland *et al.*, 2004). This led to better understanding of regional biogeography and areas of endemism, as well as to develop proposals for the conservation of biodiversity and local taxonomic (Bermingham & Moritz, 1998; Firestone *et al.*, 1999; Da Silva & Patton, 1998; Rojas, 1995). Most species are distributed in geographically structured populations, many of them may experience little or no contact for long periods of time. Other species, however, can be characterized by recent expansions, which determine close genealogical connections between the resulting populations (Ibrahim *et al.*, 1996). Historical and contemporary demographic characteristics may affect the spatial structure of conspecific populations and influence the matrilineal genealogy in multiple directions (Avisé, 2000). Two fundamental goals of conservation genetics is the prevention of problems associated with inbreeding and genetic drift in small populations and the description of population structure in order to identify conservation units. Molecular genetics has provided numerous technical access



to the genetic variability within and between populations, either through polymorphisms of proteins as through DNA (Avice, 1994).

Because they are highly polymorphic and have high mutation rates (among other features), the most widely used molecular markers in population and phylogeographic studies are mitochondrial DNA and microsatellite loci. The animal mitochondrial genome is haploid and is formed by a double-stranded circular between 15000 and 17000 base pairs in length and is present in hundreds to thousands of copies per cell (Li & Graur, 2000; Meyer, 1993). Typically each mitochondrial DNA genome (mtDNA) consists of 37 functional genes without long intergenic spaces. These loci coding for 22 transfer RNAs, two ribosomal RNAs and 13 messenger RNAs for the synthesis of subunits of specific proteins involved in oxidative phosphorylation and electron transport (Avice, 2000). The mtDNA is simple in structure and economic in size. The only major non-coding fragment (about 1000 base pairs) of the molecule is the control region (CR - Control Region) that has a regulatory role in the dynamics of the molecule and is predominantly rich in AT bases (Brown, 1985). In vertebrates this region is referred to as D-loop (displacement-loop) due to the formation of a triple ribbon that generates dynamic shifts early in the replication of mtDNA (Brown *et al.*, 1986). Due to the fast evolution mtDNA became a good molecular marker for the study of recent genetic differences due to accumulate base substitutions, insertions and deletions at an average rate 5-10 times faster than the single copy nuclear DNA (Brown *et al.*, 1979). Several hypotheses have been raised in an attempt to explain the rapid evolution of mitochondrial DNA: high mutation rate because of the inefficiency of the mechanisms of DNA repair, high exposure to free radicals (mutagens) produced inside the mitochondria or the rapid turnover inside the cell lineages. Other fact is that DNA is not complexed with histone proteins, which are evolutionarily conserved and could constrain the evolutionary rate of nuclear DNA (Avice, 2000). Another advantage of mtDNA as a marker for the study of recent genetic differences is the extremely low frequency of transpositions, inversions, rearrangements and recombination (Brown, 1985). This allows the characterization of lineages without the ambiguity caused by meiotic recombination that occurs in nuclear genes.

Fourteen field trips were made from April 2006 to March 2009 in an attempt to find individuals and collect samples for genetic analysis in Rio Grande do Sul and Uruguay. All of them were ring banded with rings provided by CEMAVE/ICMBio (National Research Center for the Conservation of Wild Birds). In Argentina, a visit was made to the Museo de Ciencias Naturales Bernardino Rivadavia from which tissue samples of eight skins were collected, one from each province where the species has occurred, and one from Uruguay. A total of 97 yellow cardinal

samples was taken in this study (Appendix A.1) and only 72 worked properly.

A microsatellite-enriched genomic DNA library of *Gubernatrix cristata* was constructed and ten polymorphic nuclear microsatellite loci (with 2-bp repeats) were selected on the basis of polymorphism levels and size of amplification products. Primers and PCR conditions are given in Chapter 2.

We extracted DNA from feather samples using the Alkali protocol for DNA extraction and attempted to sequence several mitochondrial DNA (mtDNA) loci using degenerate avian primers from Sorenson (2003) to determine if sufficient genetic variability was present for analysis. The control region, ATPase8, and ND2 were chosen, as these regions have proven useful in population genetic studies of birds (Burg & Croxall, 2001; Oyler-McCance *et al.*, 2005; Sorenson, 2003). We successfully amplified portions of the ND2, but were unable to amplify control region and ATPase8. Given the lower quality of DNA extracts from many samples we decided to amplify the gene in two pieces and run four sequencing reactions (both strands of each PCR product). The primers used and PCR conditions are described in Chapter 3.

We extracted all georeferenced *Gubernatrix cristata* records from different sources: fieldworking, museum skins, literature, and internet datasets. In total this dataset comprised 172 georeferenced records belonging to individuals, couples or a small number of birds seen on the area where the records were taken.

To model the species distributions, we selected 20 environmental predictors. We downloaded the altitude (in m) and the 19 bioclimatic predictors (1950 - 2000) of the WORLDCLIM dataset (<http://www.worldclim.org>) for South America at 30 arc-second resolution (~ 1km) (Hijmans *et al.*, 2005a).

BIOCLIM (using the DIVA-GIS platform) and MaxEnt were chosen for our two separate biogeoclimatic predictive models. DIVA-GIS is a GIS supported platform of the BIOCLIM model (Hijmans *et al.*, 2005b) which fits a minimal species habitat envelope in a multidimensional climate space. MaxEnt is a program developed for maximum entropy modeling of species geographic distributions (Phillips *et al.*, 2006) which expresses the suitability of each grid cell as a function of the environmental variables at that grid cell. High value of functions indicates predicted suitable conditions for a species. These models were selected to accommodate the presence-only data and span a range of modeling platforms (Elith *et al.*, 2006).

## 1.5. General Results

In bottlenecked populations, the observed gene diversity exceeds the expected equilibrium gene diversity under the assumption of mutation-drift equilibrium. The null hypothesis tested for heterozygosity excess using Wilcoxon sign-rank test provided ( $P < 0.05273$ ), ( $P > 0.09668$ ) and ( $P < 0.42285$ ) probabilities under the IAM, TPM, and SMM respectively. The null hypothesis is accepted under the SMM model only, implying that the *Gubernatrix cristata* has not experienced any recent genetic bottlenecks. The estimated values of the heterozygosity excess and their probabilities in sign test were 5.89 ( $P < 0.15033$ ) for the IAM, 5.90 ( $P < 0.15148$ ) for the TPM and 5.91 ( $P < 0.38936$ ) for SMM. The null hypothesis of mutation-drift equilibrium was not also accepted based on the sign test under mutation models.

Only 67 samples were successfully sequenced for the target ND2 fragment (sequences were deposited at GenBank under accession nos **tal e tal**). A total of 349 bp were aligned in *G. cristata* samples contained two variable and parsimony informative sites and a total of three haplotypes (H1 – H3). Among H2 and H1 haplotypes 75 mutational steps were found whilst 34 among H2 and H3. Measures of genetic variability as haplotype diversity ( $Hd$ ), nucleotide variation estimated based on the number of segregating sites ( $\pi$ ) and mean number of pairwise differences for sampled populations ( $k$ ) are shown in Chapter 3. Estimates of nucleotide and haplotype diversity in the sample were relatively high, with  $\pi = 0.00277$  and  $Hd = 0.6219$ . The network revealed no structure pattern at all. The  $F_{st}$  (0,00340) and  $Nm$  (73,18) shown a weak structuring with little genetic differentiation. Lack of support to a demographic expansion hypothesis was either obtained by different tests made to test the neutral theory of molecular evolution: Fu and Li's  $D^*$  test statistic (0,71916) showed no statistical significance ( $P > 0.10$ ) as well not significant was Fu and Li's  $F^*$  test statistic (1,34814) ( $P > 0.10$ ). Fu's  $F_s$  statistic was 2,248 which was significantly different from zero and is a likely evidence indicating a decrease in population size and/or balancing selection. Tajima's  $D$  was positive (2,17506) what signifies low levels of both low and high frequency polymorphisms, indicating a decrease in population size and/or balancing selection.

As the exact geographical origin of many (45%) of the sampled Yellow cardinal individuals is unknown or uncertain (many of the individuals sampled in zoos or at privates, or got from police seizures were not associated to reliable information on their exact location of capture), the structure analysis would be particularly relevant because it would allow clustering of the individuals without the need for a priori geographical information. Five independent runs for one

to six populations using an admixture model and correlated allele frequencies among populations (Falush *et al.* 2003), were run in the program Structure. The results for all runs (shown in Chapter 3) were practically the same suggesting no clustering of individuals. Due to this result no individual assignment to any geographical area was possible.

The mismatch distribution of observed haplotype variation was unimodal revealing a departure from expected variation under a Constant population size ( $k = 0,965$ , variance of  $k = 0,7205$ , Raggedness statistic  $r = 0,1319$ ).

The maps with a detailed discussion about our results on Species Distribution Model is found in Chapter 4.

Upon execution of this project we advance the knowledge about *Gubernatrix cristata*, especially with regard to their population status within its area of distribution. We were able to clearly determine where the species still survives in Brazil and which are in fact the best places to invest in its conservation. Upon execution of this project, we defined actions for the management of the species in captivity. Partnerships were established and new contacts enabled the centralization of all seizures from Police in RS in one place: the Gramado Zoo. This was the first step towards establishing a conservation breeding program, as well as a national sanitary protocol and engagement of public and private institutions and NGOs for the conservation of *G. cristata*. One of the most significant impacts of the project was to encourage the creation of the Yellow Cardinal Action Plan. Once there isn't an effective project for the conservation of an endangered species there is no way to define actions on the part of ICMBio. While the action plan has not yet been published, some of these actions have already been implemented and others are being held. Yellow Cardinal Project began in 2005 as a pilot project in the region of Pinheiro Machado and Pedras Altas. It took shape in 2006 and 2007 with the support of O Boticario Foundation for Nature Protection. And will continue for many years ahead, because many questions remain to be answered.



Figura 1.1. Macho de *Gubernatrix cristata*. Departamento Lavalleja, Uruguay em 19.03.2009. Foto de Claiton Martins-Ferreira  
Figure 1.1. *Gubernatrix cristata* male. Lavalleja Department, Uruguay, in 19.03.2009. Photo by Claiton Martins-Ferreira




Figura 1.2. Ovos de *Gubernatrix cristata* encontrados em um ninho ativo. Departamento Lavalleja, Uruguay. Foto de Alvaro Riccetto.  
Figure 1.2. *Gubernatrix cristata* eggs found in an active nest in Lavalleja Department, Uruguay. Photo by Alvaro Riccetto.



Figura 1.3. Ninho de *Gubernatrix cristata* encontrado abandonado e coletado. Departamento Lavalleja, Uruguay em 22.01.2007. Foto de Claiton Martins-Ferreira  
Figure 1.3. *Gubernatrix cristata* nest found abandoned and collected in Lavalleja Department, Uruguay in 22.01.2007. Photo by Claiton Martins-Ferreira

## CAPÍTULO DOIS

### Isolation and characterization of 10 microsatellite loci in the Yellow Cardinal *Gubernatrix cristata* En



Article published on Molecular Ecology Resources (2010) 10(4): 751-754  
<http://dx.doi.org/10.1111/j.1755-0998.2010.02871.x>

## Isolation and characterization of 10 microsatellite loci in the Yellow Cardinal *Gubernatrix cristata* En

C. Martins-Ferreira<sup>1\*</sup> · O. Haddrath<sup>2</sup> · A. J. Baker<sup>2</sup> · T. R. O. Freitas<sup>1</sup>

<sup>1</sup> PPG Biologia Animal, Departamento de Zoologia/UFRGS, 91501-970 Porto Alegre, RS, Brazil

\*e-mail: gcristata@gmail.com

<sup>2</sup> Natural History Department, Royal Ontario Museum, 80139 Toronto, ON, Canada

**Abstract** *Gubernatrix cristata* is a rare and endangered bird species from the South American Pampas grassland with a restricted and patchy distribution. Here we report the development of ten microsatellite markers isolated from this species and the characterization of their allele variability. The number of alleles observed for each locus ranged from 4 to 14, with an average of 7.5 alleles per locus. The microsatellites proved to be useful in revealing levels of diversity in *G. cristata*, and thus can be used to explore the genetic structure of scattered populations across its present geographic range.

**Keywords** *Gubernatrix cristata* · Genetic diversity · Microsatellites · Endangered species · Pampas grassland · Mitochondrial DNA ·

### Introduction

*Gubernatrix cristata*, the Yellow Cardinal, is a rare bird species from the Pampas grassland. It has a restricted geographical distribution in southern South America and is unique to this Biome (Uruguay, Argentina, and Southern Brazil). This species has undergone a dramatic population decline across its range and its conservation status is defined as endangered (Fontana *et al.* 2003). The remaining populations are small and isolated (BIRDLIFE INTERNATIONAL, 2000, 2009). The destruction and fragmentation of its habitat have been enumerated as possible additional threats for the species, as well as the extreme pressure of capture to supply the traffic in wild animals (Chebez 1994; BIRDLIFE INTERNATIONAL, 2000, 2009). The aim of this study was to

develop a set of polymorphic microsatellite markers (*Simple Sequence Repeat – SSR*) for *G. cristata* to describe the genetic structure of these endangered populations. These are the first microsatellites isolated for the Yellow Cardinal. Thirteen field trips were made from April 2006 to May 2008 in an attempt to find individuals and collect samples for genetic analysis. In Rio Grande do Sul (the southernmost Brazilian state), just three Yellow Cardinals were found in a single locality (Espinilho State Park). In Uruguay, three separate field trips to different localities located only five individuals at a farm in Minas, Lavalleja Department. In Argentina, a visit was made to the *Museo de Ciencias Naturales Bernardino Rivadavia* from which tissue samples of eight skins were collected, one from each province where

the species has occurred, and one from Uruguay. The other samples came from specimens seized by the Environmental Police and are of unknown origin.

## Material and Methods

Total DNA was extracted from 42 fresh samples of feathers or blood stored on FTA Cards (Whatman) of *G. cristata*, (using methods in Medrano *et al.* 1990; Eguchi and Eguchi, Y. 2000; Bello, N. *et al.* 2001), and from 13 museum skin samples (using methods in Mundy N. I. *et al.* 1997). Microsatellites were identified from genomic libraries partially enriched for the nucleotide repeats (CT)<sub>n</sub> and (GT)<sub>n</sub> using biotinylated oligonucleotide sequences linked to magnetic particles and recovered by streptavidin (KIJAS *et al.*, 1994; BILLOTE *et al.*, 1999). DNA fragments enriched with microsatellites were ligated in pGEM-T Easy vector (Promega), as described by the manufacturer, and used to transform cells of *E. coli* XL1Blue competent (Stratagene). A total number of 96 colonies were obtained and sequenced using the BigDye v3.1 Terminator kit and an ABI 3100 Genetic analyzer (Applied Biosystems). From these sequences, 40 clones were found to contain SSR motifs. Following some simple rules to design efficient PCR primers (WEISING, K *et al.* 2005) not all positive clones were useful for primer design. Forward and reverse sequences were aligned in SeqMan (DNASTAR packet), and primers were designed for 23 loci using the software PRIMER3 ([http://frodo.wi.mit.edu/cgi-bin/primer3/primer3\\_www.cgi](http://frodo.wi.mit.edu/cgi-bin/primer3/primer3_www.cgi)). For each SSR locus, the forward primer was synthesized with

a 5' 19 bp long M13 tail (5'-CACGACGTTGTAA AACGAC-3') which functioned as an adaptor for a fluorescently labeled primer. Following the method of Schuelke (2000), three primer amplifications were performed using a locus-specific forward M13 tailed primer, a locus specific reverse primer, and a universal fluorescent-labelled M13 primer, labeled with either 6-FAM or VIC (Applied Biosystems). All the PCR amplifications were made under the following conditions: 94°C for 4 min, followed by 36 cycles of 94°C for 30 s, 54°C for 25 s, and 72°C for 30 s, with a final extension of 7 min to 72°C. The amplification volume was 12.5 µl and consisted of: 3 µl of DNA, 6.67 µl H<sub>2</sub>O, 1.25 µl buffer (10X:100 mM Tris-HCl, pH 8.3, 500 mM KCl, 25 mM MgCl<sub>2</sub>, 0.1% gelatin), 0.28 µl dNTPs (10 mM), 5 pmol reverse primer, 2.5 pmol forward primer, 5 pmol universal M13 primer, and 0.25 U Taq polymerase (Amersham Pharmacia Biotech). The amplified products were run on an ABI 3100 Genetic Analyzer along with the LIZ 500 size standard (Applied Biosystems). The STRand v3.7 software was used for peak scoring. ARLEQUIN 3.11 (EXCOFFIER *et al.* 2005) was used to calculate the observed (H<sub>O</sub>) and the expected (H<sub>E</sub>) heterozygosities, to test if there was deviation from Hardy-Weinberg equilibrium (HWE), and to test if there was linkage disequilibrium among all pairs of loci. MICRO-CHECKER (Van Oosterhout *et al.* 2004) was used to quantify the possible genotyping errors and null alleles. Ten SSRs were found to be polymorphic, although four of them presented null alleles according to



MICRO-CHECKER (GcrisC02, GcrisE02, GcrisF12, and GcrisH09) and ARLEQUIN (Table 1). In the remaining six loci, linkage disequilibrium among any pairs of loci, genotyping mistakes because of the presence of null alleles, allele dominance, or stuttering were not detected. The number of alleles per locus varied from 4 to 14 with an average of 7.5 alleles per locus. The observed heterozygosity (HO) varied from 0.126 to 0.893 and the expected (HE) heterozygosity from 0.130 to 0.930, with means and standard errors of  $0.6824 \pm 0.1144$  and  $0.7132 \pm 0.0272$  respectively (Table 1) (GenBank accession numbers: GU237061 to GU237070). The deviation from Hardy-Weinberg equilibrium presented by 4 loci may be the result of a sampling bias. Since the species is rare and most of the material used was from seizures made by the Environmental Police, we cannot determine if all of the samples represent the same population. Samples could represent several differentiated populations combined that may be the cause of departures from HWE. For the four loci that had null alleles and departed from HWE due to heterozygote deficiencies, we estimated null allele rates (Table 2). The high level of heterozygosity observed indicates that the species has not suffered any severe bottleneck, even though this conclusion should be regarded with caution because of the low sample size due to the rarity of the species in the wild, and also because of the low DNA quality in many of these samples due to the material collected. In terms of *ex situ* conservation, these data allow us to conclude that creating a *G.*

*cristata* captive breeding program would have a high probability of success since the species is not suffering from low levels of genetic diversity as shown by the observed heterozygosity. The primers proved to be useful in revealing levels of diversity in *G. cristata*, and can be used to explore the genetic structure of scattered populations across its present geographic range.

### Acknowledgements.

We thank to Dr. A. P. Souza for the training course in microsatellite isolation. We thank to P. Rorato and C. Lopes for helping us with the software packages and valuable discussions. We thank CNPq for providing the doctorate scholarship to C. Martins-Ferreira, to CAPES for the Sandwich Scholarship, and to FBPCN for financing the project. Laboratory work was funded by NSERC operating grant to AJB. We also thank Kleber Valenti Schenk for assistance in the translation the MS from Portuguese to English.

### References

- AZPIROZ, A. B. 2003. **Aves Del Uruguay:** Lista e introducción a su biología y conservación. Montevideo, Aves Uruguay-GUPECA.
- BELLO, N.; FRANCINO, O. & SANCHEZ, A. 2001. Isolation of genomic DNA from feathers. *J Vet Diagn Invest.*, **13**(2):162-4.
- BILLOTE, N.; RISTERUCCI, A-M; BAURENS, F-C. 1999. Microsatellite enriched libraries: applied methodology for the development of SSR markers in tropical crops. **Fruits** 54:277–288

- BIRDLIFE INTERNATIONAL. 2000. **Threatened birds of the world**. Barcelona and Cambridge/UK, Lynx Edicions and BirdLife International.
- BIRDLIFE INTERNATIONAL. 2009. **Species factsheet: *Gubernatrix cristata***. Downloaded from <http://www.birdlife.org> on 25/7/2009
- CHEBEZ, J. C. 1994. **Los que se van**. Buenos Aires, Albatros.
- EGUCHI, T. & EGUCHI, Y. 2000. High yield DNA extraction from the snake cast-off skin or bird feathers using collagenase. **Biotechnology Letters**, Netherlands, **22**: 1097–1100.
- EXCOFFIER L.; LAVAL, L.G.; SCHNEIDER, S. 2005. Arlequin ver. 3.0 An integrated software package for population genetics data analysis. **Evol Bioinform Online** 1:47–50
- FONTANA, C. S.; BENCKE, G. A. & REIS, R. E. 2003. **Livro vermelho da fauna ameaçada de extinção no Rio Grande do Sul**. Porto Alegre, EDIPUCRS, 632p.
- KIJAS, J.M.; FOWLER, J.C.; GARBETT, C.A.; THOMAS, M.R. 1994. Enrichment of microsatellites from the citrus genome using biotinylated oligonucleotide sequences bound to streptavidin-coated magnetic particles. **Biotechniques** 16:656–660
- MEDRANO, J.F., AASEN, E. & SHARROW, L. 1990. DNA extraction from nucleated red blood cells. **Biotechniques**, **8**: 43.
- MUNDY, N.I.; UNIT, P.; WOODRUFF, D.S. 1997. Skin from feet of museum specimens as a non-destructive source of DNA for avian genotyping. **The Auk**, 114(1):126-129
- SCHUELKE, M. 2000. An economic method for the fluorescent labelling of PCR fragments. **Nat Biotechnol** 18:233–234
- VAN OOSTERHOUT, C.; HUTCHINSON, W.F.; WILLS, D.P.M.; SHIPLEY, P. 2004. MICRO-CHECKER: software for identifying and correcting genotyping errors in microsatellite data. **Mol Ecol Notes** 4:535–538
- WEISING, K. et al. 2005. DNA Fingerprinting in Plants: principles, methods and applications, 2nd ed. Boca Raton, CRC Press, Taylor & Francis Group.

Table 1. *Gubernatrix cristata* microsatellite loci characteristics

Locus	Primer Sequences <sup>a</sup>	Repeat	Alleles	Size (bp)	$H_o$	$H_E$	PIC values <sup>b</sup>
GcrisC02	F: TCACCCCTTTGGTTGCTTAC R: TCAGGGGTGTGTCTCTCCTT	(CA)7	4	247	0,267	0,438*	0,373
GcrisC08	F: AGCCACTCATCCGATTGTT R: AGTCTTGGCTGCCTGTCATT	(AC)8	6	174	0,911	0,791	0,748
GcrisE02	F: TTGGAGACCTTGTGGTGTG R: TCACACTGGGCACATGAGAT	(CA)8	4	227	0,229	0,710*	0,648
GcrisF02	F: TTCACTGGGTTACAGAAAGG R: GCTGTCGCATTGTTCAAGG	(CA)12	14	190	0,833	0,893	0,873
GcrisF12	F: AAGATGTGCCTTTGGTCTGG R: CCATTGCAGAAATGTCCTGA	(AC)9 (CA)7	6	230	0,600	0,784*	0,734
GcrisG10	F: CAGGATCCTCTGCCATGTCT R: TTTTCCTTTTAACGCCAAG	(CA)7	4	236	0,130	0,126	0,122
GcrisH06	F: GAGAGAAACCAGGTGCTTCG R: GTTTTAAGGCTGGGGACACA	(CA)9	10	230	0,735	0,830	0,799
GcrisH07	F: GTGTGACTTGTCCCCTTCC R: ACAGGAGCAGCCAGTTGAAT	(CA)7 (AC)5	6	213	0,739	0,755	0,709
GcrisH09	F: CAAGGTGTTGTGGAGCCTTT R: GCAACCAGCACATGAAATTG	(CA)7	10	236	0,702	0,834*	0,802
GcrisH12	F: AAGCGTGACCATGAAAAATGT R: TTCATCTGCCCTCCTTGTTT	(CA)7	5	249	0,232	0,700	0,637

<sup>a</sup> All forward primers were M13-tailed at the 5' end. Significant departures from HWE: \*  $P < 0.001$

<sup>b</sup> Polimorphism Information Content

Table 2. Estimates of null allele rates of four loci that departed from HWE using three methods of null allele estimation.

Locus	Null Present	Oosterhout	Chakraborty	Brookfield 1
C2	yes	0,2512	0,4278	0,1662
E2	yes	0,3045	0,4823	0,2682
F12	yes	0,1072	0,1248	0,0966
H9	yes	0,0694	0,0751	0,0631

## CAPÍTULO TRÊS

### **Lack of population structure in the Yellow Cardinal *Gubernatrix cristata* En and implications to its conservation**



Article to be submitted to Animal Conservation Journal.

## Lack of structuring and population shrinkage in the endangered yellow cardinal (*Gubernatrix cristata*): implications for conservation

Claiton Martins-Ferreira<sup>a,\*</sup>, Allan J. Baker<sup>b</sup>, Thales R. O. Freitas<sup>a,c</sup>

<sup>a</sup>PPG Biologia Animal, Departamento de Zoologia/UFRGS, Porto Alegre, RS, Brazil

<sup>b</sup>Natural History Department, Royal Ontario Museum, 80139 Toronto, ON, Canada

<sup>c</sup>Departamento de Genetica/UFRGS, Porto Alegre, RS, Brazil

\*E-mail address: gcristata@gmail.com

### Abstract

*Gubernatrix cristata*, the Yellow Cardinal, is a rare bird species from the Pampas grassland. It has a restricted geographical distribution in southern South America and is unique to this Biome (Uruguay, Argentina, and Southern Brazil). Fourteen field trips were made from April 2006 to March 2009 in an attempt to find individuals and collect samples for genetic analysis. A total of 72 yellow cardinal samples was taken in this study, 59 were from contemporary specimens and 13 samples were from museum specimens. We assessed the species genetic diversity through ten polymorphic nuclear microsatellite loci of *Gubernatrix cristata* and a ND2 mtDNA fragment. We found only three haplotypes with nucleotide and haplotype diversity equals to  $\pi = 0.00277$  and  $Hd = 0.6219$ . The  $F_{st}$  (0,00340) and  $Nm$  (73,18) shown a weak structuring with little genetic differentiation. Fu's  $F_s$  statistic was significantly different from zero (2,248) and Tajima's  $D$  was positive (2,17506). Both tests showed likely evidence indicating a decrease in population size and/or balancing selection. Analyses using the "no admixture model" and a larger burn-in (500000) yielded no clustering of individuals. Due to this result no individual assignment to any geographical area was possible. The present study assists in understanding the conservation needs for Yellow Cardinal by providing information on the genetic diversity and population structuring along its distribution range. Although we found no structuring within our study area, further study is needed, examining the genetic diversity and population structuring throughout the species' range, adding more samples from wild animals from La Pampa population and Corrientes.

Keywords: Endangered species, Genetic diversity, *Gubernatrix cristata*, Mitochondrial DNA, Microsatellites, ND2 gene, Pampas grassland.

## Introduction

*Gubernatrix cristata*, the Yellow Cardinal, is a rare bird species from the Pampas grassland. It has a restricted geographical distribution in southern South America and is unique to this Biome (Uruguay, Argentina, and Southern Brazil) (Azpiroz, 2003). This species has undergone a dramatic population decline across its range and its conservation status is defined as endangered (BirdLife International, 2000; Fontana *et al.* 2003). In Argentina, it is now rare except very locally (Fraga, 1997). In Uruguay, it was historically known from 13 departments, but recently from only Paysandú, Río Negro, Florida and Rocha (BirdLife International, 2000; Azpiroz, 2003). In Southern Brazil (Rio Grande do Sul State) it was already considered rare in the 1970s and 1980s (Belton, 1985). Since then it suffered a dramatic decline and now is believed it is confined to restricted spots in the SE hills, near Uruguayan border and at W tip of Rio Grande do Sul (Fontana *et al.* 2003). Once the lack of yellow cardinal sightings in the last few years and its bad situation in the adjacent areas in Uruguay, the SE hills population seems to be isolated or even extinct. (Martins-Ferreira, 2007). The remaining populations are small and isolated (BirdLife International, 2000). The only breeding wild population known in Brazil of *G. cristata* seems to be limited to Espinilho State Park (*Parque Estadual do Espinilho*) and surroundings (Damiani *et al.*, 2009). The destruction and fragmentation of its habitat have been enumerated as possible additional threats for the species, as well as the extreme pressure of capture to supply the traffic of wild animals (Chebez 1994; BirdLife International, 2000). The main reason behind the traffick is that bird is a colorful and singing passerine. The male has a bright yellow chest and back with black crest, eye line and throat. It has the full eyebrow and malar stripe yellow while the female are white with a gray breast (Belton, 1994; Sick, 1997). During the breeding time the male defends its territory aggressively attacking other males. That behavior turns to be a way how poachers capture them using a captive male as decoy (Fontana *et al.* 2003). Hudson (1920) already atested that its qualities as beautiful bird with loud and musical voice made it a favourite cage bird. The growing record of hybrids among yellow cardinal and common diuca finch (*Diuca diuca*) in Argentina seems to be a response to the shortage of male individuals in the population of the species (Bertonatti & Guerra, 1997, 2001). With the null hypothesis that Yellow Cardinal has not a population structure throughout its distribution range, we decide to investigate that. Our aim was to determine the population status of *Gubernatrix cristata* across its distribution range, taking in consideration the population structure, genetic diversity and gene flow. Only with such

information in hand we would be able to propose a specific conservation strategy to the species.

## **Methods**

### **Sampling**

Fourteen field trips were made from April 2006 to March 2009 in an attempt to find individuals and collect samples for genetic analysis. In Rio Grande do Sul (the southernmost Brazilian state), just four Yellow Cardinals were found in a single locality (Espinho State Park). In Uruguay, four separate field trips to different localities found only six individuals in Minas, Lavalleja Department, five in one farm and one in another land. All of them were ring banded with rings provided by CEMAVE/ICMBio (National Research Center for the Conservation of Wild Birds). In Argentina, a visit was made to the Museo de Ciencias Naturales Bernardino Rivadavia from which tissue samples of eight skins were collected, one from each province where the species has occurred, and one from Uruguay. Five more samples came from museum specimens: one from *Museu de Zoologia da Universidade de São Paulo* (MZUSP, São Paulo, Brazil), two from *Museu de Ciências Naturais da Fundação Zoobotânica do Rio Grande do Sul* (MCN/FZBRS, Porto Alegre, Brazil) and two from Royal Ontario Museum (ROM, Toronto, Canada) (Fig. 1). The rest of the samples came from specimens seized by the Environmental Police and are of unknown origin. A total of 97 yellow cardinal samples was taken in this study (Appendix A.1). Some of which having been maintained (but not bred) in captivity by zoos, conservation institutions, or private individuals. Of these, 25 presented a poor DNA quality that wasn't possible to proceed with any analysis. From 72 samples that worked properly 59 were from contemporary specimens. The other 13 samples were from museum specimens (years 1905–1975) selected from across the range, but with a special emphasis on the population in Argentina since many samples with unknown geographical data were supposed to come. Blood samples were stored in FTA Cards (Whatman) and feathers were placed in 95% ethanol solution. Museum tissues samples had their DNA extracted using the DNeasy Tissue Kit (QIAGEN) with the appropriate protocol according to the manufacturer for each type of



tissue (Mundy, N. I. *et al.*, 1997). Contamination with modern DNA or PCR products was monitored by including two extraction blanks in every extraction round and prevented by performing all museum samples extractions in a dedicated 'clean' laboratory, kept free of good quality DNA and PCR products.




Figure 1. Map showing the points of collected samples from wild individuals or from museum skins across *G. cristata* distribution range. Polygon used with permission from Birdlife.

### Laboratory Analyses

A microsatellite-enriched genomic DNA library of *Gubernatrix cristata* was constructed and ten polymorphic nuclear microsatellite loci (with 2-bp repeats) were selected on the basis of polymorphism levels and size of amplification products. All

samples were genotyped using a polymerase chain reaction targeting different loci (Table 1). Primers and PCR conditions are given in Molecular Ecology Resources Primer Development Consortium (2010). PCR products were purified, cycle sequenced on both strands, and run on an ABI 3730 sequencer.

We extracted DNA from feather samples using the Alkali protocol for DNA extraction and attempted to sequence several mitochondrial DNA (mtDNA) loci using degenerate avian primers from Sorenson (2003) to determine if sufficient genetic variability was present for analysis. The control region, ATPase8, and ND2 were chosen, as these regions have proven useful in population genetic studies of birds (Burg and Croxall, 2001; Sorenson, 2003; Oyeler-McCance *et al.*, 2005). We successfully amplified portions of the ND2, but were unable to amplify control region and ATPase8. Given the lower quality of DNA extracts from many samples we decided to amplify the gene in two pieces and run four sequencing reactions (both strands of each PCR product). The primers used were *ND2H* 5' CCT TGA AGC ACT TCT GGG AAT CAG A 3' (Tavares *et al.*, 2006) - *ND2F\_Passerine* 5' CCA YCC ACG AGC YAT TGA AGC 3' (new primer designed), and *MetLTF* 5' AAG CTA TCG GGC CCA TAC CCG 3' (Tavares *et al.*, 2006) - *ND2R\_Passerine* 5' GCC ATG CRT TGG TYA TGC TNG AG 3' (new primer designed). Those primers were used to amplify a 349 base-pair region that contained the variable sites in ND2. We amplified DNA via the polymerase chain reaction (PCR) in 12,5 µl reaction volumes containing 2.0 µl DNA, 7.92 µl water, 0.5 µl (10 pmol) of each primer, and 1.58 µl PCR master mix, consisting of a 1.25 µl buffer (10X:100 mM Tris-HCl, pH 8.3, 500 mM KCl, 25 mM MgCl<sub>2</sub>, 0.1% gelatin), 0.28 µl dNTPs (10 mM), and 0.05 U Taq polymerase (Amersham Pharmacia Biotech). Reactions were performed under the following conditions: initial denaturation at 94°C for three minutes, followed by 36 cycles of 94°C for 45 sec, 50°C for 45 sec, and 72°C for 1 minute and 30 sec, and a final extension of 72°C for 7 minutes. The PCR products were then electrophoresed on a 1% agarose gel containing ethidium bromide and visualized under UV light. Amplified segments were purified by excising bands from agarose gels and centrifuging each through a filter tip. The sequencing reactions were then electrophoresed on an ABI Prism 3100 automated sequencer (Applied Biosystems). We sequenced in both directions and sequences were aligned and assembled using ChromasPro V. 1.49 (Technelysium Pty Ltd) and aligned using the CLUSTAL W algorithm with default

options, implemented in MEGA 4.0.2 (Tamura *et al.*, 2007). Alignments were checked and edited by hand when necessary. The sequences were deposited in GenBank under accession numbers **tal and tal**.

Table 1. *Gubernatrix cristata* microsatellite loci used in this study.

Locus	Repeat	Alleles	Size (bp)	$H_o$	$H_E$	GenBank no.
GcrisC02	(CA)7	4	247	0,267	0,438*	GU237061
GcrisC08	(AC)8	6	174	0,911	0,791	GU237062
GcrisE02	(CA)8	4	227	0,229	0,710*	GU237063
GcrisF02	(CA)12	14	190	0,833	0,893	GU237064
GcrisF12	(AC)9 (CA)7	6	230	0,600	0,784*	GU237065
GcrisG10	(CA)7	4	236	0,130	0,126	GU237066
GcrisH06	(CA)9	10	230	0,735	0,830	GU237067
GcrisH07	(CA)7 (AC)5	6	213	0,739	0,755	GU237068
GcrisH09	(CA)7	10	236	0,702	0,834*	GU237069
GcrisH12	(CA)7	5	249	0,232	0,700	GU237070

Significant departures from HWE: \*  $P < 0.001$

## Statistical Analyses

We assessed the bottleneck history of these populations using Bottleneck v.1.2.02 (Cornuet & Luikart, 1996). This program was used to detect a heterozygote excess for individual populations, considering the two-phased model (TPM) of microsatellite mutation, a 70% stepwise-mutation model (SMM) and 30% infinite alleles model (IAM), and 1000 replications. Several other combinations of the SMM:IAM ratio were tested to establish the sensitivity of these data to the mutational mechanism. The Wilcoxon signed-rank test was used to determine if the allele frequency distribution for a population exhibited significant heterozygote excess relative to model expectations. Bayesian clustering with software Structure v.2.3.2 (Pritchard *et al.*, 2000) was used to assign individuals to populations (K) based on posterior probabilities where K is unknown. The number of groups was set to 1–7 with 3 runs per K. Posterior probabilities were calculated for all K hypothetical populations.

All analyses were based on 1,000,000 Markov Chain Monte Carlo iterations following a burn in of 50,000 iterations.

To measure mtDNA diversity, both haplotype diversity,  $H_d$ , and nucleotide diversity,  $\pi$ , and their standard deviations were estimated using DnaSP v5.10.00 (Librado & Rozas, 2009). To test for evidence of recent population expansion we calculated Fu's  $F_s$  (Fu, 1997) and Fu and Li's (1993)  $D^*$  and  $F^*$  statistics to compare with Fu's  $F_s$ . Thus, if  $F_s$  is significant and  $F^*$  and  $D^*$  are not, it is an indication of population expansion, while the opposite indicates selection (Fu, 1997). We also calculate Tajima's  $D$  (Tajima, 1989). We used ARLEQUIN 3.11 (Schneider et al., 2000) and DnaSP v5.10.00 (Librado & Rozas, 2009) to perform these calculations. We created a mismatch distribution of pairwise differences using DnaSP v5.10.00 (Librado & Rozas, 2009) to compare the expected distribution for a population. The distribution tends to be multimodal when populations are at equilibrium and unimodal in cases of recent demographic expansion or reduction (Rogers and Harpending, 1992). To graphically display the observed mismatch distribution compared to the expected distributions for populations in equilibrium and expansion, we used Roger's method of moments (Rogers, 1995) as calculated in DNASP. The topological relationship between the haplotypes was estimated using the program NETWORK 4.5.1.0 (<http://www.fluxus-engineering.com>) with median joining approach, for the three data sets (Bandelt *et al.*, 1999).

## Results

### Genetic Variation

Allelic variation for the ten nuclear microsatellite loci ranged between 4 and 14 alleles with an average of 7.5 alleles per locus. The observed heterozygosity ( $H_o$ ) varied from 0.126 to 0.893 and the expected ( $H_e$ ) heterozygosity from 0.130 to 0.930, with means and standard errors of  $0.6824 \pm 0.1144$  and  $0.7132 \pm 0.0272$  respectively (Table 1). Four loci exhibited a departure from Hardy-Weinberg expectation that may be the result of a sampling bias. Since the species is rare and most of the material used was from seizures made by the Environmental Police (Table 2), we cannot determine if all of

the samples represent the same population.

Table 2. Geographical origin and source of the samples processed in our study.

Country	Samples	Source	Samples
Argentina	8	Captive Bred	27
Brazil	16	Police Seizure	27
Uruguay	16	Wild	5
Unknown	32	Museum Skin	13
Total	72	Total	72

Results of bottlenecks detection using Wilcoxon signed-rank and sign tests under Infinite Allele Model (IAM), Two-Phase mutation (TPM) and Step-wise mutation (SMM) are presented in Table 3. In bottlenecked populations, the observed gene diversity exceeds the expected equilibrium gene diversity under the assumption of mutation-drift equilibrium. The null hypothesis tested for heterozygosity excess using Wilcoxon sign-rank test provided ( $P < 0.05273$ ), ( $P > 0.09668$ ) and ( $P < 0.42285$ ) probabilities under the IAM, TPM, and SMM respectively. The null hypothesis is accepted under the SMM model only, implying that the *Gubernatrix cristata* has not experienced any recent genetic bottlenecks. The estimated values of the heterozygosity excess and their probabilities in sign test were 5.89 ( $P < 0.15033$ ) for the IAM, 5.90 ( $P < 0.15148$ ) for the TPM and 5.91 ( $P < 0.38936$ ) for SMM. The null hypothesis of mutation-drift equilibrium was not also accepted based on the sign test under mutation models.

Not all of the 97 sampled individuals were successfully sequenced for the target ATPase 8 or Control Region fragment. Only 67 were successfully sequenced for the target ND2 fragment (sequences were deposited at GenBank under accession nos [table](#)). A total of 349 bp were aligned in *G. cristata* samples contained two variable and parsimony informative sites and a total of three haplotypes (H1 – H3). Among H2 and H1 haplotypes 75 mutational steps were found whilst 34 among H2 and H3 (Fig. 2). Estimates of nucleotide and haplotype diversity in the sample were relatively high, with  $\pi = 0.00277$  and  $Hd = 0.6219$ . The genealogical relationships between haplotypes were estimated in the median-joining networks illustrated in Figure 2. The network revealed no structure pattern at all. Haplotype 2 had the highest probability of represent ancient forms because of its central position. The  $F_{st}$  (0,00340) and  $Nm$  (73,18) shown a weak

structuring with little genetic differentiation. The  $F_{st}$  analysis was done following Nei's (1978) classification where a value is considered low when  $F_{st} < 0,05$ ; medium when  $0,05 < F_{st} < 0,15$ , and high when  $F_{st} > 0,15$ . Lack of support to a demographic expansion hypothesis was either obtained by different tests made to test the neutral theory of molecular evolution: Fu and Li's  $D^*$  test statistic (0,71916) showed no statistical significance ( $P > 0.10$ ) as well not significant was Fu and Li's  $F^*$  test statistic (1,34814) ( $P > 0.10$ ). Fu's  $F_s$  statistic was 2,248 which was significantly different from zero and is a likely evidence indicating a decrease in population size and/or balancing selection. Tajima's  $D$  was positive (2,17506) what signifies low levels of both low and high frequency polymorphisms, indicating a decrease in population size and/or balancing selection.

Table 3. Genetic bottleneck detection using Wilcoxon signed-rank sign tests under Infinite Allele (IAM) , Two-Phase (TPM), and Step-wise (SMM) mutations models of microsatellite evolution

Test	Model		
	IAM	TPM	SMM
<i>Wilcoxon signed-rank test:</i>			
Probability of Heterozygosity excess	$P > 0.05273$	$P > 0.09668$	$P < 0.42285$
<i>Sign test:</i>			
Number of loci with heterozygosity excess			
Observed	8	8	5
Expected (probability)	5.89	5.90	5.91
Probability	$P > 0.15033$	$P > 0.15148$	$P > 0.38936$




Figure 2. Genealogical relationships between haplotypes estimated in the median-joining networks. H\_1, H\_2 and H\_3 – Haplotypes; 75 – mutational steps between H2 and H1; 34 – mutational steps between H2 and H3.

### Population Structure


As the exact geographical origin of many (45%) of the sampled Yellow cardinal individuals is unknown or uncertain (many of the individuals sampled in zoos or at privates, or got from police seizures were not associated to reliable information on their exact location of capture), the structure analysis would be particularly relevant because it would allow clustering of the individuals without the need for a priori geographical information. Five independent runs for one to six populations using an admixture model and correlated allele frequencies among populations (Falush *et al.*, 2003), were run. Figure 3 shows the results for  $k$  equals to 3 populations as bar plot and triangle plot. Analyses using the “no admixture model” and a larger burn-in (500000) yielded very similar results (data not shown). The results for all runs were practically the same suggesting no clustering of individuals (Fig. 2). Due to this result

no individual assignment to any geographical area was possible.

The mismatch distribution of observed haplotype variation was unimodal revealing a departure from expected variation under a Constant population size (Fig. 3) ( $k = 0,965$ , variance of  $k = 0,7205$ , Raggedness statistic  $r = 0,1319$ ).



a) Bar plot showing no structure for  $k = 3$  populations.



b) Triangle plot showing no structure for  $k = 3$  populations.

Figure 3. Results of Structure run. Here is depicted when  $k = 3$  populations. a) bar plots and b) triangle plot.






Figure 4. Mismatch distribution of the observed haplotype variation from 67 Yellow Cardinal mtDNA samples compared to the expected under a population expansion.

## Discussion

Endangered birds usually have small and fragmented populations. A commonly concern about population studies is a proper sample size to rightly infer genetic diversity measures. Despite the relatively small sample size of this study it has been recommended that at least 20 to 30 individuals be sampled in microsatellite studies (Pruett and Winker, 2008). Our 72 sample size is more than enough to run studies about genetic diversity and population structure estimations. The findings about no structuring appear to be related to other issues.

The lack of population structure showed by microsatellites with low  $F_{st}$  and large  $Nm$  apparently tell us the history about only one huge population across Yellow cardinal range with an intense gene flow. The same is truth for ND2 mitochondrial DNA gene.

The fact that both Fu's  $F_s$  and Tajima's  $D$  indicates a decrease in population size and/or balancing selection shows us a possible explanation of what is happening.

The yellow cardinal has suffered an intense hunting pressure by almost one hundred years now (Hudson, 1920). This time period is not sufficient to be detected at

the molecular level, however is enough to drastically reduce the species population size or even driven to extinct some subpopulations (Martins-Ferreira, 2007).

The species had a wide distribution area in the past that has been reduced over time, both due to changes in habitat, as the hunting pressure. Today it seems to be limited to a narrow continuous irregular arc-shaped band in Argentina, from San Luís, passing through La Pampa, Córdoba, Santa Fé, Entre Ríos and Corrientes (Marchiori, 2004; Martins-Ferreira, 2007). Following the center south of Uruguay to the west end of Rio Grande do Sul, Brazil. That reduction in its distribution range to only the optimal environment for the species, also ended up reducing their genetic variability and, possibly, the haplotype diversity. The Whooping Crane (*Grus americana*) experienced a severe decline in their population size in early 1900. Reducing its population to 14 individuals in 1938, resulted in the loss of two thirds of its haplotype diversity. Only three haplotypes were identified in the current population (Glenn *et al.*, 1999).

As shown in Whooping Cranes, a severe reduction in population size can cause a genetic bottleneck, resulting in loss of genetic diversity (Nei *et al.*, 1975). However, the impact of a bottleneck on population genetic diversity depends on how rapidly the population declines, the size of the bottleneck population and the duration of the bottleneck. When a population suffers a genetic bottleneck, rare alleles initially are lost, but the population can maintain diversity if the population recovery is rapid (Allendorf , 1986). If population abundance remains low, diversity can continue to be lost due to genetic drift (Allendorf ,1986). There are several possible explanations for the relatively high genetic diversity observed in Yellow Cardinals along Southern South America. First, the Yellow Cardinal decline may have been less severe than that of the Whooping Crane. The population status in the early 1900s is unknown, but never drop low before than today. Secondly, population recovery may have been sufficiently rapid so as not to have lost diversity due to genetic drift. Although Yellow Cardinal in Uruguay and Brazil has recently experienced several episodes of population decline, each time population recovery appears to have been rapid. Finally, recolonization by individuals that emigrated from areas where genetic diversity remained high, such as Argentina, could have contributed to the high genetic diversity. The observed genetic similarity within the region is consistent with an open population, as gene flow is expected to homogenize populations (Charlesworth, 2003). We detected no pattern in

the distribution of haplotypes to suggest that genetic structuring exists in Yellow Cardinals along its range.

Thus, our results support managing Yellow Cardinals along its distribution range as a single breeding population. It is possible that the Yellow Cardinal may form one large panmictic population. The existence of one large population could explain the high level of genetic diversity found in the population by allowing the effective population size to remain high during periods of population fluctuation and serve as a source of immigrants. Although the Yellow Cardinal's demographic history does not show drastic reductions in genetic diversity, the population size fluctuations appear to have had no detectable effects on the differentiation and frequency of haplotypes as evidenced by the mismatch distribution and  $F_s$ 's  $F_s$ . The minimum spanning network (Fig. 2) shows that the 3 haplotypes are not closely related (within 75 and 34 mutational steps). A triangle-shaped phylogeny, as shown in Fig. 2, tell us only that a population has experienced a severe decline on its genetic diversity. The Yellow Cardinal population still appears smaller than historical accounts suggest. Historical accounts described Yellow Cardinal as common in Argentina in the 1900s (Hudson, 1920; Wetmore, 1926). Thus, it seems that the population may have being suffering a decline in its numbers from the last century until now.

## **Conclusions**

The present study assists in understanding the conservation needs for Yellow Cardinal by providing information on the genetic diversity and population structuring along its distribution range.

Although we found no structuring within our study area, further study is needed, examining the genetic diversity and population structuring throughout the species' range, adding more samples from wild animals from La Pampa population and Corrientes.

The main implication for Yellow Cardinal conservation with these results is that a captive breeding program could be established on an international level, without much concern about preserving specific haplotypes, since all populations share the same ones.

## Acknowledgements

We thank to Oliver Haddrath for the tremendous help in conducting the lab work at ROM. We thank to Carla Lopes, Paula Rorato, and Nelson Fagundes for helping us with the software packages and valuable discussions. We thank to Darlise Lopes for the last-minute help in the lab. We thank CNPq for providing the doctorate scholarship to CMF, to CAPES for the Sandwich Scholarship, and to FBPCN for financing the project. Laboratory work was funded by NSERC operating grant to AJB.

## References

- Allendorf, F. W. (1986) Genetic drift and the loss alleles vs. heterozygosity. *Zoo Biology*, **5**, 181-190.
- Avise, J. C. (2000) *Phylogeography: the history and formation of species*. Harvard University Press, Cambridge, Massachusetts.
- Azpiroz, A. B. (2003) *Aves Del Uruguay: lista e introducción a su biología y conservación*. Aves Uruguay-GUPECA, Montevideo.
- Bandelt, H.J.; Forster, P. & Röhl, A. (1999) Median-joining networks for inferring intraspecific phylogenies. *Molecular Biology and Evolution*, **16**, 37-48.
- Belton, W. (1985) Birds of Rio Grande do Sul, Brazil. Part 2: Formicariidae through Corvidae. *Bulletin of the American Museum of Natural History*, **180**, (1): 1-242.
- Belton, W. (1994) *Aves do Rio Grande do Sul: distribuição e biologia*. Editora UNISINOS, São Leopoldo.
- Bertonatti, C. & Guerra, A. L. (1997) Hybridization between the yellow cardinal (*Gubernatrix cristata*) and the common diuca finch (*Diuca diuca*) in the wild in Argentina. *Hornero*, **14**,(4):235-42.
- Bertonatti, C. & Guerra, A. L. (2001) Nuevos registros de híbridos entre cardenal amarillo (*Gubernatrix cristata*) y diuca común (*Diuca diuca*). *Nuestras Aves*, **41**,27.
- BIRDLIFE INTERNATIONAL. (2000) *Threatened birds of the world*. Lynx Edicions and BirdLife International, Barcelona and Cambridge/UK.

- Burg, T. M. & Croxall, J. P. (2001) Global relationships amongst black-browed and grey-headed albatrosses: analysis of population structure using mitochondrial DNA and microsatellites. *Molecular Ecology*, **10**, 2647-2660
- Charlesworth, D. 2003. Effects of inbreeding on the genetic diversity of populations. *Philosophical Transactions of the Royal Society B: Biological Sciences*, **358**, 1051-1070.
- Chebez, J. C. (1994) *Los que se van*. Albatros, Buenos Aires.
- Chromas Pro. Available at <<http://www.technelysium.com.au/ChromasPro.html>> Downloaded in: [23/09/2009].
- Cornuet, J.M. & Luikart, G. (1996) Description and power analysis of two tests for inferring recent population bottlenecks from allele frequency data. *Genetics*, **144**:2001-2014
- Damiani, R.V.; Ribeiro,A.C. & Martins-Ferreira,C. (2009) Projeto Cardeal-Amarelo, resultados preliminares do monitoramento de *Gubernatrix cristata* no Parque Estadual do Espinilho – RS. In: CONGRESSO BRASILEIRO DE ORNITOLOGIA, XVII, 2009, Aracruz. Biogeografia das aves da Mata Atlântica. TEC ART, Aracruz/ES.
- Fontana, C. S.; Bencke, G. A. & Reis, R. E. (2003) *Livro vermelho da fauna ameaçada de extinção no Rio Grande do Sul*. EDIPUCRS, Porto Alegre.
- Fraga, R. M. (1997) Sección Aves. In: Garcia-Fernandez, J. J.; Ojeda, R. A.; Fraga, R. M.; Díaz, G. B. & Baigún, R. J. (Ed.). Libro rojo de mamíferos y aves amenazadas de la Argentina. FUCEMA, Buenos Aires.
- Fu, Y.-X. and Li, W.-H. (1993) Statistical tests of neutrality of mutations. *Genetics*, **133**, 693–709.
- Fu, Y.-X. (1997) Statistical tests of neutrality of mutations against population growth, hitchhiking and background selection. *Genetics*, **147**, 915–925.
- Glenn, T. C., W. Stephan and M. J. Braun. 1999. Effects of a population bottleneck on whooping crane mitochondrial DNA variation. *Conservation Biology*, **13**, 1097-1107.
- Hudson, W.H. (1920) *Birds of La Plata*. E. P. Dutton & Co., New York.
- Librado, P. & Rozas, J. (2009) DnaSP v5: A software for comprehensive analysis of DNA polymorphism data. *Bioinformatics*, **25**: 1451-1452.
- Marchiori, J. N. C. (2004) Fitogeografia do Rio Grande do Sul: *campos sulinos*. EST, Porto Alegre.
- Martins-Ferreira, C. (2007) Biogeografia e ecologia do Cardeal-Amarelo *Gubernatrix cristata* (Emberizidae) – resultados preliminares / Biogeography and ecology of Yellow Cardinal *Gubernatrix cristata* (Emberizidae) – preliminary results. In: Congresso Brasileiro de Ornitologia (15: 2007: Porto Alegre, RS) [programação e resumos]. EDIPUCRS, Porto Alegre.
- Molecular Ecology Resources Primer Development Consortium (2010) Permanent Genetic Resources added to the Molecular Ecology Resources Database 1 February 2010-31 March 2010. *Molecular Ecology Resources*, **10**, 751-754.

- Mundy, N. I., Unitt, P. & Woodruff, D. S. 1997a Skin from feet of museum specimens as a non-destructive source of DNA for avian genotyping. *Auk*, **114**, 126±129.
- Nei, M.; Maruyama, T. & Chakraborty, R. (1975) The bottleneck effect and genetic variability in populations. *Evolution*, **29**, 1–10.
- Oyler-McCance, S. J.; St. John, J.; Knopf, F. L. & Quinn, T. W. (2005) Population genetic analysis of mountain plover using mitochondrial DNA sequence data. *Condor*, **107**, 353-362.
- Pritchard, J.K.; Stephens, M. & Donnelly, P. (2000) Inference of population structure using multilocus genotype data. *Genetics*, **155**, 945–959.
- Pruett, C. L. & Winker, K. (2008) The effects of sample size on population genetic diversity estimates in song sparrows. *Journal of Avian Biology*, **39**, 252-256.
- Rogers, A.R. (1995) Genetic evidence for a Pleistocene population explosion. *Evolution*, **49**, 608-615.
- Rogers, A.R. & Harpending, H. (1992) Population growth makes waves in the distribution of pairwise genetic differences. *Molecular Biology and Evolution*, **9**, 552-569.
- Schneider, S.; Roessli, D. & Excoffier, L. (2000) Arlequin: a software for population genetics data analysis User manual ver 2.000. Genetics and Biometry Lab, Dept. of Anthropology, University of Geneva, Geneva.
- Sick, H. (1997) *Ornitologia brasileira*. Nova Fronteira, Rio de Janeiro.
- Sorenson, M. D. (2003) *Avian mtDNA primers*. Boston University. Available at <http://people.bu.edu/msoren/Bird.mt.Primers.pdf>, accessed 01 July 2008. Sutherland, W. J.; Newton, I. & Green, R. E. (2004) Bird ecology and conservation: *a handbook of techniques*. Oxford University Press, Oxford.
- Tajima, F. (1989) Statistical method for testing the neutral mutation hypothesis by DNA polymorphism. *Genetics*, **123**, 585–595.
- Tamura, K.; Dudley, J.; Nei, M. & Kumar, S. (2007) MEGA4: Molecular Evolutionary Genetics Analysis (MEGA) software version 4.0. *Molecular Biology and Evolution*, **24**:1596-1599.
- Tavares, E.S.; Baker, A.J.; Pereira, S.L. & Miyaki, C.Y. (2006) Phylogenetic relationships and historical biogeography of neotropical parrots (Psittaciformes: Psittacidae: Arini) inferred from mitochondrial and nuclear DNA sequences. *Systematic Biology*, **55** (3): 454-470.
- Wetmore, A. (1926) *Observations on the Birds of Argentina, Paraguay, Uruguay, and Chile*. Government Printing Office, Washington.

## CAPÍTULO QUATRO

**Potential Distribution Model to Yellow Cardinal  
*Gubernatrix cristata*: how much effort we should put to  
preserve this Endangered bird?**



Article to be submitted to Diversity and Distribution Journal.

## Potential Distribution Model to Yellow Cardinal *Gubernatrix cristata*: how much effort we should put to preserve this Endangered bird?

Claiton Martins-Ferreira<sup>a,\*</sup>, Thales R. O. Freitas<sup>a,b</sup>

<sup>a</sup>PPG Biologia Animal, Departamento de Zoologia/UFRGS, Porto Alegre, RS, Brazil

<sup>b</sup>Departamento de Genética/UFRGS, Porto Alegre, RS, Brazil

\*E-mail address: gcristata@gmail.com

### ABSTRACT

**Aim** The yellow cardinal distribution range map seems to be outdated because of the increasingly critical situation the species is facing in nature. Given the recent lack of knowledge regarding its distribution and its state of preservation in its natural habitat we decided to model the species potential distribution. The main idea behind this information was to propose strategies for their conservation. To better achieve that target we specifically focused our research on a) locating wild populations of this species in Rio Grande do Sul, Uruguay and Argentina, and b) identifying areas of occurrence of yellow cardinal that could be turned into protected areas.

**Location** Southern South America (Argentina, Uruguay and Southern Brazil)

**Methods** We extracted all georeferenced *Gubernatrix cristata* records from different sources: field working, museum skins, literature, and internet datasets. To model the species distributions, 20 environmental predictors were selected. To model the species distributions we selected the modeling application MaXent v. 3.3.2. Historical and current distribution maps of *G. cristata* were created using DIVA GIS.

**Results** The model that best predicts the relative index of environmental suitability for the species is the model where resulting AUC on test data was 0.868. The results of the jackknife test of variable importance showed that the environmental variable with highest gain when used in isolation is bio1\_sa\_30s\_cut (Annual Mean Temperature), which therefore appears to have the most useful information by itself. The environmental variable that decreases the gain the most when it is omitted is alt\_sa\_30s\_cut (Altitude), which therefore appears to have the most information that isn't present in the other variables.

**Main conclusions** If the objective of future surveys of the yellow cardinal in its distribution range is to detect other populations, then areas predicted to have high relative suitability in the spatial model is a good starting point for further targeted survey. The hunting pressure on the species is so great that the yellow cardinal nowadays seems to be only found in areas of difficult access. Many of the areas suitable for the species shown by the model may still contain populations and can also be used as key areas for future reintroductions into the yellow cardinal conservation program.

**Keywords** *Gubernatrix cristata*, Endangered species, Espinal, Pampa, Species Distribution Model.



## INTRODUCTION

The Yellow Cardinal, is a rare bird species from the Pampas grassland. It has a restricted geographical distribution in Southern South America and is unique to this Biome (Uruguay, Argentina, and Southern Brazil) (Azpiroz, 2003). This species has undergone a dramatic population decline across its range and its conservation status is defined as endangered (BirdLife International, 2000; Fontana *et al.*, 2003).

In Argentina, it is now rare except very locally (Fraga, 1997). It is found in San Luis, Buenos Aires, La Pampa and Río Negro (especially among General Conesa, San Antonio Oeste and Viedma). There are important populations in Corrientes (Pay Urbe and Estancia San Antonio), Entre Ríos (recently in the area of Montiel, Ceibas and Estancia La Choza) and Cordoba [recently in Chancaní (Chebez *et al.*, 1998)] (Di Giacomo, 2005). At the same time Cordoba present a huge lowering of Yellow Cardinal population, despite some new records (Kopta, 1999). There are records prior to 1975 from Salta, Tucumán, Santiago del Estero, Santa Fe and San Juan, and it was recorded with no details to Formosa, Chaco, La Rioja and (wrongly) in Misiones (Chebez, 1996).

In Uruguay, it was historically known from 13 departments, but recently from only Paysandú, Río Negro, Florida, Rocha, Lavalleja and Artigas (BirdLife International, 2000; Azpiroz, 2003; Lapitz, 2010; Martins-Ferreira, 2007). In 1999 the Uruguayan population of yellow cardinal was estimated less than 300 individuals, mostly concentrated in the Uruguay River Basin (Azpiroz *in litt.* 1999).

Two individuals collected in Paraguay in 1905 were probably caged birds escaped from captivity (Di Giacomo, 2005).

In Southern Brazil (Rio Grande do Sul State) it was already considered rare in the 1970s and 1980s (Belton, 1985). Since then it suffered a dramatically decline and now is believed it is confined to restricted spots in the SE hills, near Uruguayan border and at W tip of Rio Grande do Sul (Fontana *et al.*, 2003).

The constant and chronic exploitation as songbird for the cage bird market remains the most significant threat. The destruction and fragmentation of its habitat have been enumerated as possible additional threats for the species, as well as the

extreme pressure of capture to supply the traffic of wild animals (Chebez, 1994; BirdLife International 2000). The main reason behind the traffick is that bird is a colorful and singing passerine. In Rio Grande do Sul, Yellow Cardinal is caught mainly in the municipalities of Piratini and Pinheiro Machado. The mischaracterization and destruction of habitat have been listed as potential additional threats to the species (Chebez, 1994; BirdLife International, 2000). In Rio Grande do Sul, most of the original landscape of the south and west has been converted into rice fields and pastures for cattle. However, it is unknown to what degree these activities affected the species. Presumably its habitat suffered with conversion for livestock grazing, timber extraction for firewood and furniture (Chebez, 1994) and especially by the rapid forestation with *Eucalyptus* plantations.

The species distribution range map better known and used can be accessed on the website of Birdlife (2010). This map, however, seems to be outdated because of the increasingly critical situation the species is facing in nature. Given the recent lack of knowledge regarding its distribution and its state of preservation in its natural habitat we decided to model the species potential distribution.

The main idea behind this information was to propose strategies for their conservation. To better achieve that target we specifically focused our research on a) locating wild populations of this species in Rio Grande do Sul, Uruguay and Argentina, and b) identifying areas of occurrence of yellow cardinal that could be turned into protected areas.

## **METHODS**

### **Species Data**

We extracted all georeferenced *Gubernatrix cristata* records from different sources: field working, museum skins, literature, and internet datasets. Fourteen field trips were made from April 2006 to March 2009 in an attempt to find individuals. In Rio Grande do Sul (the Southernmost Brazilian State), just four yellow cardinals were found in a single locality (Espinilho State Park). In Uruguay, four separate field trips to different localities found only six individuals in Minas, Lavalleja Department, five in one farm and one in another land. All of them were ring banded with rings provided by

CEMAVE/ICMBio (National Research Center for the Conservation of Wild Birds) and have samples collected for genetic analysis. In Argentina, a visit was made to the *Museo de Ciencias Naturales Bernardino Rivadavia* to better know which localities yellow cardinals were found, the distribution range, and past records. Nine records came from Belton's field trips (Belton, 1985) to Southern Brazil. One more record came from *Museu de Zoologia da Universidade de São Paulo* (MZUSP, São Paulo, Brazil), and one from Royal Ontario Museum (ROM, Toronto, Canada). Some other records were extracted from Red Data Book (Collar *et al.*, 1992) and IBAS from Argentina (Di Giacomo, 2005). The remainder records were accessed through ORNIS data portal. In total this dataset comprised 172 georeferenced records belonging to individuals, couples or a small number of birds seen on the area where the records were taken. To avoid duplicate location names and confusion about co-ordinate system, a data cleaning was made (Hijmans *et al.*, 1999). After that we got 150 georeferenced records used for training gain.

### **Environmental Predictors**

To model the species distributions, 20 environmental predictors were selected. We downloaded the altitude (in m) and the 19 bioclimatic predictors (1950 - 2000) of the WORLDCLIM dataset (<http://www.worldclim.org>) for South America at 30 arc-second resolution (~ 1km) (Hijmans *et al.*, 2005a). Data-layer manipulations were performed with DIVA GIS.

Environmental layers used: alt\_sa\_30s\_cut (Altitude), bio1\_sa\_30s\_cut (Annual Mean Temperature), bio2\_sa\_30s\_cut (Mean Diurnal Range [Mean (period max-min)]), bio3\_sa\_30s\_cut (Isothermality [P2/P7]), bio4\_sa\_30s\_cut (Temperature Seasonality [Coefficient of Variation]), bio5\_sa\_30s\_cut (Max Temperature of Warmest Period), bio6\_sa\_30s\_cut (Min Temperature of Coldest Period), bio7\_sa\_30s\_cut (Temperature Annual Range [P5-P6]), bio8\_sa\_30s\_cut (Mean Temperature of Wettest Quarter), bio9\_sa\_30s\_cut (Mean Temperature of Driest Quarter), bio10\_sa\_30s\_cut (Mean Temperature of Warmest Quarter), bio11\_sa\_30s\_cut (Mean Temperature of Coldest Quarter), bio12\_sa\_30s\_cut (Annual Precipitation), bio13\_sa\_30s\_cut (Precipitation of Wettest Period), bio14\_sa\_30s\_cut

(Precipitation of Driest Period), bio15\_sa\_30s\_cut (Precipitation Seasonality [Coefficient of Variation]), bio16\_sa\_30s\_cut (Precipitation of Wettest Quarter), bio17\_sa\_30s\_cut (Precipitation of Driest Quarter), bio18\_sa\_30s\_cut (Precipitation of Warmest Quarter), bio19\_sa\_30s\_cut (Precipitation of Coldest Quarter).

### **Species Distribution Model (SDM) building**

To model the species distributions we selected the modeling application MaxEnt v. 3.3.2 (Phillips *et al.*, 2006). MaxEnt was specifically developed to model species distributions with presence-only data, has shown to outperform most other modelling applications (Elith *et al.*, 2006; Pearson *et al.*, 2007), is least affected by location errors in occurrences (Graham *et al.*, 2008), and best performs when few presence records are available (Wisz *et al.*, 2008). MaxEnt was set to use all species presence records for model building (explained below), by setting the “random test percentage” to zero. The modelling rules were set to use auto features. As measure of SDM accuracy we used the threshold independent and prevalence insensitive area under the curve (AUC) of the receiver operating characteristic (ROC) plot (Fielding & Bell, 1997; McPherson *et al.*, 2004; Raes & ter Steege, 2007), produced by MaxEnt. All measures of SDM accuracy require absences (Fielding & Bell, 1997). When these are lacking, as is the case here, they are replaced by pseudo-absences or sites randomly selected at localities where no species presence was recorded (Ferrier *et al.*, 2002, Phillips *et al.*, 2006). The random points were drawn from cells where in the past collections were made, and hence were corrected for any geographical sampling bias.

MaxEnt was implemented using version 3.3.2 of the software made freely available for download (<http://www.cs.princeton.edu/~schapire/maxent>). Recommended default values for convergence threshold ( $10^{-5}$ ), maximum iterations (500), and 10,000 background points were accepted. The regularization value that is appropriate for the types of environmental variables used and the number of presence points was automatically selected by the program (used to reduce overfitting), as was the selection of environmental variables that is also dependent on the number of presences. Linear and quadratic features were considered. The output of MaxEnt is a

continuous variable ranging from 0 to 100, where high values indicate higher relative suitability.

These models were selected to accommodate the presence-only data and span a range of modeling platforms (Elith *et al.*, 2006).

Using different datasets and scenarios allows us capacity to bracket predicted futures based on an ensemble of models that capture the extremes.

To select the highest performing yet most accurate predictive model, the species was tested in different ways.

We initially used all default features of the program. We also build models using bootstrap replicated type run with 20 replicas. The bootstrapping method was used to obtain an estimate of the AUC for the model. This procedure is recommended when an independent data set is lacking, and provides the least biased estimate of predictive performance of any of the model evaluation methods that are based on resampling (Wintle *et al.*, 2005b).

To all models we selected random seed otherwise the generated models would be the same.

The default method of determining AUC for the MaxEnt model was also applied, after partitioning the data by randomly assigning 30% of occurrences to an evaluation (test) data set, and the remaining 70% to a training data set. Additionally, 10,000 pixels were drawn randomly from the study area as background samples.

We also generate a different test file, using DIVA GIS (Hijmans *et al.*, 2005b), to test the model generated independently. Thus we had then a file with 150 original records and another with 300 generated by DIVA.

Predicted habitat, as well as mean AUC scores, improved with using fewer, better-performing parameters.

Parameters were evaluated in a jackknife test in the MaxEnt program to assess individual parameter contribution, which had to be positive to include the parameter. Final models were run with all the data.

The final selection criteria for MaxEnt were (a) lack of significant strong relationships, (b) AUC value of >0.8, and (c) positive contribution to the model outcome in an internal jackknife test procedure.

## Model evaluation

The predictive performance of the model was evaluated using the area under the receiver operating characteristic curve (ROC). A ROC curve is a plot of true-positive cases (or sensitivity) against corresponding false-positive cases (or  $1 - \text{specificity}$ ) across a range of threshold values (Fielding & Bell, 1997). The area under the curve (AUC) provides a measure of discrimination ability, and varies from 0.5 for a model with discrimination ability no better than random, to 1.0 for a model with perfect discriminatory ability (Pearce & Ferrier, 2000). The specificity is defined using predicted area, rather than true commission (Phillips *et al.*, 2006). This implies that the maximum achievable AUC is less than 1. If test data is drawn from the Maxent distribution itself, then the maximum possible test AUC would be 0.903 rather than 1.

Although this evaluation technique typically requires presence and absence data, it has also been applied to presence-only data by substituting the absences with randomly selected pseudo-absences (Graham & Hijmans, 2006; Hernandez *et al.*, 2006; Lütolf *et al.*, 2006; Phillips *et al.*, 2006). We similarly use it to evaluate and compare the resulting MaxEnt models.

## Distribution Maps

Historical and current distribution maps of *G. cristata* were created using DIVA GIS (Figure 1). IBAs (Important Bird Areas) and conservation units points (Parks, Private Reserves, etc.) were used to compare the species distribution and protected areas across its range. Each area received a different color according to its protection degree: red for no protection, yellow for partial protection and green for total protection.

Our dataset was divided into current and historical records, according to the date on which they were collected. Historical records were considered from the first record of a collected yellow cardinal (1802) until 1989. The recent records were considered from 1990 to the present. This distinction was made for two reasons: 1) to have a clear idea of how diminished the area of occurrence and in which ways and 2) because of intense hunting pressure experienced by species, define the last 20 years as

a current distribution, in an environment where a population could be extinct in that period, it is quite reasonable.

Thus we created maps showing the yellow cardinal historical distribution and the recent one. On top of these data we also used data from the Birdlife IBAs (Important Bird Areas) that was proposed to the region, and the existing map for the species distribution (Birdlife, 2010). The birdlife data was partially used, only for comparison purposes, in full agreement with birdlife guidelines and disclaimers.

We build a map based on these data records to the proposed update of the map currently in use. From the model generated by MaxEnt we built a map that would show environments probably best suited to meet the species needs or to use as future areas for reintroduction programs.

## **RESULTS**

### **Maxent**

From the generated models, those which had the best response in both the AUC as in the test gain were generated with the following settings used during the run: 103 presence records used for training, 43 for testing.

10103 points used to determine the Maxent distribution (background points and presence points).

Environmental layers used: all of them.

Regularization values: linear/quadratic/product: 0.050, categorical: 0.250, threshold: 1.000, hinge: 0.500

Feature types used: product linear quadratic hinge threshold

Random seed: true

Random test points: 30

Auto feature: false

A representation of the resultant best Maxent model is shown in Fig. 1. This output represents cumulative probabilities where the value of a given grid cell (or pixel) is the sum of that cell and all other cells with equal or lower probability,

multiplied by 100 to give a percentage (Phillips *et al.*, 2006). The image uses colors to indicate suitable conditions of the predicted probability. Red indicates high probability of suitable conditioning for the species; green indicates typical conditions of those where the species is found and lighter shades of blue indicating low predicted probability of suitable conditions (Phillips *et al.*, 2008).




Figure 1. Model output.



Figure 2 shows the results of the jackknife test of variable importance. The environmental variable with highest gain when used in isolation is bio1\_sa\_30s\_cut (Annual Mean Temperature), which therefore appears to have the most useful information by itself. The environmental variable that decreases the gain the most when it is omitted is alt\_sa\_30s\_cut (Altitude), which therefore appears to have the most information that isn't present in the other variables.

All the other models made with the complete set of variables had an AUC scoring greater than 0.8, means that no noticeable different was find taking any of the parameters off the model.




Figure 2. Jackknife test of variable importance of regularized training gain.

The Maxent model with independent data test yielded an AUC of 0.889 for the test data.

The models generated with the independent test file were very similar to models generated with the file in which 30% were separated for testing, demonstrating that both the number of points were enough to generate good models as the points themselves have no bias associated to them.

The bootstrapped estimates of the ROC curve for the model resulted in an averaged training AUC for the 20 replicate runs of 0.961, with standard deviation of 0,004 (Fig. 3).

The model that best predicts the relative index of environmental suitability for the species is the model shown in Figure 1. Specificity is defined using the area instead of the commission predicted true. This implies that the maximum AUC achieved is less than 1. In other words, as the test data was obtained from the Maxent distribution itself, and then the maximum possible test AUC was 0.903 instead of 1. So, the resulting AUC of this model on test data was 0.868 (Fig. 4).




Figure 3. Receiver operating characteristic (ROC) curve for the data. The average training AUC for the replicate runs is 0.961.




Figure 4. Receiver operating characteristic (ROC) curve for the data. The test AUC is 0.868.

### Distribution Maps

A map of the historical records of *G. cristata* is shown in fig. 5 and another one with recent records is shown in fig. 6. Both maps were designed plotting together with suggested areas as IBAs by BirdLife and already existing conservation units. A distinction was made between categories of protection for each unit by assigning a different color: red for areas without any protection, yellow for areas with partial protection and green for fully protected areas (most times within state or federal parks).

A polygon based on the current species records was created and plotted on the map with the model generated by Maxent (Fig. 7).




Figure 5. Map showing the historical records of *Gubernatrix cristata* with plotted IBAs and Parks across its range.




Figure 6. Map showing the current records of *Gubernatrix cristata* with plotted IBAs and Parks across its range.




Figure 7. Map showing the species distribution model with a plotted polygon of current records.

## DISCUSSION

### Habitat suitability

This study sought to explore environmental attributes that may define habitat suitability of the yellow cardinal at a broad spatial scale.

The model agrees on the relationship between yellow cardinal occupancy and each environmental variable. Even though Mean Diurnal Range (`bio2_sa_30s_cut`) was included in the MAXENT model, Fig. 2 shows that this variable contributed almost no information to the model. The positive associations with elevation, temperature seasonality, and precipitation of driest quarter, suggest that the yellow cardinal prefers areas not high in altitude (below 1000m), with a temperature seasonality between 4°C and 30 °C and precipitation below 300 mm a quarter. Those conditions are according to what is found in the ecoregion of Espinal (Marchiori, 2004; Parodi, 2002).

A very low association with increasing temperature diurnal range also suggests that this species prefers habitats with low daily temperature variations. This apparent

selection for low variations in temperature is again likely to be related to the vegetation type associated with this ecoregion.

### **Modelling approach**

One problem often encountered when dealing with a rare species, is that the number of presence sites are likely to be relatively small and this then limits the number of candidate variables that should be used in model development. A high ratio of candidate models to the number of species observations can lead to overfitting of the model, i.e. inclusion of spurious variables (Burnham & Anderson, 2002; Harrell, 2001), thereby emphasizing the importance of using expert scientific knowledge in the a priori selection of variables. A plausible variable set would be in the bounds of the rule of thumb of Harrell (2001) where no more than  $n/10$  variables should be included in the final model, where  $n$  is the total sample size or. In this case, this number would be 15. As none of the variables decrease strongly the fit of the model (fig. 2) nor any publications says which ones are the best, we decided to model with the complete set of variables (20).

While MAXENT uses background information, pixels without species records are not interpreted as absences (Phillips et al., 2006), and therefore this method may not suffer from biases resulting from the inclusion of false absences in the model. A study by Elith et al. (2006) comparing various methods for modelling presence-only data showed that MAXENT outperformed most other methods when evaluated using independent presence and absence data. MAXENT has also been shown to be superior to other approaches that use presence data only (i.e. not requiring pseudo-absences) such as Domain, GARP, and BIOCLIM (Elith et al., 2006; Hernandez et al., 2006; Pearson et al., 2007; Phillips et al., 2006). Even though post hoc evaluation of predictive performance should involve a comparison with independent data (Fielding & Bell, 1997; Manel et al., 1999; Pearce & Ferrier, 2000; Scott et al., 2002), with studies involving rare species, such data are difficult and costly to collect. Resampling techniques, such as bootstrap used in the current study, are popular methods for dealing with the lack of an independent data set, as is the MAXENT approach of partitioning the data set into training and test data (Guisan & Zimmermann, 2000). A

further complication when evaluating model performance with presence-only data is the difficulty of evaluating false-positive prediction errors (Hernandez et al., 2006). The approach used in this study where ROC plots were generated using presence and randomly selected pseudo-absence data (or background samples) is one solution to this problem. But here the AUC is interpreted as a measure of the ability of the model to discriminate between suitable habitat and a random background sample, rather than suitable and unsuitable habitat (as would be the case with real absence data) (Hernandez et al., 2006; Phillips et al., 2006). The similarity of the AUC derived from the ROC plots for MAXENT model suggests that it performed well and values greater than 0.8 also indicates that its ability to discriminate suitable habitat from random was high. The good performance of this presence-only method suggests that the environmental variables used here describe the habitat where the species occurs within its range quite well. However, assuming the species is in disequilibrium, there are clearly additional factors not included in the variable set that account for the inability of the bird to occupy all suitable habitats. This highlights one of the problems of distributional models, as most rely on environmental attributes, and rarely are factors such as dispersal limitations or interspecific interactions included in such models.

### **Distribution Maps**

When analyzing the species historical records map, we noticed that yellow cardinal is distributed over a wide area, covering different phytogeographical zones (Di Giacomo, 2005). Many of these records were taken in areas without any protection (Fig. 5). When this map is compared with that of recent records is clear the shrinkage that populations have suffered in recent years. This shrinkage makes an irregular arc shape along the species distribution, resembling much to the arc-shape made by the ecoregion Espinal (Marchiori, 2004; Parodi, 2002). When the polygon drawn from the recent record is plotted on the map generated from the model of potential distribution (Fig. 7) this arc is also formed. What can be analyzed from this is an apparent association of the yellow cardinal to a particular vegetation type. Throughout its geographic range and especially in the area where the arc forms, the characteristic

vegetation type is savanna steppe, with abundant plants of the genus *Prosopis* (Marchiori, 2004; Di Giacomo, 2005). It is not surprising that this type of environment is the last refuge of the yellow cardinal, since it is a type of vegetation that also occurs in areas of difficult access.

## **CONCLUSION**

While models derived from occurrence information have been shown to overestimate the extent of suitable habitat compared to models based on presence and absence data (Zaniewski et al., 2002; Brotons et al., 2004; Engler et al., 2004; Pearce & Boyce, 2006), predictive modelling is usually an iterative process, and this model can be used as a starting point to approximate habitat suitability, and hence guide future surveys for further data collection and model refinement. If the objective of future surveys of the yellow cardinal in its distribution range is to detect other populations, then areas predicted to have high relative suitability in the spatial model (Fig. 1) is a good starting point for further targeted survey.

The hunting pressure on the species is so great that the yellow cardinal nowadays seems to be only found in areas of difficult access. Many of the areas suitable for the species shown by the model may still contain populations and can also be used as key areas for future reintroductions into the yellow cardinal conservation program.

## **ACKNOWLEDGEMENTS**

We thank to Dr. J. M. Alexandrino for the very first lesson in how to use MaxEnt. We thank to Dr. M. B. Martins for the discussions in the model analysis and to C. Zank for the good tips and help with DIVA GIS. We thank CNPq for providing the doctorate scholarship to C. Martins-Ferreira, to CAPES for the Sandwich Scholarship, and to FBPCN for financing the project.



## REFERENCES

- Azpiroz, A. B. (2003) *Aves Del Uruguay: lista e introducción a su biología y conservación*. Aves Uruguay-GUPECA, Montevideo.
- Belton, W. (1985) Birds of Rio Grande do Sul, Brazil. Part 2: Formicariidae through Corvidae. *Bulletin of the American Museum of Natural History*, **180** (1): 1-242.
- BirdLife International (2000) *Threatened birds of the world*. Lynx Edicions and BirdLife International, Barcelona and Cambridge/UK.
- BirdLife International (2010) Species factsheet: *Gubernatrix cristata*. Downloaded from <http://www.birdlife.org> on 29/7/2010.
- Burnham, K.P. & Anderson, D.R. (2002) *Model selection and multimodel inference*. A practical information-theoretic approach, 2nd edn. Springer, New York.
- Collar, N. J.; Gonzaga, L. P.; Krabbe, N.; Madroño Nieto, A.; Naranjo, L. G.; Parker III, T. A. & Wege, D. C. (1992) *Threatened birds of the Americas: The I.C.B.P. / I.U.C.N. Red Data Book*. Third edition, part 2. Smithsonian Inst. Press, ICBP, Cambridge.
- Chebez, J. C. (1994) *Los que se van*. Albatros, Buenos Aires.
- Chebez, J. C. (1996) Campos misioneros: algo más que el confín de la selva. *Nuestras Aves* 34: 4-16.
- Chebez, J. C.; Rey, N. R.; Babarskas, M. & Di Giacomo, A.G. (1998) *Las Aves de los Parques Nacionales de la Argentina*. Monografía Especial L.O.L.A. Nº 12. Administración de Parques Nacionales y Asociación Ornitológica del Plata, Buenos Aires.
- Di Giacomo, A. S. (editor) (2005) Áreas importantes para la conservación de las aves en Argentina: *sítios prioritários para la conservación de la biodiversidad*. Temas de Naturaleza y Conservación 5:1-514. Aves Argentinas/Asociación Ornitológica del Plata, Buenos Aires.
- Elith, J.; Graham, C.H.; Anderson, R.P.D.M.; Ferrier, S.; Guisan, A.; Hijmans, R. J.; Huettmann, F.; Leathwich, J.R.; Lehmann, A.; Li, J.; Lohmann, L.G.; Loiselle, B.A.; Manion, G.; Moritz, C.; Nakamura, M.; Nakazawa, Y.; Overton, J.M.; Peterson, A.T.; Phillips, S.J.; Richardson, K.; Scachetti-Pereira, R.; Schapire, R.E.; Soberon, J.; Williams, S.; Wisz, M.S. & Zimmermann, N.E. (2006) Novel methods improve prediction of species' distributions from occurrence data. *Ecography*, **29**, 129-152.
- Ferrier, S.; Watson, G; Pearce, J. & Drielsma, M. (2002) Extended statistical approaches to modeling spatial pattern in biodiversity in northeast New South Wales. I. Species-level modelling. *Biodiversity and Conservation*, **11**, 2275-2307.
- Fielding, A. H. & Bell, J. F. (1997) A review of methods for the assessment of prediction errors in conservation presence/absence models. *Environmental Conservation*, **24**, 38\_49.
- Fontana, C. S.; Bencke, G. A. & Reis, R. E. (2003) *Livro vermelho da fauna ameaçada de extinção no Rio Grande do Sul*. EDIPUCRS, Porto Alegre.

- Fraga, R. M. (1997) Sección Aves. *In*: Garcia-Fernandez, J. J.; Ojeda, R. A.; Fraga, R. M.; Díaz, G. B. & Baigún, R. J. (Ed.). Libro rojo de mamíferos y aves amenazadas de la Argentina. FUCEMA, Buenos Aires.
- Graham, C.H. & Hijmans, R.J. (2006) A comparison of methods for mapping species ranges and species richness. *Global Ecology and Biogeography*, **15**, 578–587.
- Graham, M. H.; Elith, J.; Hijmans, R.J.; Guisan, A.; Peterson, A.T.; Loiselle, B.A. & Nceas Predicting Species Distributions Working Group (2008) The influence of spatial errors in species occurrence data used in distribution models. *Journal of Applied Ecology*, **45**, 239–247.
- Guisan, A. & Zimmermann, N.E. (2000) Predictive habitat distribution models in ecology. *Ecological Modelling*, **135**, 147–186.
- Harrell, F.E. (2001) *Regression modelling strategies with applications to linear models, logistic regression and survival analysis*. Springer Series in Statistics. Springer-Verlag, New York.
- Hernandez, P.A., Graham, C.H., Master, L.L. & Albert, D.L. (2006) The effect of sample size and species characteristics on performance of different species distribution modeling methods. *Ecography*, **29**, 773–785.
- Hijmans, R.J.; Schreuder, M.; De la Cruz, J. & Guarino, L. (1999) Using GIS to check coordinates of genebank accessions. *Genetic Resources and Crop Evolution*, **46**, 291–96.
- Hijmans, R.J.; Cameron, S.E.; Parra, J.L.; Jones, P.G. & Jarvis, A. (2005a) Very high resolution interpolated climate surfaces for global land areas. *International Journal of Climatology*, **25**, 1965–1978.
- Hijmans, R. J.; Guarino, L.; Jarvis, A.; O'brien, R. & Mathur, P.; Bussink, C.; Cruz, M.; Barrantes, I. & Rojas, E. (2005b) *DIVA-GIS v. 5.2. Manual*. Available at: <<http://www.diva-gis.org/>>. Accessed in: 10 June 2006.
- Kopta, Federico (1999) Disminución de la fauna autóctona. *In*: Problemática ambiental con especial referencia a la Provincia de Córdoba – Capítulo 5. Fundación Ambiente, Cultura y Desarrollo – ACUDE. Edición auspiciada por UNESCO y financiada por la Embajada Real de los Países Bajos. 203 págs. Córdoba, Argentina.
- Lapitz, Fernando (2010) Cardenal Amarillo – *Gubernatrix cristata*. Achará Digital. Año 1 (1) pág. 8. AVES URUGUAY, Montevideo.
- Lütolf, M., Kienast, F. & Guisan, A. (2006) The ghost of past species occurrences: improving species distribution models for presence-only data. *Journal of Applied Ecology*, **43**, 802–815.
- Manel, S., Dias, J.M., Buckton, S.T. & Ormerod, S.J. (1999) Alternative methods for predicting species distribution: an illustration with Himalayan river birds. *Journal of Applied Ecology*, **36**, 734–747.
- Martins-Ferreira, C. (2007) Biogeografía e ecologia do Cardeal-Amarelo *Gubernatrix cristata* (Emberizidae) – resultados preliminares / Biogeography and ecology of Yellow Cardinal *Gubernatrix cristata* (Emberizidae) – preliminary results. *In*:

- Congresso Brasileiro de Ornitologia (15: 2007: Porto Alegre, RS) [programação e resumos]. EDIPUCRS, Porto Alegre.
- McPherson, J. M.; Jetz, W. & Rogers, D. J. (2004) The effects of species' range sizes on the accuracy of distribution models: ecological phenomenon or statistical artefact? *Journal of Applied Ecology*, **41**, 811-823.
- ORNIS. *Data portal*. Accessed at (<http://ornisnet.org>) on 10/11/2009.
- Parodi, L. R. (2002) Las regiones fitogeográficas Argentinas. *Ciência & Ambiente*, **24**, 25-34.
- Pearce, J. & Ferrier, S. (2000) Evaluating the predictive performance of habitat models developed using logistic regression. *Ecological Modelling*, **133**, 225-245.
- Pearson, R. G.; Raxworthy, C.J.; Nakamura, M. & Peterson, A.T. (2007) Predicting species distributions from small numbers of occurrence records: a test case using cryptic geckos in Madagascar. *Journal of Biogeography*, **34**, 102-117.
- Phillips, S. J.; Anderson, R. P. and Schapire, R. E. (2006) Maximum entropy modeling of species geographic distributions. *Ecological Modelling*, **190**, 231-259.
- Phillips, S.J. Maxent V. 3.3.2. *A brief tutorial on Maxent*. Downloaded from <http://www.cs.princeton.edu/~schapire/maxent/> on 10/11/2009.
- Raes, N. & ter Steege, H. (2007) A null-model for significance testing of presence-only species distribution models. *Ecography*, **30**, 727-736.
- Scott, J.M., Heglund, P.J., Morrison, M.L., Haufler, J.B., Raphael, M.G., Wall, W.A. & Samson, F.B. (2002) Predicting species occurrences: *issues of accuracy and scale*. Island Press, Washington.
- Wintle, B.A., Kavanagh, R.P., McCarthy, M.A. & Burgman, M.A. (2005a) Estimating and dealing with detectability in occupancy surveys for forest owls and arboreal marsupials. *Journal of Wildlife Management*, **69**, 905-918.
- Wisz, M. S. ; Hijmans, R. J.; Li, J.; Peterson, A. T.; Graham, C. H.; Guisan, A. & NCEAS Predicting Species Distributions Working Group (2008) Effects of sample size on the performance of species distribution models. *Diversity and Distributions*, **14**, 763-773.

## BIOSKETCH

Claiton Martins-Ferreira is a Ph.D student at Graduate Program in Animal Biology, based in the Zoology Department at the Federal University of Rio Grande do Sul (UFRGS). See our website for more information (in Portuguese) (<http://www.ufrgs.br/zoologia/>).

Author contributions: This study was conceived by C.M.F., who programmed and implemented the modeling experiments, analyzed the results and prepared the manuscript. The study was supervised by T.R.O.F. All authors discussed the results and commented on the manuscript.

## **CAPÍTULO CINCO**

### **CONCLUSÕES E RECOMENDAÇÕES PARA A CONSERVAÇÃO DA ESPÉCIE**



### 5.1. Viabilidade do cardeal-amarelo no RS, Uruguai e Argentina

A partir das expedições feitas durante a execução do projeto, pudemos constatar que as melhores áreas para serem transformadas em UCs, no RS, se localizam na serra do sudeste, mais especificamente no entorno da cidade de Herval, na região da várzea do rio Jaguarão, localidade de Passo São Diogo. Naquela área encontramos uma fitofisionomia muito semelhante àquela encontrada em áreas que ainda possuem populações de cardeal-amarelo, tanto na Argentina quanto no RS (Parque Estadual do Espinilho). A região de Herval já foi incluída dentro da IBA RS08 – Região de Pinheiro Machado e, de acordo com nossos dados, essa região possuiria ainda habitats adequados para espécie. Embora não tenhamos avistado o cardeal-amarelo nas cidades vizinhas de Pedras Altas e Pinheiro Machado, é importante estimular a criação de uma rede de RPPNs naquela região. Tanto para a proteção de alguns exemplares errantes ou para um programa de reintrodução, conforme mostra o modelo de distribuição potencial para a espécie. Como a região sofreu muito, e ainda sofre, com o impacto da coleta/caça da avifauna, há que se estabelecer um programa de educação ambiental junto à comunidade, paralelo à criação de UCs e de programa de reintrodução/translocação da espécie.

Outra área que cabe destacar é o entorno do Parque Estadual do Espinilho. Embora o parque já seja uma Unidade de Conservação, carece de melhor fiscalização. Os relatos de avistamentos anteriores ao nosso reportam a observação de *G. cristata* também em áreas fora do parque. Há vários habitats interessantes para a espécie tanto dentro quanto fora do parque, o que torna aquela região como um todo extremamente importante para a sobrevivência da espécie.

Encontramos também áreas adequadas para a proteção integral da espécie no Uruguai, no interior do município de Minas, Departamento de Lavalleja, onde as áreas de proteção seriam dentro de propriedades particulares, estimulando a fiscalização por meio dos próprios fazendeiros, que se mostraram dispostos a colaborar com a conservação da espécie.

As áreas mais adequadas para se encontrar cardeal-amarelo na Argentina seriam em Corrientes, Entre Ríos, sul da província de Buenos Aires, Río Negro, La Pampa, San Luís, Córdoba e Tucumán. Muitas dessas áreas tem registros apenas

históricos de cardeal-amarelo, como é o caso de Tucumán, embora aparentemente ainda tenham áreas adequadas para a espécie. Áreas adequadas mas que não possuam mais populações da espécie seriam áreas potenciais para integrarem um futuro programa de reintrodução da espécie.

O status de conservação de *Gubernatrix cristata* é avaliado mundialmente pela IUCN como Em Perigo; no Brasil é considerado Criticamente em Perigo e no RS, Em Perigo. Os resultados obtidos no projeto nos levam a concluir que deveria ser feita uma reavaliação desse status, tanto para o RS, passando a Criticamente em Perigo, quanto em nível mundial, passando também para a categoria de Criticamente em Perigo.

A caça intensa continua sendo a principal pressão contra as populações na natureza e foi o principal fator na provável extinção da espécie na região da serra do sudeste. Eliminando essa pressão e estimulando a criação de UCs é possível que *G. cristata* volte a recolonizar a região.

A principal implicação para a conservação da espécie a partir da falta de estruturação populacional é que um programa conservacionista de reprodução em cativeiro tem grandes chances de obter sucesso sem a preocupação extra de manter uma diversidade haplotípica associada a áreas geográficas.

Os ambientes encontrados no RS (Herval e Pedras Altas) apresentam espécies do gênero *Scutia* sp. e *Prosopis* sp.. No Parque Estadual do Espinilho (PEE) predominam *Prosopis* sp., *Celtis* sp., *Acacia* sp. e *Aspidosperma quebracho-blanco*. No PEE a formação tipo parque domina a paisagem, enquanto que em Herval e Pedras Altas o mesmo não ocorre. Isso se deve, provavelmente, ao pastejo de gado intenso nas áreas de fazendas e não protegidas, encontradas no sul do RS.

Em Corrientes, Argentina, a fitofisionomia é bastante semelhante aquela encontrada no PEE, com espécies similares. No Departamento de Lavalleja, Uruguai, onde foram capturados exemplares de cardeal-amarelo, a fitofisionomia encontrada é bastante similar à do sul do RS. Não parece haver uma correlação direta entre a presença/ausência de *Gubernatrix cristata* e a presença/ausência de gado. Essa correlação está mais relacionada, possivelmente, ao uso do hábitat pela ave, tanto para construção de ninhos quanto para alimentação. Ou seja, onde há determinadas espécies botânicas é possível encontrar cardeal-amarelo.

O que se observou em campo, além do descrito acima, é que a pressão de caça

causou a rarefação da espécie, mesmo nos habitats considerados ideais.

Os resultados do trabalho também nos levam a concluir que a espécie está fortemente associada ao ambiente conhecido como savana tipo parque. O mapa construído a partir dos registros históricos da espécie e do modelo de distribuição potencial mostra claramente essa associação quando avaliamos os registros mais recentes da espécie. Ou seja, o cardeal-amarelo ainda sobrevive nas áreas mais remotas e com vegetação tipo espinilho. Ao compararmos os mapas da distribuição histórica com o mapa da distribuição recente de *G. cristata* pode-se concluir claramente que houve uma redução significativa da área de distribuição da espécie.

## **5.2. Recomendações para a conservação: um primeiro passo para a construção de um plano internacional para a conservação do cardeal-amarelo no Sul da América do Sul.**

A partir dos resultados encontrados durante a execução do projeto, recomendamos as seguintes ações para a conservação da espécie:

- 1) Parceria multi-institucional (órgãos governamentais, universidades, ONGs) para o estabelecimento de um protocolo sanitário de manejo da população em cativeiro da espécie a nível nacional;
- 2) Parceria multi-institucional para o estabelecimento oficial de um programa de reprodução em cativeiro com a criação e manutenção de um *studbook* da espécie;
- 3) Acompanhamento e monitoramento dos indivíduos encontrados em vida livre com o intuito de conhecer melhor a respeito da biologia reprodutiva, uso do habitat e comportamento da espécie a fim de dar bases para a elaboração de uma análise de viabilidade populacional;
- 4) Prosseguimento com as análises genéticas incluindo amostras das populações de La Pampa e de outras províncias argentinas, possivelmente através de um programa de parceria internacional, que possibilitarão o conhecimento mais aprofundado a respeito das populações em vida livre ainda existentes;
- 5) Designar unidades de conservação de proteção integral ao longo da distribuição da espécie dentro da Argentina, Uruguai e RS, naqueles locais que se mostraram mais adequados para a sobrevivência de cardeal-amarelo e que ainda não possuem unidade

de conservação alguma ou se possuem, não é de proteção integral;

6) Proteger bosques de *Prosopis* sp. no sul e oeste do RS, oeste e norte do Uruguai e centro e sul da Argentina;

7) Coibir o tráfico de animais silvestres, que é efetivamente a maior ameaça para a sobrevivência da espécie na natureza, utilizando tanto medidas legais quanto socioeducativas através de campanhas de educação ambiental das populações do entorno de áreas onde ainda existem cardeais-amarelos na natureza, especialmente os municípios de Barra do Quaraí, no RS e Bella Unión, no Uruguai;

8) Envolver os criadores particulares de cardeal-amarelo no programa de conservação da espécie. Muitas dessas pessoas têm boas intenções. São apenas mal orientadas. O contingente de criadores de cardeal-amarelo dentro do SISPASS/IBAMA não é algo a se descartar.



### 5.1. Viability of yellow cardinal in RS, Uruguay and Argentina

During the expeditions made while the project execution, we found that the best areas to be turned into conservation areas in the RS, are located in the southeast hills, specifically around the city of Herval in the region of the river banks Jaguarão, Passo de São Diogo town. The vegetation type found in that area was quite similar to that one found in areas that still have populations of yellow cardinal, both in Argentina and in the RS (Espinilho State Park). The Herval region have already been included within the IBA RS08 - Região de Pinheiro Machado, and according to our data, this region still possess suitable habitats to the species. Although we have not seen the yellow cardinal in the neighboring towns of Pedras Altas and Pinheiro Machado, it is important to stimulate the creation of a network of RPPNs (Private Reserves) there. Both for the protection of some wandering specimens or for a reintroduction program, as shown in the species distribution model. As the region has suffered much, and still suffers, the impact of sample/hunting birds, we must establish an environmental education program in the community, alongside the creation of UCs and reintroduction/translocation species program.

Another noteworthy area is the surrounding area of Espinilho State Park. Although the park is already a conservation area, needs better control. The sightings accounts prior to ours also reported the observation of *G. cristata* in areas outside the park. There are several interesting habitats for the species both within and outside the park, which makes the region as a whole extremely important for the species survival.

We also found suitable areas for species full protection in Uruguay, within the municipality of Minas, Department of Lavalleja, where protected areas would be within private properties, stimulating the supervision by the very farmers who were willing to collaborate with the species conservation.

The most suitable areas to find yellow cardinal in Argentina would be in Corrientes, Entre Ríos, the southern of Buenos Aires province, Río Negro, La Pampa, San Luís, Córdoba and Tucumán. Many of these areas have only historical records of yellow cardinal, as is the case of Tucumán, although apparently still have suitable areas for the species. Suitable areas but having no more species populations would be potential areas to join a future species reintroduction program.

The conservation status of *Gubernatrix cristata* is valued worldwide by IUCN as Endangered, in Brazil is considered Critically Endangered and in RS, Endangered. The results obtained in the project lead us to conclude that it should be make a reassessment of this status for both RS as worldwide to become Critically Endangered.

Hunting remains the main intense pressure against the wild populations and was a major factor in the likely of species extinction in the southeastern hills region. Eliminating the pressure and stimulating the creation of PAs is possible that *G. cristata* re-colonize the region.

The lack of population structure has one important implication for the species conservation: the conservation breeding program has great chances of success without the extra worry of maintaining an haplotype diversity associated with geographic areas.

The environments found in the RS (Herval and Pedras Altas) have species of the genus *Scutia* sp. and *Prosopis* sp. In Espinilho State Park (PEE) predominates *Prosopis* sp., *Celtis* sp., *Acacia* sp. and *Aspidosperma quebracho-blanco*. In the PEE savanna-steppe formation dominates the landscape, while in Herval and Pedras Altas the same is not true. This is probably due to the intensive grazing livestock in not protected farming areas, found in southern Brazil.

In Corrientes, Argentina, the vegetation type is very similar to that found in PEE, with similar species. Department of Lavalleja, Uruguay, where we captured yellow cardinal specimens, the vegetation type found is very similar to southern Brazil. There seems not to be a direct correlation between the presence/absence of *Gubernatrix cristata* and the presence/absence of cattle. This correlation is possibly more related to the bird habitat use, both for nest building and for food. In other words, where there are certain plant species it's likely to find yellow cardinal.

What was observed in the field, beyond what is described above, is that hunting pressure caused the species thinning, even in considered optimal habitats.

The results of the study also lead us to conclude that the species is strongly linked to the environment known as the savannah steppe. The map built from the species historical records and the potential distribution model clearly shows that association when evaluating the most recent species records. The yellow cardinal still survive in remote areas and vegetation type espinilho. When comparing the *G. cristata* historical distribution map with the recent distribution map of can be clearly concluded

that there was a significant shrinkage in the species range.

## **5.2. Conservation Recommendations: A first step in building a regional action plan for the conservation of yellow cardinal in Southern South America.**

From the results found during the project running, we recommend the following actions for the species conservation:

- 1) A multi-institutional partnership (government agencies, universities, NGOs) to define a health protocol for species captive population management at national level;
- 2) Multi-institutional Partnership for the establishment of a official captive breeding program with the creation and maintenance of a yellow cardinal studbook;
- 3) monitoring and tracking of individuals found in the wild in order to learn more about the reproductive biology, habitat use and species behavior to provide bases for developing a population viability analysis;
- 4) Continuing with the genetic analysis including samples of La Pampa populations and other Argentine provinces, possibly through a program of international partnership that will allow a deeper knowledge about the populations that still exist in the wild;
- 5) designate conservation units fully protected along the species distribution within Argentina, Uruguay and RS, at those sites that were more suitable for yellow cardinal survival and still do not have any conservation area or if they have, is not fully protected;
- 6) Protect *Prosopis* sp. woodlands in the south and west of the RS, west and north of Uruguay and central and southern Argentina;
- 7) Repress the wild animal trade, which is actually the biggest threat to the species survival in nature, using both socio-legal measures as environmental education campaigns. We suggest mostly, work with the surrounding populations from areas where still there are yellow cardinals in the wild, e.g. the municipalities of Barra do Quaraí in RS and Bella Union, Uruguay;
- 8) Involve the private yellow cardinal breeders in the species conservation program. Many of these people have good intentions. They're just misguided. The number of yellow cardinal owners within the SISPASS/IBAMA is not something to dismiss.

## 6. REFERENCIAS

- Avise, J.C. (1994) *Molecular markers, natural history and evolution*. Chapman & Hall, New York.
- Avise, J.C. (2000) *Phylogeography: the history and formation of species*. Harvard University Press, Cambridge.
- Avise, J.C.; Arnold, J.; Ball JR., R.M.; Bermingham, E.; Lamb, T.; Neigel, J.E.; Ree, C.A. & Saunders, N.C. (1987a) Intraespecific phylogeography: the mitochondrial DNA bridge population genetics and systematics. *Annu. Rev. Ecol. Syst.*, **18**, 489-522.
- Azpiroz, A. B. (2003) *Aves Del Uruguay: lista e introducción a su biología y conservación*. Aves Uruguay-GUPECA, Montevideo.
- Belton, W. (1985) Birds of Rio Grande do Sul, Brazil. Part 2: Formicariidae through Corvidae. *Bulletin of the American Museum of Natural History*, **180**, (1): 1-242.
- Belton, W. (1994) *Aves do Rio Grande do Sul: distribuição e biologia*. Editora UNISINOS, São Leopoldo.
- Bencke, G. A. (2001). *Lista de referência das aves do Rio Grande do Sul*. Fundação Zoobotânica do Rio Grande do Sul, Porto Alegre.
- Bermingham, P. & Moritz, C. (1998) Comparative phylogeography: concepts and applications. *Molecular Ecology*, **7**, (4):367-370.
- Bertonatti, C. & Guerra, A. L. (1997) Hibridization between the yellow cardinal (*Gubernatrix cristata*) and the common diuca finch (*Diuca diuca*) in the wild in Argentina. *Hornero*, **14**,(4):235-42.
- Bertonatti, C. & Guerra, A. L. (2001) Nuevos registros de híbridos entre cardenal amarillo (*Gubernatrix cristata*) y diuca común (*Diuca diuca*). *Nuestras Aves*, **41**,27.
- BIRDLIFE INTERNATIONAL. (2000) *Threatened birds of the world*. Lynx Edicions and BirdLife International, Barcelona and Cambridge/UK.
- Broders, H.G.; Mahoney, S.P.; Montevecchi, W.A. & Davidson, W.S. (1999) Population genetic structure and the effect of founder events on the genetic variability of moose, *Alces alces*, in Canada. *Molecular Ecology*, **8**, 1309-1315.
- Brown, W.M.; George, M.J.R. & Wilson, A.C. (1979) Rapid Evolution of animal mitochondrial DNA. *Proc. Natl. Acad. Sci. USA*, **76**, 1967-1971.
- Brown, W.M. (1985) The mitochondrial genome of animals. *In*: Mac Intyre, R.J. (ed). *Molecular Evolutionary Genetics*. Plenum, New York.
- Brown, G.G.; Gadalata, G.; Pepe, G.; Saccone, C. & Sbisà, E. (1986) Structural conservation and variation in the D-loop containing region of vertebrate mitochondrial DNA. *Journal of Molecular Biology*, **192**, 503-511.
- Burg, T. M. & Croxall, J. P. (2001) Global relationships amongst black-browed and grey-headed albatrosses: analysis of population structure using mitochondrial DNA and microsatellites. *Molecular Ecology*, **10**, 2647-2660

- Cabrera, A.L. & Willink, A. (1973) *Biogeografía de América Latina*. Secretaría General de la Organización de los Estados Americanos, Washington.
- Chebez, J. C. (1994) *Los que se van*. Albatros, Buenos Aires.
- Chebez, J. C. (1996) Campos misioneros: algo más que el confín de la selva. *Nuestras Aves* 34: 4-16.
- Chebez, J. C.; Rey, N. R.; Babarskas, M. & Di Giacomo, A.G. (1998) *Las Aves de los Parques Nacionales de la Argentina*. Monografía Especial L.O.L.A. Nº 12. Administración de Parques Nacionales y Asociación Ornitológica del Plata, Buenos Aires.
- Comitê Brasileiro de Registros Ornitológicos – CBRO (2009) *Listas das aves do Brasil. Versão 9/8/2009*. Available at <<http://www.cbro.org.br>>. Accessed in: [25/08/2009].
- Craig, J.L. (1994) Meta-populations: is management as flexible as nature? p. 50-63 In: Olney, P.J.S., Mace, G.M. & Feistner, A.T.C. (eds.) *Creative Conservation: interactive management of wild and captive animals*. Chapman & Hall, London.
- Cristi, Gustavo A. (1958) Filaridos em *Gubernatrix cristata* (Cardenal amarillo) – su comprobacion en el Uruguay. **Anales de la Facultad de Veterinaria**. Pag. 11 – 17. Republica Oriental Del Uruguay.
- Damiani, R.V.; Ribeiro, A.C. & Martins-Ferreira, C. (2009) Projeto Cardeal-Amarelo, resultados preliminares do monitoramento de *Gubernatrix cristata* no Parque Estadual do Espinilho – RS. In: CONGRESSO BRASILEIRO DE ORNITOLOGIA, XVII, 2009, Aracruz. *Biogeografia das aves da Mata Atlântica*. TEC ART, Aracruz/ES.
- Da Silva, M.N & Patton J.L. (1998) Molecular phylogeography and the evolution and conservation of Amazonian mammals. *Molecular Ecology*, **7**, 475-486.
- Di Giacomo, A. S. (editor) (2005) Áreas importantes para la conservación de las aves en Argentina: *sítios prioritários para la conservación de la biodiversidad*. Temas de Naturaleza y Conservación 5:1-514. Aves Argentinas/Asociación Ornitológica del Plata, Buenos Aires.
- Elith, J.; Graham, C.H.; Anderson, R.P.D.M.; Ferrier, S.; Guisan, A.; Hijmans, R. J.; Huettmann, F.; Leathwich, J.R.; Lehmann, A.; Li, J.; Lohmann, L.G.; Loiselle, B.A.; Manion, G.; Moritz, C.; Nakamura, M.; Nakazawa, Y.; Overton, J.M.; Peterson, A.T.; Phillips, S.J.; Richardson, K.; Scachetti-Pereira, R.; Schapire, R.E.; Soberon, J.; Williams, S.; Wisz, M.S. & Zimmermann, N.E. (2006) Novel methods improve prediction of species' distributions from occurrence data. *Ecography*, **29**, 129-152.
- Firestone, K.B.; Elphinstone, M.S.; Sherwin, W.B. & Houlden, B.A. (1999) Phylogenetical population structure of tiger quolls *Dasyurus maculates* (Dasyuridae: Marsupialia), an endangered carnivorous marsupial. *Molecular Ecology*, **8**, 1613-1625.
- Fontana, C. S.; Bencke, G. A. & Reis, R. E. (2003) *Livro vermelho da fauna ameaçada de extinção no Rio Grande do Sul*. EDIPUCRS, Porto Alegre.
- Fraga, R. M. (1997) Sección Aves. In: Garcia-Fernandez, J. J.; Ojeda, R. A.; Fraga, R. M.; Díaz, G. B. & Baigún, R. J. (Ed.). *Libro rojo de mamíferos y aves amenazadas de la Argentina*. FUCEMA, Buenos Aires.

- Frankham, R.; Ballou, J. D. & Briscoe, D. A. (2002) *Introduction to conservation genetics*. Cambridge University Press, Cambridge.
- Gaines, M.S.; Diffendorfer, J.E.; Tamarin, R.H. & Whittam, T.S. (1997) The effects of habitat fragmentation on the genetic structure of small mammal populations. *Journal of Heredity*, **88**, 294-304.
- Goodman, D. (1987) The demography of chance extinction. p.11-34 *In*: Soulé, M.E. (ed.) *Viable Populations for Conservation*. Cambridge University Press, Cambridge.
- Hartl, D.L. & Clark, A.G. (1997) *Principles of population genetics* (3a. ed.) Sinauer Associates, Sunderland.
- Hedrick, P.W., Lacy, R.C., Allendorf, F.W. & Soulé, M.E. (1996) Directions in conservation biology: comments on Caughley. *Conservation Biology*, **10**, 1312-1320.
- Hijmans, R.J.; Cameron, S.E.; Parra, J.L.; Jones, P.G. & Jarvis, A. (2005a) Very high resolution interpolated climate surfaces for global land areas. *International Journal of Climatology*, **25**,1965-1978.
- Hijmans, R. J.; Guarino, L.; Jarvis, A.; O'brien, R. & Mathur, P.; Bussink, C.; Cruz, M.; Barrantes, I. & Rojas, E. (2005b) *DIVA-GIS v. 5.2. Manual*. Available at: <<http://www.diva-gis.org/>>. Accessed in: 10 June 2006.
- Hoelzel, A.R., Halley, J. & Campagna, C. (1993) Elephant seal genetic variation and the use of simulation models to investigate historical population bottlenecks. *Journal of Heredity*, **84**,443-49.
- Hudson, W.H. (1920) *Birds of La Plata*. E. P. Dutton & Co., New York.
- Ibrahim, K.M.; Nichols, R.A. & Hewitt (1996) Spatial patterns of genetic variation generated by different forms of dispersal during range expansion. *Heredity*, **77**, 282-291.
- Kopta, Federico (1999) Disminución de la fauna autóctona. *In*: Problemática ambiental con especial referencia a la Provincia de Córdoba – Capítulo 5. Fundación Ambiente, Cultura y Desarrollo – ACUDE. Edición auspiciada por UNESCO y financiada por la Embajada Real de los Países Bajos. 203 págs. Córdoba, Argentina.
- Lacy, R.C. (1997) Importance of genetic variation to the viability of mammalian population. *Journal of Mamalogy*, **78**, 320-335.
- Lande, R. (1995) Breeding plans for small populations based on the dynamics of quantitative genetic variance. p.318-340 *In*: Ballou, J.D.; Gilpin, M. & Foose, T.J. (eds.) *Population Management for Survival and Recovery*. Columbia University Press, New York.
- Lapitz, Fernando (2010) Cardenal Amarillo – *Gubernatrix cristata*. Achará Digital. Año 1 (1) pág. 8. AVES URUGUAY, Montevideo.
- Li, W. H. & Graur, D. (2000) *Fundamentals of molecular evolution*. 2<sup>nd</sup> ed. Sinauer Associates, Sunderland.
- Loydi, Alejandro (2008) Situación actual del comercio ilegal de aves en la ciudad de Bahía Blanca. *BioScriba*, **1**,(1):9-16.

- Matteucci, S.D.; Herrera, P.; Miñarro, F.; Adámoli, J.; Torrella, S. & Ginzburg, R. (2007) Herramientas de toma de decisiones en la zonificación para el uso sustentable en los humedales del sudeste de la región chaqueña. *In: Memorias XII Conferencia Iberoamericana de Sistemas de Información Geográfica. Sociedad Iberoamericana de Sistemas de Información Geográfica (SIBSIG)-Universidad Nacional de Luján, Luján.*
- Marchiori, J. N. C. (2004) Fitogeografia do Rio Grande do Sul: *campos sulinos*. EST, Porto Alegre.
- Martins-Ferreira, C. (2007) Biogeografia e ecologia do Cardeal-Amarelo *Gubernatrix cristata* (Emberizidae) – resultados preliminares / Biogeography and ecology of Yellow Cardinal *Gubernatrix cristata* (Emberizidae) – preliminary results. *In: Congresso Brasileiro de Ornitologia (15: 2007: Porto Alegre, RS) [programação e resumos]*. EDIPUCRS, Porto Alegre.
- Meyer, A. (1993) Evolution of mitochondrial DNA in fishes. *In: Hochachka & Mommsen (eds). Biochemistry and molecular biology of fishes, vol.2. Elsevier Science Publishers.*
- Muniz-pereira, L.C.; Vieira, F.M. & Luque, J.L. (2009) Checklist of helminth parasites of threatened vertebrate species from Brazil. *Zootaxa*, **2123**, 1-45.
- Oyler-McCance, S. J.; St. John, J.; Knopf, F. L. & Quinn, T. W. (2005) Population genetic analysis of mountain plover using mitochondrial DNA sequence data. *Condor*, **107**, 353-362.
- Pautasso, Andres A. (2003) Aprovechamiento de la fauna Silvestre por pobladores rurales en la fraccion norte de los bajos submeridionales da la provincial de Santa Fe, Argentina. *Com. Mus. Prov. Cs. Naturales*, **8**,(2):1-61
- Pereira, G. A. & Brito, M. T. (2005) Diversidade de aves silvestres brasileiras comercializadas nas feiras livres da Região Metropolitana do Recife, Pernambuco. *Atualidades Ornitológicas*, 126 (jul/ago): 14-20. (Available at: <http://www.ao.com.br/ao126.htm>)
- Pessino, M. & Tittarelli, R. F. (2006) Cardenal Amarillo (*Gubernatrix cristata*): diagnóstico de su situación con respecto al comercio ilegal en la provincia de La Pampa, Argentina. *Gestión Ambiental*, **12**,69-76.
- Phillips, S. J.; Anderson, R. P. and Schapire, R. E. (2006) Maximum entropy modeling of species geographic distributions. *Ecological Modelling*, **190**, 231-259.
- Rojas, M. (1995) The species problem and conservation: what are we protecting? *In: Genes, population, species: readings from conservation biology. Blackwell Science/Society for Conservation Biology, Cambridge.*
- Shaffer, M.L. (1987) Minimum viable populations: coping with uncertainty. p.69-86. *In: Soulé, M.E. (ed.) Viable populations for conservation. Cambridge University Press, Cambridge.*
- Scherer, J.F.M.; Scherer, A.L. & Petry, M.V. (2010) Estrutura trófica e ocupação de hábitat da avifauna de um parque urbano em Porto Alegre, Rio Grande do Sul, Brasil. *Biotemas*, **23**,(1):169-180.

- Seewald, P. A. & Perez, C. H. F. (2009) Aves nuevas o poco frecuentes del alto valle de río negro, Río Negro, Argentina. *Nuestras Aves*, **54**,25-27.
- Shields, W.M. (1994) The Natural and Unnatural History of Inbreeding and Outbreeding. *In*: Thornhill, N.W. (ed.) The Natural History of Inbreeding and Outbreeding. University of Chicago Press, Chicago.
- Sick, H. (1997) *Ornitologia brasileira*. Nova Fronteira, Rio de Janeiro.
- Sorenson, M. D. (2003) *Avian mtDNA primers*. Boston University. Available at <http://people.bu.edu/msoren/Bird.mt.Primers.pdf>, accessed 01 July 2008.
- Sutherland, W. J.; Newton, I. & Green, R. E. (2004) Bird ecology and conservation: *a handbook of techniques*. Oxford University Press, Oxford.
- Wetmore, A. (1926) *Observations on the Birds of Argentina, Paraguay, Uruguay, and Chile*. Government Printing Office, Washington.



## 7. ANEXOS

### 7.1. Appendix A.1. *Gubernatrix cristata* samples used in this study.

Sample #	Sample Code	Sex	Country	Locality	Date	Source	Tissue type	mtDNA Samples	SSRs Samples	Haplotypes
1	PCA 002	F	ND	Canoas Zoo	12. mai. 2006	Police Seizure	Feathers/Blood	mtDNA	SSR	Hap_1
2	PCA 003	M	ND	Canoas Zoo	12. mai. 2006	Police Seizure	Feathers/Blood	mtDNA	SSR	Hap_3
3	PCA 004	M	ND	Canoas Zoo	23. set. 2006	Police Seizure	Feathers/Blood	mtDNA	SSR	Hap_3
4	PCA 005	M	ND	Canoas Zoo	23. set. 2006	Police Seizure	Feathers/Blood	mtDNA	SSR	Hap_1
5	PCA 006	M	ND	Canoas Zoo	23. set. 2006	Police Seizure	Feathers/Blood	mtDNA	SSR	Hap_3
6	PCA 007	M	ND	Canoas Zoo	23. set. 2006	Police Seizure	Feathers/Blood	mtDNA	SSR	Hap_3
7	PCA 008	F	ND	Canoas Zoo	23. set. 2006	Police Seizure	Feathers/Blood	mtDNA	SSR	Hap_3
8	PCA 009	F	ND	Canoas Zoo	23. set. 2006	Police Seizure	Feathers/Blood	mtDNA	SSR	Hap_3
9	PCA 010	M	ND	Canoas Zoo	5. out. 2006	Police Seizure	Feathers/Blood	mtDNA	SSR	Hap_1
10	PCA 011	F	ND	Canoas Zoo	7. abr. 2007	Police Seizure	Feathers/Blood	mtDNA	SSR	Hap_3
11	PCA 012	?	ND	Canoas Zoo	7. abr. 2007	Police Seizure	Feathers/Blood	mtDNA	SSR	Hap_3
12	PCA 051	F	ND	Canoas Zoo	7. abr. 2007	Police Seizure	Feathers/Blood	mtDNA	SSR	Hap_3
13	PCA 052	M	ND	Canoas Zoo	22. jun. 2007	Police Seizure	Feathers/Blood	mtDNA	SSR	Hap_1
14	PCA 053	F	ND	Canoas Zoo	22. jun. 2007	Police Seizure	Feathers/Blood	mtDNA	SSR	Hap_3
15	PCA 054	M	ND	Canoas Zoo	22. jun. 2007	Police Seizure	Feathers/Blood	mtDNA	SSR	Hap_3
16	PCA 055	?	ND	Canoas Zoo	22. jun. 2007	Police Seizure	Feathers/Blood	mtDNA	SSR	Hap_1
17	PCA 056	?	ND	Canoas Zoo	22. jun. 2007	Police Seizure	Feathers/Blood	mtDNA	SSR	Hap_1
18	001/02	?	Uruguay	Durazno	8. mai. 2008	Captive Bred	Blood	mtDNA	SSR	Hap_3
19	002/02	?	Uruguay	Durazno	8. mai. 2008	Captive Bred	Blood	mtDNA	SSR	Hap_1
20	003/02	?	Uruguay	Durazno	8. mai. 2008	Captive Bred	Blood	mtDNA	SSR	Hap_3
21	004/02	?	Uruguay	Durazno	8. mai. 2008	Captive Bred	Blood	mtDNA	SSR	Hap_1
22	005/02	?	Uruguay	Durazno	8. mai. 2008	Captive Bred	Blood	mtDNA	SSR	Hap_3
23	006/02	?	Uruguay	Durazno	8. mai. 2008	Captive Bred	Blood	mtDNA	SSR	Hap_3
24	007/02	?	Uruguay	Durazno	8. mai. 2008	Captive Bred	Blood	mtDNA	SSR	Hap_3
25	0800/04	?	Uruguay	Colonia	8. mai. 2008	Captive Bred	Blood	mtDNA	SSR	Hap_3
26	932/06	?	Uruguay	Durazno	8. mai. 2008	Captive Bred	Blood	mtDNA	SSR	Hap_3
27	453/07	?	Uruguay	Durazno	8. mai. 2008	Captive Bred	Blood	mtDNA	SSR	Hap_3
28	458/07	?	Uruguay	Durazno	8. mai. 2008	Captive Bred	Blood	mtDNA	SSR	Hap_3
29	459/07	?	Uruguay	Durazno	8. mai. 2008	Captive Bred	Blood	mtDNA	SSR	Hap_3

## 7.1. Appendix A.1. *Gubernatrix cristata* samples used in this study.

Sample #	Sample Code	Sex	Country	Locality	Date	Source	Tissue type	mtDNA Samples	SSRs Samples	Haplotypes
30	G 63205	M	Uruguay	Lavalleja	22. jan. 2007	Wild	Feathers/Blood	mtDNA	SSR	Hap_1
31	G 63206	?	Uruguay	Lavalleja	22. jan. 2007	Wild	Feathers/Blood	mtDNA	SSR	Hap_3
32	G 63207	?	Uruguay	Lavalleja	22. jan. 2007	Wild	Feathers/Blood	mtDNA	SSR	Hap_3
33	G 63209	F	Brasil	PEE/RS	24. mai. 2008	Wild	Blood	mtDNA	SSR	Hap_3
34	G 63210	F	Brasil	PEE/RS	24. mai. 2008	Wild	Blood	mtDNA	SSR	Hap_3
35	BRIAO 001	M	Brasil	Pedras Altas/RS	19. abr. 2006	Captive Bred	Feathers	mtDNA	SSR	Hap_1
36	BRIAO 002	M	Brasil	Bagé/RS	19. abr. 2006	Captive Bred	Feathers	No	SSR	-
37	PCA 001	M	ND	Canoas Zoo	12. mai. 2006	Police Seizure	Feathers	mtDNA	SSR	Hap_1
38	190907	?	ND	Canoas Zoo	19. set. 2007	Police Seizure	Feathers	mtDNA	SSR	Hap_1
39	URU 001	?	ND	Private Owner	30. jun. 1905	Captive Bred	Feathers	mtDNA	SSR	Hap_1
40	URU 002	?	ND	Private Owner	30. jun. 1905	Captive Bred	Feathers	mtDNA	SSR	Hap_1
41	URU 003	?	ND	Private Owner	30. jun. 1905	Captive Bred	Feathers	mtDNA	SSR	Hap_1
42	URU 004	?	ND	Private Owner	30. jun. 1905	Captive Bred	Feathers	mtDNA	SSR	Hap_1
43	MACN 384a	M	Argentina	Tucumán	7. abr. 1905	Museum Skin	Footpads	mtDNA	SSR	Hap_3
44	MACN 826a	M	Argentina	San Luis	15. dez. 1925	Museum Skin	Footpads	No	SSR	-
45	MACN 1462a	M	Uruguay	Colonia	15. abr. 1927	Museum Skin	Footpads	mtDNA	SSR	Hap_2
46	MACN 3388a	M	Argentina	Cordoba	13. mai. 1933	Museum Skin	Footpads	mtDNA	No	Hap_2
47	MACN 43559	M	Argentina	Entre Ríos	21. abr. 1961	Museum Skin	Footpads	mtDNA	SSR	Hap_3
48	MACN 48179	M	Argentina	Corrientes	26. dez. 1962	Museum Skin	Footpads	mtDNA	SSR	Hap_3
49	MACN 49697	M	Argentina	La Pampa	6. jun. 1963	Museum Skin	Footpads	mtDNA	SSR	Hap_1
50	MACN 51721	M	Argentina	Buenos Aires	18. abr. 1967	Museum Skin	Footpads	mtDNA	SSR	Hap_1
51	MZUSP 70522	M	Brasil	Arroio Xuí/RS	20. jul. 1966	Museum Skin	Footpads	No	SSR	-
52	MCN 01682	M	Brasil	Arroio Quaraí-chico/RS	12. jul. 1973	Museum Skin	Footpads	No	SSR	-
53	MCN 01683	M	Brasil	Pedras Altas/RS	25. mar. 1975	Museum Skin	Footpads	No	SSR	-
54	ROM 59090	M	Argentina	Cordoba	2. jun. 1913	Museum Skin	Skin	mtDNA	SSR	Hap_1
55	ROM 22.1.2.4	M	ND	Metro Toronto Zoo	?	Museum Skin	Skin	mtDNA	SSR	Hap_3
56	P 2151	?	ND	Private Owner	.2006	Police Seizure	Muscle	mtDNA	SSR	Hap_3
58	PCA014	M	ND	Gramado Zoo	24. abr. 2009	Police Seizure	Feathers/Blood	mtDNA	No	Hap_2
59	PCA015	M	ND	Gramado Zoo	24. abr. 2009	Police Seizure	Feathers/Blood	mtDNA	No	Hap_2
60	PCA016	M	ND	Gramado Zoo	24. abr. 2009	Police Seizure	Feathers/Blood	mtDNA	No	Hap_1
63	PCA057	F	ND	Gramado Zoo	24. abr. 2009	Police Seizure	Feathers/Blood	mtDNA	No	Hap_2
64	PCA058	F	ND	Gramado Zoo	24. abr. 2009	Police Seizure	Feathers/Blood	mtDNA	No	Hap_2

**7.1. Appendix A.1.** *Gubernatrix cristata* samples used in this study.

Sample #	Sample Code	Sex	Country	Locality	Date	Source	Tissue type	mtDNA Samples	SSRs Samples	Haplotypes
65	PCA059	F	ND	Gramado Zoo	24. abr. 2009	Police Seizure	Feathers/Blood	mtDNA	No	Hap_2
67	PCA061	M	ND	Gramado Zoo	24. abr. 2009	Police Seizure	Feathers/Blood	mtDNA	No	Hap_2
75	DA32105	F	Brasil	CITOCEL	3. abr. 2008	Captive Bred	DNA	mtDNA	No	Hap_3
77	DA32107	M	Brasil	CITOCEL	3. abr. 2008	Captive Bred	DNA	mtDNA	No	Hap_1
78	DA32108	M	Brasil	CITOCEL	3. abr. 2008	Captive Bred	DNA	mtDNA	No	Hap_1
79	DA32109	F	Brasil	CITOCEL	3. abr. 2008	Captive Bred	DNA	mtDNA	No	Hap_1
84	DA32387	F	Brasil	CITOCEL	20. mai. 2008	Captive Bred	DNA	mtDNA	No	Hap_3
87	DA33531	M	Brasil	CITOCEL	18. dez. 2008	Captive Bred	DNA	mtDNA	No	Hap_3
88	DA33532	M	Brasil	CITOCEL	18. dez. 2008	Captive Bred	DNA	mtDNA	No	Hap_2
89	DA33793	F	Brasil	CITOCEL	20. jan. 2009	Captive Bred	DNA	mtDNA	No	Hap_2
90	DA33794	F	Brasil	CITOCEL	20. jan. 2009	Captive Bred	DNA	mtDNA	No	Hap_2

## 7.2. Author guidelines. Animal Conservation and Diversity and Distributions

### Animal Conservation

The rapid publication journal for quantitative studies in conservation

Published on behalf of the Zoological Society of London

Edited by:

Trenton Garner, Matthew Gompper, Iain Gordon, Todd Katzner, Karen Mock and Stephen Redpath  
Reviews Editor: Nathalie Pettorelli

Print ISSN: 1367-9430

Online ISSN: 1469-1795

Frequency: Bi-monthly

Current Volume: 13 / 2010

ISI Journal Citation Reports® Ranking: 2009: Ecology: 49 / 127; Biodiversity Conservation: 9 / 28

Impact Factor: 2.358

### Author Guidelines

#### Instructions to Authors

Papers should be submitted online at <http://mc.manuscriptcentral.com/acv> Full upload instructions and support are available online from the submission site via the 'Get Help Now' button. Please submit your covering letter or comments to the editor when prompted online.

The submissions listed below are considered:

Research papers: must be limited to 4000 words, excluding references, tables and figures

Reviews: Review articles should provide a succinct analysis of the subject and can include quantitative meta-analysis, syntheses and modeling approaches, and critical evaluation. Prospects for future research directions should be described. Reviews should be no more than 4000 words with a maximum of 80 references, 70% of which must have been published within 5 years of the submission. Extensive literature reviews will not be considered.

Prospective authors should submit a 300-word abstract, a list of up to 20 key references, and a cover letter (two pages maximum) outlining what will be discussed in the article, to the Reviews Editor (Nathalie.pettorelli@ioz.ac.uk). The editor will supply guidelines on manuscript preparation if the proposal is accepted. The submission of completed manuscripts without prior consultation with the Editor is strongly discouraged. Authors should note that all articles in Animal Conservation are peer-reviewed and publication cannot be guaranteed.

Correspondence: criticisms or additions to papers already published; limited to 1000 words.

Papers that report experimental work must comply with the standards and procedures laid down by British national or equivalent legislation. Attention is drawn to the 'Guidelines for the Use of Animals in Research' published in each January issue of the journal Animal Behaviour. Papers will not be accepted if they are based on work involving cruelty to animals or if the work may have put at risk endangered populations, species or habitats. The Editors may seek advice from the Ethical Committee of the Zoological Society of London on ethical matters.

#### Conflict of interest

Authors must declare details of any potential conflict of interest. A conflict of interest exists when professional judgement concerning a primary interest (such as animal welfare or the validity of research) may be influenced by secondary interests (personal matters such as financial gain, personal relationships or professional rivalry).

#### Submission

Submission of a manuscript will be taken to imply that the material is original and that no similar paper is being, or will be, submitted elsewhere, either in whole or substantial part. Serialized studies should not be submitted and titles should not contain part numbers. The Editors reserve the right to accelerate the publication of high-profile papers, commissioned reviews or topical papers. Authors are required to provide the names of 2 - 4 suitable referees, however the Editor reserves the right to choose referees other than those suggested.

#### OnlineOpen

Authors of articles in the Animal Conservation Journal can now choose to make their articles open access and available free online for all readers through the payment of an author fee. A copy of the OnlineOpen form can be found [here](#).

#### DNA sequence data

DNA sequence data published in Animal Conservation should be deposited in one of the EMBL/GenBank/DDBJ nucleotide sequence databases, and accession numbers must be included in the final version of the manuscript. Genetic data from sources other than the manuscript must either be publicly available (with accession numbers or citations provided) or the authors must have appropriate permission to use unpublished data. Such permission should be described explicitly in the covering letter

#### Presentation

Typescripts must be typed in double spacing, and pages should be numbered consecutively, including those containing acknowledgements, references, tables and figures. Lines must be numbered, preferably within pages.

Manuscripts for review must consist of no more than two files and should, ideally be a single file with figures embedded in the text (please note that separate high resolution figure files will be required upon acceptance - please see below). Typescripts must be in English (both English and American English are acceptable).

The Editors reserve the right to modify accepted manuscripts that do not conform to scientific, technical, stylistic or grammatical standards, and these minor alterations may not be seen by the authors until the proof stage.

#### Conventions

The Metric system must be used and SI units where appropriate. For further details see Baron, D.N. (1988). Units, symbols and abbreviations. 5th edition. London: Royal Society of Medicine Series. Whole numbers one to nine should be spelled out and number 10 onwards given in numerals. If a new taxon is described, the institution in which the type material is deposited must be given, together with details of the registration assigned to it. Full binomial names should be given on the first occasion an organism is mentioned (and abbreviated thereafter), except at the beginning of a sentence. Avoid footnotes except to add information below the body of a table. Do not use initial capitals for the common names of animals unless derived from a proper noun.

#### Typescript

The typescript should follow the conventional form and must include:

- (1) Title page giving a concise title (do not include scientific names in the title), followed by a list of authors' names and the institutions where the work was carried out. The name, address and email address of the corresponding author should also be given. A short title for page headings must be provided (maximum 8 words).
- (2) Abstract of not more than 300 words which should list the main results and conclusions. The abstract should also explain the importance of the paper in a way that is accessible to non-specialists and should describe the novel aspects of the research and highlight the relevance of the findings to other taxa or general principles in conservation biology. Authors may submit non-English abstracts for online publication to allow the international research community greater access to published articles. Translated abstracts should be submitted in pdf format as supplementary material. The Editors have no input into the content of supplementary material, therefore accuracy is the sole responsibility of the authors.
- (3) Keywords. A maximum of eight keywords may be suggested.
- (4) Introduction, which should not provide a review of the area of work but should introduce the reader to the aims and context for the work described.
- (5) Materials and Methods should be sufficient to allow the work to be replicated, but should not repeat information described fully elsewhere.
- (6) Results should be restricted to a factual account of the findings obtained and the text must not duplicate information given in Tables and Figures. (7) Discussion. This should point out the significance of the results in relation to the reasons for undertaking the research, and describe the novel aspects of the research and the relevance of the findings to a range of taxa or general principles in conservation biology.

Please note that appendices are no longer published in the printed version of the journal. Supplementary material may be published online only. References Accuracy of references is the responsibility of the author(s).

#### References

must be checked against the text to ensure (a) that the spelling of authors' names and the dates given are consistent and (b) that all authors quoted in the text (in date order if more than one) are given in the reference list and vice versa. The full title of the paper must be given together with the first and last pages.

Journal titles should be abbreviated in accordance with the Zoological Record Serial Sources, published annually by BIOSIS.

Book titles should be followed by the place of publication and the publisher. Please give the name of the editor(s) if different from the author cited.

In the text, references must be arranged chronologically with the surname(s) of the author(s) followed by the date. Use a, b, etc. after the year to distinguish papers published by the same author(s) in the same year. Reference should not be made to unpublished data.

(i) Two authors: use both names and the year. Do not use et al.

(ii) Three authors: on first citation use all authors' names and the year. Thereafter it is usually sufficient to give the name of the first author followed by et al. and the date.

(iii) More than three authors: on first citation and thereafter give the name of the first author followed by et al. and the date.

In the list, references must be arranged first alphabetically under author(s) name(s) and then in chronological order if several papers by the same author(s) are cited.

#### Examples

Lemelin, P. (1996a). Relationships between hand morphology and feeding strategies in small-bodied prosimians. *Am. J. phys. Anthropol. (Suppl.)* 22, 148.

Lemelin, P. (1996b). The evolution of manual prehensility in primates: a comparative study of prosimians and didelphid marsupials. PhD thesis, State University of New York at Stony Brook.

Pianka, E. R. (1978). *Evolutionary ecology*. 2nd edn. New York: Harper & Row.

Whitear, M. (1992). Solitary chemosensory cells. In *Fish chemoreception: 103-125*. Hara, T. J. (Ed.). London: Chapman & Hall.

#### References in Articles

We recommend the use of a tool such as EndNote or Reference Manager for reference management and formatting. EndNote reference styles can be searched for here: <http://www.endnote.com/support/enstyles.asp>

Reference Manager reference styles can be searched for here: <http://www.refman.com/support/rmstyles.asp>

#### Figures

Illustrations may be line drawings or photographs and should be numbered consecutively in the text as Fig. 1, Fig. 2 etc. Component parts of figures should be labelled (a), (b), (c) etc. Captions for figures, which should be self-explanatory, must be typed, double spaced, on a separate page and must not contain details of results.

Our preferred electronic file type is vector-format encapsulated post script (EPS) because these images are scaleable and therefore do not lose quality in the online PDF. All line drawings or photographs with added labelling should be supplied in EPS format. Half tones without any labelling should be supplied in TIFF format at 300 dots per inch minimum. If line drawings cannot be supplied as EPS files then they must be in TIFF format with a minimum resolution of 800 dpi. These resolutions also apply to any images embedded into an EPS file.

Line drawings should not be larger than twice the final size and in no circumstances should exceed 168 x 220 mm. The axes of graphs should be carefully chosen so as to occupy the space available to the best advantage. When reduced, the drawing should fit into either one (80 mm) or two (168 mm) columns, preferably the former.

Lines should be bold enough to stand reduction to about 0.25-0.35 mm. Line drawings should be as simple as possible and many computer-generated figures, such as 3-dimensional graphs, fine lines, gradations of stippling and unusual symbols, cannot be reproduced satisfactorily when reduced. Unsatisfactory line drawings will have to be redrawn at the author's expense. Preferred symbols are open and filled circles, boxes and triangles, and these should be used consistently. Lettering should be kept to a minimum and should be self-explanatory and unambiguous and of sufficiently high quality and size to be clearly visible after reduction to final size. Lettering of all figures within the manuscript should be of uniform style in a sans serif typeface (Helvetica) and capitals should be used for the initial letter of the first word only. Bold lettering should not be used.

Photographs should be the same size as they will appear in the journal and should be selected to fit neatly into one column (80 mm) or two columns (168 mm). Photographs should be labelled and numbered as for line drawings. For microscopical preparations, scale bars with appropriate units must be provided; statements of magnification are not acceptable.

Colour figures may be accepted provided that they are of a very high quality. The cost of reproduction must be met by the author(s) and a binding agreement to meet the costs will be required before the manuscript can be accepted for publication. For colour figures, the instructions for the preparation of photographs should be followed. Original illustrations should not be sent until the paper has been accepted and will only be returned on request. Any article received by Blackwell Publishing with colour work will not be published until the colour work agreement form has been returned.

#### Tables

These must fit the page size (220 x 168 mm) without undue reduction. Oversize tables will not be accepted. Tables are referred to as Table 1, Table 2, etc., and any sub-sections as (a), (b), etc. Footnotes in tables should be indicated by superscript a, b.

#### Decisions and invitations to revise

All submissions are subject to peer review and authors can expect a decision, or an explanation for the delay, within 3 months of receipt. If a revision is requested, the corresponding author should submit the revised manuscript within 2 months unless there are special reasons for a delay, agreed in advance with the Editor. Papers not received within 2 months may be treated as new submissions and sent for further evaluation by new referees.

#### Pre-submission English-language editing

Authors for whom English is a second language may choose to have their manuscript professionally edited before submission to improve the English. A list of independent suppliers of editing services can be found at [www.blackwellpublishing.com/bauthor/english\\_language.asp](http://www.blackwellpublishing.com/bauthor/english_language.asp). All services are paid for and arranged by the author, and use of one of these services does not guarantee acceptance or preference for publication.

#### Exclusive Licence Form

Authors will be required to sign an Exclusive Licence Form (ELF) for all papers accepted for publication. Signature of the ELF is a condition of publication and papers will not be passed to the publisher for production unless a signed form has been received. Please note that signature of the ELF does not affect ownership of copyright in the material. (Government employees need to complete the Author Warranty sections, although a licence does not need to be assigned in such cases). After submission authors will retain the right to publish their paper in various medium/circumstances (please see the form for further details). To assist authors an appropriate form will be supplied by the editorial office.

## Diversity and Distributions

A Journal of Conservation Biogeography

Edited by:

David M. Richardson

Print ISSN: 1366-9516

Online ISSN: 1472-4642

Frequency: Bi-monthly

Current Volume: 16 / 2010

ISI Journal Citation Reports® Ranking: 2009: Ecology: 18 / 127; Biodiversity Conservation: 4 / 28

Impact Factor: 4.224

### Author Guidelines

Average time between receipt of manuscript and first editorial decision is now 43 days!

Diversity and Distributions has five main categories of articles: -

1. Biodiversions. These are editorial items solicited directly by the Editor. Unsolicited material will not normally be considered. If you have an idea for such a contribution (up to 2000 words), please contact the Editor, who will provide you with the necessary guidance for submission.
2. Biodiversity Viewpoints. This section contains short essays (usually up to 3000 words) considering biodiversity from a particular disciplinary, regional, political, or other standpoint. If you would like to contribute such an essay, please contact the Editor outlining the distinctive character of your proposed essay, its length, the number of references, and the character of any illustrations to be used.
3. Biodiversity Research and Reviews This is the core section of the journal and presents research or review articles up to 5000 words in length, but preferably shorter (the word limit refers to text from the start of the introduction to the end of the acknowledgements - i.e. excluding the title, abstract, references, figure captions, and tables). Tables should not be overlong and complicated. The Editor reserves the right to publish long tables and appendices on the journal's website, rather than in the printed version. Such a decision will only be taken after consultation with the author. A short running title should be provided. The manuscript must include an abstract of no more than 300 words structured under the headings: Aim, Location, Methods, Results, Main conclusions, and ending with a list of 6-10 keywords or phrases, arranged in alphabetical order. Three different weights of headings are available: authors should indicate the relative importance of a heading by the use of ringed capital letters. i.e. (A) for main headings; (B) for secondary headings; and, (C) for tertiary headings. The correct nomenclatural authorities for all taxa must be given on the first appearance in the text, in Tables, and in the captions to Figures, unless a general reference to a standard source is provided at an appropriate place in the manuscript. A biosketch entry should be included after the references section (see below).
4. Biodiversity Letters. This section presents short items (normally less than 1000 words) of general news interest with respect to biodiversity, conferences or events, computer hardware and software developments, films and videos, the law, and political debates. Brief letters to the editor are also most welcome. Lively titles are encouraged, and material should be as topical as possible. Longer letters (up to 2500 words in total) prompted by papers previously published in this or occasionally other journals are also encouraged. Such longer communications should include a one-paragraph abstract (150 word maximum), and a list of 6-10 keywords.

### Manuscript preparation and submission

Diversity and Distributions requires online submission of manuscripts at <http://mc.manuscriptcentral.com/ddi>. Submission online is an intuitive, step-by-step process. By submitting online, you will benefit from quicker peer-review, web-based manuscript tracking, online reviewing and faster response. You will need your manuscript and figures in a digital format. When submitting, authors should upload a single file that contains all text (including a short running title, references, tables, figure captions and appendices) and figures, which should be embedded into the document. A PDF file will then automatically be created for reviewing purposes. Full instructions and support for authors can be found at the Site. To use the Site you will need a user ID and password. Go to the Journal's submission homepage (<http://mc.manuscriptcentral.com/ddi>) and click 'Create a new account' if you have not registered before, or click 'Check for existing account' if you have submitted online or reviewed online before for the Journal (or if you have forgotten your details). If you at any time experience difficulty with your online submission, please contact the Editorial Assistant at [ddi@sun.ac.za](mailto:ddi@sun.ac.za).



Contributing authors are requested to submit, at the time of submission of their manuscripts, a list of at least five persons that they consider well qualified to review the submitted work (e-mail addresses should be included). The list should NOT include any current or recent collaborators in work that is closely related to the topic of the submitted paper, or any persons within the same organization as any of the authors of the submitted work.

All enquiries should be directed to:

Prof. David M. Richardson  
Centre for Invasion Biology (CIB)  
Science Faculty  
University of Stellenbosch  
Private Bag X1  
Matieland 7602  
South Africa

Tel: +2782 902-9024  
Fax: +2721 808-2995  
E-mail: rich@sun.ac.za  
CIB website: www.sun.ac.za/cib

Pages and lines should be numbered to aid cross-referencing (in MSWord, go to "Page Setup" then "Layout"; select "Line numbers"; click on "Add line numbers" and select "continuous").

Only papers written in English will be accepted. The journal cannot provide detailed editing of manuscripts to correct English. Where necessary, authors should have their manuscripts checked by a native English speaker before submitting their work.

#### Abbreviations and units

SI units (metre, kilogram, etc.) are essential. Statistics and measurements should be given in figures, i.e. 10<sup>+</sup>mm, except where the number begins the paragraph. When the number does not refer to a unit of measurement, it is spelt out, except where the number is greater than 10. A list of preferred abbreviations and naming conventions is available here.

#### Tables

Tables must be positioned on separate sheets and numbered consecutively (Table 1, Table 2, etc.). Column headings should be brief: with units of measurement in parentheses. Tables should be typed as text, using 'tabs' (not spaces) to align columns. The use of table editors should be avoided. Do not use graphics software to create tables.

#### Methods

Please ensure that this section is entitled 'METHODS', and not 'MATERIALS AND METHODS'.

#### Figures, Illustrations and Maps

All illustrations (including photographs) are classified as figures and should be numbered consecutively (Fig. 1, Fig. 2, etc.). When submitting a manuscript to Manuscript Central, authors should upload a single text file with embedded figures. Upon your manuscript being accepted for publication, please supply separate files containing electronic versions of your figures (see File Formats, below). Please note that your paper will go through production more quickly if instructions on content and format are followed carefully. Each figure must have a legend that makes the material completely understandable. Legends should be presented separately from the figures, in a list at the end of the manuscript. Label multi-panel figures (a), (b), (c), etc., preferably in the upper left corner, and refer to them in the text as, for example, Fig. 1(a). Please ensure that electronic artwork is prepared such that, after reduction to fit across one or two columns or two-thirds width (80 mm, 169 mm or 110 mm, respectively) as required, all lettering and symbols will be clear and easy to read, i.e. no labels should be too large or too small. Avoid using tints if possible; if they are essential to the understanding of the figure, try to make them coarse.

Maps that display area data and organism distribution at a continental, hemispheric, or world scale must always use an equal-area map projection (e.g. Mollweide or Aitoff's). Note especially that Mercator's projection is not acceptable for such data. Please indicate the precise projection employed in the caption. On these maps, the equatorial scale should be indicated, while scale information should be provided, preferably as a scale bar within the figure, for all maps of whatever size and area.

File Formats: After acceptance of your manuscript for publication, figure files should be supplied as follows.

Photographic figures should be saved in tif format at 300 d.p.i. (or failing that in jpg format with low compression).

Line figures should be saved as vector graphics (i.e. composed of lines, curves, points and fonts; not pixels) in eps or pdf format, or embedded as such in Word, as this enhances their display when published online. Combination

figures (those composed of vector and pixel/raster elements) should also be saved in eps or pdf format where possible (or embedded as such in Word). If line figures and combination figures cannot be saved in vector graphics format, they should be saved in tif format at high resolution (i.e. 600 d.p.i.) (do not save them in jpg format). If you are unsure about the resolution of your tif files, please zoom in and check that fonts, curves and diagonal lines are smooth-edged and do not appear blocky when viewed at high magnification. Note that line and combination figures supplied in tif format are downsampled for online publication and so authors should preferentially opt for vector graphic formats for these figure types (full resolution tif files are used for print publication).

If there is colour artwork in your manuscript when it is accepted for publication, Wiley-Blackwell require you to complete and return a Colourwork Agreement Form before your paper can be published. Once completed, please return the form to the Production Editor at the address below:

Production Editor  
Diversity and Distributions  
Wiley-Blackwell  
600 North Bridge Road #05-01  
Parkview Square  
Singapore 188778.  
Fax: +65 6295 6202  
E-mail: ddi@wiley.com

Any article received by Wiley-Blackwell with colour work will not be published until the form has been returned. Under exceptional circumstances, authors may request the above charges to be waived. This must be done, in writing, at the time of submission of the manuscript, and authors must justify to the Editor that inclusion of the figure(s) in colour is essential for interpretation of the results presented. If authors wish to apply for funds to cover the costs of colour printing, the Editor will provide relevant support letters to funding bodies, indicating acceptance of the paper. Note that we offer a free Colour on the Web option whereby authors can have figures printed in black and white in the journal but in colour in the online version, free of charge.

#### Appendices and Supporting Information

Appendices may be provided for important primary data, which needs to be included in the paper. If, however, these data are very extensive, or if they are of only indirect relevance to the paper, they will normally be made available in an electronic form through the Journal's web pages. Mention of the first supporting appendix, table or figure ,etc., in the text should be in the form 'see Appendix S1 in Supporting Information' [where 'S' indicates Supporting], subsequent mention should be in the form 'see Appendix S2'. Authors should then include a Supporting Information section after the References section, which should be in the following form (text in curly brackets is for completion by the author, see instructions below):

-----  
Supporting Information

Additional Supporting Information may be found in the online version of this article:

Appendix S1 {Insert short legend to online Appendix S1}  
Figure S1 {Insert short legend to online Figure S1}  
Table S1 {Insert short legend to online Table S1}

As a service to our authors and readers, this journal provides supporting information supplied by the authors. Such materials are peer-reviewed and may be re-organized for online delivery, but are not copy-edited or typeset. Technical support issues arising from supporting information (other than missing files) should be addressed to the authors.

-----  
For reasons of space, only short titles to Supporting Information should be given in this section; full titles (if different) can be given with the Supporting Information itself; full titles can include a fuller description of content, definition of abbreviations, etc. Supporting Information files are hosted by the Publisher in the format supplied by the author and are not copy-edited by the Publisher. It is the responsibility of the author to supply Supporting Information in an appropriate file format and to ensure that it is accurate and correct. Authors should therefore prepare Supporting Information with the same rigour as their main paper, including adherence to journal style (e.g. formatting of references). Supporting Information can be provided as separate files or as one combined file. Authors are discouraged from supplying very large files or files in non-standard file formats, both of which may reduce their use to the readership. Files should be prepared without line numbers or wide line spacing, and with all track-change edits accepted. Further information on Supporting Information is available here.

At proof correction stage authors will be given access to their Supporting Information (via the web) and should check it for accuracy and updates. If changes are required corrected versions of the files received with the proof must be emailed to the Production Editor, with a brief description of the changes made. Supporting Information must be checked alongside the main proof and corrections for both returned to the Production Editor at the same time.

#### Biosketch/Biosketches

A short Biosketch/Biosketches entry (30-100 words for one author/150 words for the first three authors, respectively) describing the research interests of the author(s) should be provided. For papers with 4 or more authors, biosketch details should be supplied for the first author only; alternatively, a general statement of the focus of the research team (a link to a group web page is encouraged) should be provided, together with a statement of author roles, e.g. Author contributions: A.S. and K.J. conceived the ideas; K.J. and R.L.M. collected the data; R.L.M. and P.A.K. analysed the data; and A.S. and K.J. led the writing.

#### References

We recommend the use of a tool such as EndNote for reference management and formatting. Click [here](#) to download the most up to date EndNote reference style for Diversity and Distributions. References should be made by giving the author's name with the year of publication in parentheses. When reference is made to a work by three authors or more, only the first name and et al. should be given in the citation. All authors' names should be listed in the reference itself. If several papers by the same author and from the same year are cited, a, b, c, etc., should be inserted after the year of publication. References must be listed in alphabetical order at the end of the paper in the following standard form:

Cox, C. B. & Moore, P. D. (1999) *Biogeography: an ecological and evolutionary approach*, 6th edn. Blackwell Science Ltd, Oxford.

May, R.M. (1994) The effects of spatial scale on ecological questions and answers. *Large-scale ecology and conservation biology* (ed. by P.J. Edwards, R.M. May and N.R. Webb), pp. 1-17. Blackwell Scientific Publications, Oxford.

Prentice, I.C., Guiot, J., Huntley, B., Jolly, D. & Cheddadi, R. (1996) Reconstructing biomes from palaeoecological data: a general method and its application to European pollen data at 0 and 6 ka. *Climate Dynamics*, 12, 185-194.

Please note that titles of journals should be written in full. Unpublished data, works in preparation and papers submitted but not yet accepted may be cited in the text, giving the author's initials and surname, but should not be included in the reference list.

# Livros Grátis

( <http://www.livrosgratis.com.br> )

Milhares de Livros para Download:

[Baixar livros de Administração](#)

[Baixar livros de Agronomia](#)

[Baixar livros de Arquitetura](#)

[Baixar livros de Artes](#)

[Baixar livros de Astronomia](#)

[Baixar livros de Biologia Geral](#)

[Baixar livros de Ciência da Computação](#)

[Baixar livros de Ciência da Informação](#)

[Baixar livros de Ciência Política](#)

[Baixar livros de Ciências da Saúde](#)

[Baixar livros de Comunicação](#)

[Baixar livros do Conselho Nacional de Educação - CNE](#)

[Baixar livros de Defesa civil](#)

[Baixar livros de Direito](#)

[Baixar livros de Direitos humanos](#)

[Baixar livros de Economia](#)

[Baixar livros de Economia Doméstica](#)

[Baixar livros de Educação](#)

[Baixar livros de Educação - Trânsito](#)

[Baixar livros de Educação Física](#)

[Baixar livros de Engenharia Aeroespacial](#)

[Baixar livros de Farmácia](#)

[Baixar livros de Filosofia](#)

[Baixar livros de Física](#)

[Baixar livros de Geociências](#)

[Baixar livros de Geografia](#)

[Baixar livros de História](#)

[Baixar livros de Línguas](#)

[Baixar livros de Literatura](#)  
[Baixar livros de Literatura de Cordel](#)  
[Baixar livros de Literatura Infantil](#)  
[Baixar livros de Matemática](#)  
[Baixar livros de Medicina](#)  
[Baixar livros de Medicina Veterinária](#)  
[Baixar livros de Meio Ambiente](#)  
[Baixar livros de Meteorologia](#)  
[Baixar Monografias e TCC](#)  
[Baixar livros Multidisciplinar](#)  
[Baixar livros de Música](#)  
[Baixar livros de Psicologia](#)  
[Baixar livros de Química](#)  
[Baixar livros de Saúde Coletiva](#)  
[Baixar livros de Serviço Social](#)  
[Baixar livros de Sociologia](#)  
[Baixar livros de Teologia](#)  
[Baixar livros de Trabalho](#)  
[Baixar livros de Turismo](#)